

1-1-2008

Antioxidant and Vasorelaxant Activities of Flavonoids from *Amygdalus lycioides* var. *horrida*

HOSSEIN BABAEI

OMID SADEGHPOUR

LUTFUN NAHAR

ABBAS DELAZAR

HOSSEIN NAZEMIYEH

See next page for additional authors

Follow this and additional works at: <https://journals.tubitak.gov.tr/biology>



Part of the [Biology Commons](#)

Recommended Citation

BABAEI, HOSSEIN; SADEGHPOUR, OMID; NAHAR, LUTFUN; DELAZAR, ABBAS; NAZEMIYEH, HOSSEIN; MANSOURI, MOHAMMAD REZA; POURSAEID, NASER; ASNAASHARI, SOLMAZ; MOGHADAM, SEDIGHEH BAMDAD; and SARKER, SATYAJIT DEY (2008) "Antioxidant and Vasorelaxant Activities of Flavonoids from *Amygdalus lycioides* var. *horrida*," *Turkish Journal of Biology*. Vol. 32: No. 3, Article 9. Available at: <https://journals.tubitak.gov.tr/biology/vol32/iss3/9>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Biology by an authorized editor of TÜBİTAK Academic Journals. For more information, please contact academic.publications@tubitak.gov.tr.

Antioxidant and Vasorelaxant Activities of Flavonoids from *Amygdalus lycioides* var. *horrida*

Authors

HOSSEIN BABAEI, OMID SADEGHPOUR, LUTFUN NAHAR, ABBAS DELAZAR, HOSSEIN NAZEMIYEH, MOHAMMAD REZA MANSOURI, NASER POURSAEID, SOLMAZ ASNAASHARI, SEDIGHEH BAMDAD MOGHADAM, and SATYAJIT DEY SARKER

Antioxidant and Vasorelaxant Activities of Flavonoids from *Amygdalus lycioides* var. *horrida*

Hossein BABAEI¹, Omid SADEGHPOUR², Lutfun NAHAR³, Abbas DELAZAR¹, Hossein NAZEMIYEH¹,
Mohammad Reza MANSOURI⁴, Naser POURSAEID⁴, Solmaz ASNAASHARI¹, Sedigheh Bamdad MOGHADAM¹,
Satyajit Dey SARKER^{3,*}

¹Faculty of Pharmacy, Drug Applied Research Center, Tabriz University of Medical Sciences, Tabriz - IRAN

²Research Institute for Islamic and Complementary Medicine, Iran University of Medical Sciences, Tehran - IRAN

³School of Biomedical Sciences, University of Ulster, Cromore Road, Coleraine BT52 1SA, Co. Londonderry, Northern Ireland - UK

⁴Faculty of Agriculture, Azad University of Khorasgan, Isfahan - IRAN

Received: 28.03.2008

Abstract: *Amygdalus lycioides* var. *horrida* (Spach) Browicz (Rosaceae), also known as *Prunus lycioides* (Spach.) Schneid., is an endemic Iranian species of the genus *Amygdalus*. In Iranian traditional medicine, the aerial parts and roots of *A. lycioides* are used in the treatment of diabetes. Six flavonoids, i.e. quercetin 3-*O*-rhamnoside (1), luteolin 7-*O*-rhamnoside (2), isorhamnetin 3-*O*-rutinoside (3), kaempferol 3-*O*-rhamnoside (4), apigenin (5), and naringenin (6), have been isolated from the aerial parts of this plant. The structures of these compounds were elucidated by UV, MS, and NMR spectroscopic data analyses. While the antioxidant activity of these compounds was assessed by the 2,2-diphenyl-1-picryl-hydrazyl (DPPH) assay, the vasorelaxant effect was determined using the rat aortic vascular smooth muscle. Compounds 1-6 displayed significant antioxidant activity, with the RC_{50} values ranging from 0.0033 to 0.5186 mg/ml. Compound 2 showed a considerable vasorelaxant activity on rat aortic vascular smooth muscle in a dose-dependent manner.

Key Words: *Amygdalus lycioides* var. *horrida*, Rosaceae, flavonoid, antioxidant, DPPH, vasorelaxant

Amygdalus lycioides var. *horrida* Bitkisinin Flavonitlerinin Antioksidant ve Vasorelaksant Aktiviteleri

Özet: *Prunus lycioides* (Spach.) Schneid olarak isimlendirilen *Amygdalus lycioides* var. *horrida* (Spach) Browicz (Rosaceae) *Amygdalus* genusunda İran'a ait endemic bir bitkidir. İran halk tebabatinde şeker hastalığının tedavisinde *A. lycioides* bitkisinin hava kökleri kullanılır. Bu bitkinin hava köklerinden kuersetin 3-*O*-rhamnosit (1), luteolin 7-*O*-rhamnosit (2), isorhamnetin 3-*O*-rutinosit (3), kaempferol 3-*O*-rhamnosit (4), apigenin (5) ve naringenin (6) adlı altı flavonit izole edilmiştir. Bileşiklerin yapısı UV, MS ve NMR spektroskopisi ile belirlenmiştir. Bileşiklerin antioksidant aktiviteleri 2,2-difenil-1-pikril-hidrazil (DPPH) ile, vasorelaksant etki ise rat aortic vasküler düz kası kullanılarak belirlenmiştir. Bileşikler önemli derecede antioksidant aktivite gösterirken, RC_{50} değerleri 0,0033 ile 0,5186 mg/ml arasında gözlenmiştir. 2 nolu Bileşik ise doza bağımlı olarak rat vasküler düz kası üzerine önemli derecede vasorelaksant aktivite göstermiştir.

Anahtar Sözcükler: *Amygdalus lycioides* var. *horrida*, Rosaceae, flavonit, antioksidant, DPPH, vasorelaksant

Introduction

Amygdalus lycioides var. *horrida* (Spach) Browicz (Rosaceae), (*syn.* *Prunus lycioides* (Spach.) Schneid.), known as wild almond, is an endemic Iranian species of the genus *Amygdalus* (1,2). In Iranian traditional medicine, the aerial parts and roots of *A. lycioides* are used in the treatment of diabetes. Amygdalin, a

characteristic compound of the genus *Amygdalus*, was previously reported from this plant (3). However, there are no reports available to date on any phytochemical or bioactivity studies on *A. lycioides*. As a part of our on-going studies on the Iranian flora (4-14), we now report the isolation, identification, and antioxidant and vasorelaxant properties of 6 flavonoids, namely quercetin

3-*O*-rhamnoside (1), luteolin 7-*O*-rhamnoside (2), isorhamnetin 3-*O*-rutinoside (3), kaempferol 3-*O*-rhamnoside (4), apigenin (5), and naringenin (6), from the aerial parts of *A. lycioides*.

Materials and Methods

General experimental procedures

UV spectra were obtained in methanol (MeOH) using a Shimadzu UV-1650PC UV-visible spectrometer. NMR spectra were recorded in CD₃OD on a Bruker 200 MHz NMR Spectrometer (200 MHz for ¹H and 50 MHz for ¹³C) using residual solvent peak as internal standard. HPLC separation was performed in a Shimadzu photodiode-array detector (SPD-M20A). A Shim-Pack ODS preparative HPLC column (15 m, 250 mm × 20 mm) was used. A Sep-Pak Vac 35 cc (10 g) C₁₈ cartridge (Waters) was used for pre-HPLC fractionation. FAB-MS analyses were performed on a Finnigan MAT95 spectrometer.

Plant material

The aerial parts of *Amygdalus lycioides* var. *horrida* (Spach) Browicz were collected during April-May 2006 from Naeine in Esfahan province in Iran. A voucher specimen (TUM-ADE 0241) has been retained in The Herbarium of the Faculty of Pharmacy, Tabriz University of Medical Sciences, Iran.

Extraction and isolation of compounds (1-6)

The dried and ground aerial parts of *A. lycioides* (100 g) were Soxhlet extracted with *n*-hexane, dichloromethane, and methanol (MeOH), successively (1.1 l each), 10 cycles each. The MeOH extract (2 g) was subjected to Sep-Pack fractionation using a step gradient of MeOH-water mixture (10:90, 20:80, 40:60, 60:40, 80:20, and 100:0). The preparative reversed-phase HPLC analysis (linear gradient: 30% to 70% methanol in water in 50 min, flow rate: 20 ml/min) of the 40% methanolic Sep-Pack fraction yielded 2 flavonoids: quercetin 3-*O*-rhamnoside (1, 18.3 mg, *t_R* = 28.1 min) and luteolin 7-*O*-rhamnoside (2, 17.4 mg, *t_R* = 36.5 min). Similar purification of the 60% methanolic Sep-Pack fraction (linear gradient: 50% to 90% ACN in water in 50 min, flow rate: 20 ml/min) afforded 4 flavonoids: isorhamnetin 3-*O*-rutinoside (3, 1.7 mg, *t_R* = 8.2 min), kaempferol 3-*O*-rhamnoside (4, 3.5 mg, *t_R* = 11.8 min), apigenin (5, 2.1 mg, *t_R* = 21.1 min), and naringenin (6, 7.3 mg, *t_R* = 21.9 min). All of the compounds (1-6) were identified by spectroscopic means.

Quercetin 3-O-rhamnoside (1): Yellow amorphous solid; 18.3 mg; UV I_{max} (MeOH): 256, 268 sh, 299 sh, 362; +AlCl₃: 275, 305 sh, 331 sh, 437; +AlCl₃/HCl: 268, 299 sh, 366 sh, 405; +NaOMe: 272, 327, 409; +NaOAc: 274, 324, 380; +NaOAc/H₃BO₃: 262, 298 sh, 378 nm; FAB-MS *m/z* 471 [M + Na]⁺; ¹H-NMR (200 MHz, CD₃OD) and ¹³C-NMR (50 MHz, CD₃OD) as in accordance with the published data (15-18).

Luteolin 7-O-rhamnoside (2): Yellow amorphous solid; 17.4 mg; UV I_{max} (MeOH): 255, 267 sh, 348; +AlCl₃: 274, 299 sh, 329, 432; +AlCl₃/HCl: 273, 293 sh, 358, 387; +NaOMe: 263, 300 sh, 394; +NaOAc: 259, 266 sh, 365 sh, 405; +NaOAc/H₃BO₃: 259, 372 nm; FAB-MS *m/z* 455 [M + Na]⁺; ¹H-NMR (200 MHz, CD₃OD) and ¹³C-NMR (50 MHz, CD₃OD) as in accordance with the published data (15-18).

Isohamnetin 3-O-rutinoside (3): Yellow amorphous solid; 1.7 mg; UV I_{max} (MeOH): 254, 266 sh, 305 sh, 356; +AlCl₃: 268, 279 sh, 300 sh, 369 sh, 402; +AlCl₃/HCl: 267, 275 sh, 300 sh, 359 sh, 399; +NaOMe: 271, 328, 413; +NaOAc: 272, 320, 396; +NaOAc/H₃BO₃: 254, 267 sh, 304 sh, 360 nm; FAB-MS *m/z* 647 [M + Na]⁺; ¹H-NMR (200 MHz, CD₃OD) and ¹³C-NMR (50 MHz, CD₃OD) as in accordance with the published data (15-18).

Kaempferol 3-O-rhamnoside (4): Yellow amorphous solid; 11.8 mg; UV I_{max} (MeOH): 264, 317, 345; +AlCl₃: 268, 300, 335, 395; +AlCl₃/HCl: 274, 300, 335, 396; +NaOMe: 267, 326; +NaOAc: 266, 315 sh, 345; +NaOAc/H₃BO₃: 264, 315 sh, 345 nm; FAB-MS *m/z* 455 [M + Na]⁺; ¹H-NMR (200 MHz, CD₃OD) and ¹³C-NMR (50 MHz, CD₃OD) as in accordance with the published data (15-18).

Apigenin (5): Yellow amorphous solid; 2.1 mg; UV I_{max} (MeOH): 267, 296 sh, 336; +AlCl₃: 276, 302, 348, 384; +AlCl₃/HCl: 276, 299, 340, 380; +NaOMe: 276, 324, 392; +NaOAc: 274, 301, 376; +NaOAc/H₃BO₃: 268, 302 sh, 338 nm; FAB-MS *m/z* 293 [M + Na]⁺; ¹H-NMR (200 MHz, CD₃OD) and ¹³C-NMR (50 MHz, CD₃OD) as in accordance with the published data (15-18).

Naringenin (6): Yellow amorphous solid; 7.3 mg; UV I_{max} (MeOH): 287, 326 sh; +AlCl₃: 311, 378; +AlCl₃/HCl: 310, 373; +NaOMe: 248, 325; +NaOAc: 286 sh, 326; +NaOAc/H₃BO₃: 291, 332 sh nm; FAB-MS *m/z* 295 [M + Na]⁺; ¹H-NMR (200 MHz, CD₃OD) and ¹³C-NMR (50 MHz, CD₃OD) as in accordance with the published data (15-18).

Antioxidant activity

2,2-Diphenyl-1-picrylhydrazyl (DPPH), with the molecular formula of $C_{18}H_{12}N_5O_6$, was obtained from Fluka Chemie AG, Bucks, UK. Trolox[®] (6-hydroxy-2,5,7,8-tetramethylchroman-2-carboxylic acid) was obtained from Sigma-Aldrich, UK. The method used by Takao et al. (19) was adopted with appropriate modifications (20,21). DPPH (4 mg) was dissolved in MeOH (50 ml) to obtain a concentration of 80 mg/ml.

Qualitative assay: Compounds 1-6 were applied to a precoated silica gel TLC plate (0.25 mm thickness) and sprayed with DPPH solution using an atomizer. It was allowed to develop for 30 min. The colour change (purple on white) was noted.

Quantitative assay: Compounds 1-6 were dissolved in MeOH to obtain a concentration of 0.5 mg/ml. Dilutions were made to obtain concentrations of $5 \cdot 10^{-2}$, $5 \cdot 10^{-3}$, $5 \cdot 10^{-4}$, $5 \cdot 10^{-5}$, $5 \cdot 10^{-6}$, $5 \cdot 10^{-7}$, $5 \cdot 10^{-8}$, $5 \cdot 10^{-9}$, and $5 \cdot 10^{-10}$ mg/ml. Diluted solutions (1 ml each) were mixed with DPPH (1 ml) and allowed to stand for half an hour for any reaction to occur. The absorbance was recorded at 517 nm. The experiment was performed in triplicate and the average absorption was noted for each concentration. The same procedure was followed for the positive control, a well-known antioxidant Trolox[®]. The RC_{50} value, which is the concentration of the test material that reduces 50% of the free radical concentration, was calculated as mg/ml.

Vasorelaxant activity

The vasorelaxant activity of compounds 1, 2 and 4 was assessed in triplicate following the method described by Chan et al. (22) and Kim et al. (23). Rings of rat (Wistar 250-300 g) thoracic aorta 3-5 mm in length were trimmed from adjacent tissues. Two stainless steel triangular hooks were introduced through the lumen of the ring. One hook was fixed to the bottom of the organ bath and the other was connected to a force-displacement transducer. Each aortic ring was set up in a 10 ml bath containing modified Krebs Ringer-bicarbonate solution of the following composition (mM): NaCl 118, KCl 4.7, KH_2PO_4 1.2, $NaHCO_3$ 25, $MgSO_4 \cdot 7H_2O$ 1.2, $CaCl_2$ 2.5, glucose 11.1. The solution was equilibrated with a mixture of 95% O_2 and 5% CO_2 to give a pH of 7.3 to 7.4. Temperature was held at 37 °C. The optimal resting tension was adjusted to 2 g, which was obtained in preliminary tests and maintained throughout the experiments. Tissues were allowed to attain a steady level of tension during a 60 min accommodation period before being tested. During this

time, the bathing solution was changed every 15 min. Functional integrity of the endothelium was confirmed routinely at the beginning of the experiment by the presence or absence of relaxation induced by acetylcholine (6 μ M) on contraction induced by phenylephrine (0.1 μ M). Changes in isometric tension were recorded on a computer assisted data acquisition system (ADInstrument, Power Lab/4SP) with force displacement transducers (LETICA, Spain). In the various experiments, the vascular rings were contracted with prostaglandin F2 (PGF2, 10 μ M). When contraction was stable, the test compound (highest concentration: 800 M) was applied to the bath. Relaxation was expressed as percentage reversal of contraction induced by vasoactive agents.

Results and Discussion

Reversed-phase preparative HPLC analysis of the methanol extract of the aerial parts of *A. lycioides* var. *horrida* produced 6 flavonoids, which were identified as quercetin 3-*O*-rhamnoside (1), luteolin 7-*O*-rhamnoside (2), isorhamnetin 3-*O*-rutinoside (3), kaempferol 3-*O*-rhamnoside (4), apigenin (5), and naringenin (6), on the basis of UV, MS, and NMR data analyses (Figure 1).




Figure 1. Structures of the flavonoids isolated from *A. lycioides* var. *horrida*.

The UV absorption spectra in MeOH of 1-6 were typical for flavonoids (15). Further UV analyses using various shift reagents confirmed the presence/absence of free hydroxyls as well as the site of conjugations in these compounds (15). The FABMS spectra of all compounds (1-6) established the molecular mass and thereby the molecular formula of these compounds. The UV, ^1H -, and ^{13}C -NMR spectroscopic data of 1-6 were identical to the literature data for quercetin 3-*O*-rhamnoside (1), luteolin 7-*O*-rhamnoside (2), isorhamnetin 3-*O*-rutinoside (3), kaempferol 3-*O*-rhamnoside (4), apigenin (5), and naringenin (6) (15-18).

To our knowledge, this is the first report on the occurrence of flavonoids 1-6 in the aerial parts of *A. lycioides* var. *horrida*. The flavonoids found predominantly in the genus *Prunus*, which has a close taxonomical relation to *Amygdalus*, are isorhamnetin 3-*O*-rutinoside, isorhamnetin 3-*O*-glucoside, kaempferol 3-*O*-rutinoside, quercetin 3-*O*-galactoside, and isorhamnetin 3-*O*-galactoside (24-26). The co-occurrence of similar flavonoids both in *A. lycioides* and in *Prunus* species might have some chemotaxonomic implications; at least, this finding justifies the synonym *Prunus lycioides* for *Amygdalus lycioides*.

All compounds (1-6) exhibited significant levels of antioxidant activity in the DPPH assay. The RC_{50} (the concentration of the test material that reduces 50% of the free radical concentration, calculated as mg/ml) values of these compounds are presented in the Table. Among these flavonoids (1-6), quercetin 3-*O*-rhamnoside (1) and luteolin 7-*O*-rhamnoside (2) were the most potent ones

($\text{RC}_{50} = 0.0035$ and 0.0033 mg/ml), while naringenin (6) was the least potent ($\text{RC}_{50} = 0.5186$ mg/ml). The antioxidant properties of 1-6 indicated that not only the phenolic hydroxyls but also the C2-C3 double bond, as in compounds 1-5, was an important contributor to the antioxidant activity of these compounds. However, generally the antioxidant activity of 1-6, like that of other natural phenolic compounds, is a consequence of the presence of the phenolic moieties in the structures. The antioxidant activity of phenolic natural products is predominantly owing to their redox properties, i.e. the ability to act as reducing agents, hydrogen donors, and singlet oxygen quenchers, and to some extent could also be due to their metal chelation potential (21). The presence of these antioxidant compounds (1-6) in *A. lycioides* var. *horrida* might be significant in relation to this plant's various medicinal uses.

Due to the paucity of the samples, only compounds 1, 2, and 4 were assessed for their possible vasorelaxant activity. None of the flavonoids tested, except for luteolin 3-*O*-rhamnoside (2), showed any vasorelaxant property at the test concentrations (highest concentration: 800 M). Luteolin 3-*O*-rhamnoside (2) relaxed rat aortic vascular smooth muscle in a dose-dependent manner (Figure 2). This finding was in agreement with the results of previous studies on flavonoids conducted by other researchers (22,23,27,28). It is interesting to note that, although compounds 1, 2, and 4 are structurally quite similar, luteolin 3-*O*-rhamnoside (2) lacks any *O*-glycosylation at C-3. Thus, it is possible that *O*-glycosylation at C-3 rendered compounds 1 and 4 inactive.

Table. Antioxidant properties of compounds 1-6 in the DPPH assay.

Compounds	RC_{50} * value (mg/ml)
Quercetin 3- <i>O</i> -rhamnoside (1)	0.0035
Luteolin 7- <i>O</i> -rhamnoside (2)	0.0033
Isorhamnetin 3- <i>O</i> -rutinoside (3)	0.0403
Kaempferol 3- <i>O</i> -rhamnoside (4)	0.1605
Apigenin (5)	0.0043
Naringenin (6)	0.5186
Positive control: Trolox®	0.0026

* The concentration of the test material that reduces 50% of the free radical concentration was calculated as mg/ml




Figure 2. Vasorelaxant effect of luteoline 7-O-rhamnoside (Lu. 7-O-Rh) (2) in rat aortic rings precontracted by PGF_2 (10 M) in normal Krebs' solution.

Acknowledgement

We thank the EPSRC National Mass Spectrometry Service Centre (Department of Chemistry, University of Wales Swansea, Swansea, Wales, UK) for MS analyses.

Corresponding Author:

Satyajit Dey SARKER
School of Biomedical Sciences,
University of Ulster, Cromore Road,
Coleraine BT52 1SA, Co. Londonderry,
Northern Ireland, UK

References

1. Browicz K, Zohary D. The genus *Amygdalus* L. (Rosaceae): species relationships, distribution and evolution under domestication. *Genetic Resources and Crop Evolution* 43: 229-247, 1996.
2. GRIN Database. USDA, ARS, National Genetic Resources Program, Germplasm Resources Information Network - (GRIN) [Online Database], National Germplasm Resources Laboratory, Beltsville, Maryland. Available on-line at: <http://www.ars-grin.gov/cgi-bin/npgs/html/taxon.pl?3027>, 2008.
3. Siami A, Heidari R, Mohseni M. Comparative study of amygdalin, fat and total protein of 7 species of wild almond in West Azerbaijan (Iran). *Proceedings of the 3rd International Symposium on Pistachios and Almond* 591: 181-187, 2002.
4. Delazar A, Naseri M, Nazemiyeh H et al. Flavonol 3-methyl ether glucosides and a tryptophylglycine dipeptide from *Artemisia fragrans* (Asteraceae). *Biochem Syst Ecol* 35: 52-56, 2007.
5. Delazar A, Naseri M, Nahar L et al. GC-MS analysis and antioxidant activities of essential oils of two cultivated *Artemisia* species. *Chemistry of Natural Compounds* 43: 112-114, 2007.
6. Delazar A, Biglari F, Nazemiyeh H et al. GC-MS analysis of the essential oils, and the isolation of phenylpropanoid derivatives from the aerial parts of *Pimpinella aurea*. *Phytochemistry* 67: 2176-2181, 2006.
7. Delazar A, Modarresi M, Shoeb M et al. Eremostachiin: A new furanolabdane diterpene glycoside from *Eromostachys glabra*. *Natural Product Research* 20: 167-172, 2006.
8. Delazar A, Talischi B, Nazemiyeh Z et al. Chrozophorin: a new acylated flavone glucoside from *Chrozophora tinctoria*. *Revista Brasileira de Farmacognosia (Brazilian Journal of Pharmacognosy)* 16: 286-290, 2006.
9. Delazar A, Gibbons S, Kosari AR et al. Flavonoid C-glycosides and cucurbitacin glycosides from *Citrullus colocynthis*. *DARU* 14: 109-114, 2006.
10. Delazar A, Celik S, Gokturk RS et al. Two acylated flavonoids from *Stachys bombycina* and their free radical scavenging activity. *Die Pharmazie* 60: 878-880, 2005.
11. Delazar A, Reid RG, Sarker SD. GC-MS analysis of essential oil of the oleoresin from *Pistacia atlantica* var *mutica*. *Chemistry of Natural Compounds* 40: 24-27, 2004.
12. Delazar A, Byres M, Gibbons S et al. Iridoid glycosides from *Eremostachys glabra*. *J Nat Prod* 67: 1584-1587, 2004.
13. Nazemiyeh H, Delazar A, Ghahramani MA et al. Phenolic glycosides from *Phlomis lanceolata* (Lamiaceae). *Natural Product Communications* 3: 53-56, 2008.
14. Nazemiyeh H, Maleki N, Mehmani F et al. Assessment of anti-inflammatory properties ethyl acetate extract of *Stachys schtschegleevii* Sosn. *DARU* 4: 174-182, 2007.
15. Mabry TJ, Markham KR, Thomas MB. *The Systematic Identification of Flavonoids*, Springer-Verlag, New York, 1970.
16. Markham KR. *Techniques of Flavonoid Identification*, Academic Press, London, 1982.
17. Harborne JB, Mabry TJ. *The Flavonoids: Advances in Research*, Chapman and Hall, London, 1982.
18. Agrawal PK. *Carbon-13 NMR of Flavonoids*, Elsevier, Amsterdam, 1989.
19. Takao T, Watanabe N, Yagi I et al. A simple screening method for antioxidants and isolation of several antioxidants produced by marine-bacteria from fish and shellfish. *Biosci Biotech Biochem* 58: 1780-1783, 1994.
20. Kumarasamy Y, Fergusson M, Nahar L et al. Biological activity of moschamindole from *Centaurea moschata*. *Pharm Biol* 40: 307-310, 2002.
21. Kumarasamy Y, Byres M, Cox PJ et al. Isolation, structure elucidation and biological activity of flavone C-glycosides from the seeds of *Alliaria petiolata*. *Chemistry of Natural Compounds* 40: 122-128, 2004.

22. Chan EC, Pannangpetch P, Woodman OL. Relaxation to flavones and flavonols in rat isolated thoracic aorta: mechanism of action and structure-activity relationships. *J Cardiovasc Pharmacol* 35: 326-333, 2000.
23. Kim TJ, Kim JH, Jin YR et al. The inhibitory effect and mechanism of luteolin 7-glucoside on rat aortic vascular smooth muscle cell proliferation. *Arch Pharm Res* 29: 67-72, 2006.
24. Combined Chemical Dictionary. Chapman & Hall/CRC Press LLC. URL: <http://www.chemnetbase.com/>, 2008.
25. ISI database. ISI Web of Knowledge, Thomson ISI, London. Available on-line at <http://wok.mimas.ac.uk/>, 2008.
26. Milbury PE, Chen CY, Dolnikowski GG et al. Determination of flavonoids and phenolics and their distribution in almonds. *Journal of Agriculture and Food Chemistry* 54: 5027-5033, 2006.
27. Lemos VS, Cotes SF, dos Santos MH et al. Structure and vasorelaxant activity of floranol, a flavonoid isolated from the roots of *Dioclea grandiflora*. *Chemistry and Biodiversity* 3: 635-645, 2006.
28. Morello S, Vellecco V, Alfieri A et al. Vasorelaxant effect of the flavonoid galangin on isolated rat thoracic aorta. *Life Sciences* 78: 825-830, 2006.