

1-1-2006

## Antioxidant and Free Radical Scavenging Properties of Twelve Traditionally Used Indian Medicinal Plants

FARRUKH AQIL

IQBAL AHMAD

ZAFAR MEHMOOD

Follow this and additional works at: <https://journals.tubitak.gov.tr/biology>



Part of the [Biology Commons](#)

---

### Recommended Citation

AQIL, FARRUKH; AHMAD, IQBAL; and MEHMOOD, ZAFAR (2006) "Antioxidant and Free Radical Scavenging Properties of Twelve Traditionally Used Indian Medicinal Plants," *Turkish Journal of Biology*. Vol. 30: No. 3, Article 11. Available at: <https://journals.tubitak.gov.tr/biology/vol30/iss3/11>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Biology by an authorized editor of TÜBİTAK Academic Journals. For more information, please contact [academic.publications@tubitak.gov.tr](mailto:academic.publications@tubitak.gov.tr).

## Antioxidant and Free Radical Scavenging Properties of Twelve Traditionally Used Indian Medicinal Plants

Farrukh AQIL, Iqbal AHMAD, Zafar MEHMOOD

<sup>1</sup>Department of Agricultural Microbiology, Faculty of Agricultural Sciences, Aligarh Muslim University, Aligarh-202002, INDIA

<sup>2</sup>Himalaya Drug Company, Najaf Garh Road, New Delhi-110015, INDIA

Received: 01.06.2006

**Abstract:** The methanolic crude extracts of 12 traditionally used Indian medicinal plants were screened for their antioxidant and free radical scavenging properties using  $\alpha$ -tocopherol and butylated hydroxy toluene (BHT) as standard antioxidants. Antioxidant activity was measured by ferric thiocyanate (FTC) assay and compared with the thiobarbituric acid (TBA) method. Free radical scavenging activity was evaluated using diphenyl picryl hydrazyl (DPPH) radicals. The overall antioxidant activity of *Lawsonia inermis* was the strongest, followed in descending order by *Ocimum sanctum*, *Cichorium intybus*, *Piper cubeba*, *Punica granatum*, *Allium sativum*, *Delonix regia*, *Terminalia chebula*, *Terminalia bellerica*, *Mangifera indica*, *Camellia sinensis*, and *Trigonella foenum-graecum*. Seven plants, namely *Terminalia chebula*, *Mangifera indica*, *Terminalia bellerica*, *Punica granatum*, *Ocimum sanctum*, *Cichorium intybus*, and *Camellia sinensis*, showed strong free radical scavenging activity with the DPPH method. Phytochemical analysis of plant extracts indicated the presence of major phytochemicals, including phenolics, alkaloids, glycosides, flavonoids, and tannins. The phenolic concentrations in the above plants ranged from 28.66 to 169.67 mg/g of dry plant extract. A fair correlation between antioxidant/free radical scavenging activity and phenolic content was observed among 9 plants; however, in 3 plants (*Piper cubeba*, *Lawsonia inermis* and *Trigonella foenum-graecum*), no such relationship was observed. The tested plant extracts showed promising antioxidant and free radical scavenging activity, thus justifying their traditional use.

**Key Words:** Medicinal plant, antioxidant activity, free radical scavenging activity, phytochemicals, phenolics

### Introduction

Reactive oxygen species (ROS) are an entire class of highly reactive molecules derived from the metabolism of oxygen. ROS, including superoxide radicals, hydroxyl radicals, and hydrogen peroxide, are often generated as byproducts of biological reactions or from exogenous factors. In vivo, some of these ROS play positive roles in cell physiology; however, they may also cause great damage to cell membranes and DNA, inducing oxidation that causes membrane lipid peroxidation, decreased membrane fluidity, and DNA mutations leading to cancer, degenerative, and other diseases (1-4).

Mammalian cells possess elaborate defense mechanisms for radical detoxification. Key metabolic steps are superoxide dismutase (SOD), catalase (CAT), and glutathione peroxidase (GPX), which destroy toxic peroxides. In addition to antioxidant enzymes, non-enzymatic molecules, including thioredoxin, thiols, and disulfide-bonding play important roles in antioxidant defense systems. Some of the compounds are of an

exogenous nature and are obtained from food, such as  $\alpha$ -tocopherol,  $\beta$ -carotene, and ascorbic acid, and such micronutrient elements as zinc and selenium (5). If cellular constituents do not effectively scavenge free radicals, they lead to disease conditions as described above.

Antioxidant-based drugs/formulations for the prevention and treatment of complex diseases like atherosclerosis, stroke, diabetes, Alzheimer's disease, and cancer have appeared during the last 3 decades (6). This has attracted a great deal of research interest in natural antioxidants. Subsequently, a worldwide trend towards the use of natural phytochemicals present in berry crops, tea, herbs, oilseeds, beans, fruits, and vegetables has increased (7-9).

Several herbs and spices have been reported to exhibit antioxidant activity, including rosemary, sage, thyme, nutmeg, turmeric, white pepper, chili pepper, ginger, and several Chinese medicinal plant extracts (10-13). The majority of the active antioxidant compounds are

flavonoids, isoflavones, flavones, anthocyanins, coumarins, lignans, catechins, and isocatechins. In addition to the above compounds found in natural foods, vitamins C and E,  $\beta$ -carotene, and  $\alpha$ -tocopherol are known to possess antioxidant potential (14-16). A direct relationship between antioxidant activity and phenolic content of plant extracts has been reported (16,17). Epidemiological studies have shown that the consumption of foods and beverages rich in phenolic content can reduce the risk of heart disease (18).

Many Indian medicinal plants are considered potential sources of antioxidant compounds. In some cases, their active constituents are known. *Terminalia chebula*, *T. bellerica*, *T. muelleri*, and *Phyllanthus emblica*, all of which have antioxidant activity, showed high content of phenolics like gallic acid (19,20). Conversely, the antioxidant activity of *Hemidesmus indicus*, *Cichorium intybus*, *Withania somnifera*, *Ocimum sanctum*, *Mangifera indica*, and *Punica granatum*, as determined by several methods, has been poorly documented (19-23).

There is an increased quest to obtain natural antioxidants with broad-spectrum actions. The majority of the rich diversity of Indian medicinal plants are yet to be scientifically evaluated for such properties. Furthermore, the relationship between phenolic content and antioxidant activity is largely not examined in Indian medicinal plants. Among the above 12 medicinal plants that are commonly used in the Indian system of medicine, Ayurveda, Siddha, and Unani were selected for study. We have previously evaluated these plants for their broad-spectrum antimicrobial activity (24,25). In this report, we investigated the crude methanolic extracts of these plants for their potential antioxidant activity with the thiobarbituric acid (TBA), ferric thiocyanate (FTC), and 1, 1, diphenyl picryl hydrazyl (DPPH) radical scavenging methods.

## Materials and Methods

### Preparation of crude plant extracts

Test plants were collected locally or obtained from the Himalaya Drug Company, New Delhi-15. All the plant materials were further identified by the Department of Botany, Aligarh Muslim University (AMU), Aligarh. Voucher specimens of these plants were deposited in the department of Agricultural Microbiology, AMU, Aligarh. About 800 g of dried, ground plant material was soaked in 2.5 l of 98% methanol for 8-10 days, stirring every 18

h using a sterile glass rod. The final extract was passed through No.1 Whatman filter paper (Whatman Ltd., England). The filtrate obtained was concentrated under vacuum on a rotary evaporator at 40 °C and stored at 4 °C for further use. The crude extract was obtained by dissolving a known amount of the dry extract in 98% methanol to obtain a stock solution of 40 mg/ml concentrations.

### Antioxidant assay

The antioxidant activity of each plant extract was tested using the ferric thiocyanate (FTC) and thiobarbituric acid (TBA) methods. The FTC method was used to measure the amount of peroxide at the beginning of lipid peroxidation, in which peroxide will react with ferrous chloride and form ferric ions. Ferric ions will then unite with ammonium thiocyanate and produce ferric thiocyanate. The substance is red, and denser color is indicative of higher absorbance. The TBA method measures free radicals present after peroxide oxidation.

#### (a) Ferric thiocyanate (FTC) method

The standard method as described by Kikuzaki and Nakatani (12) was used. A mixture of 4.0 mg of plant extract in 4 ml of absolute ethanol, 4.1 ml of 2.52% linolenic acid in absolute ethanol, 8.0 ml of 0.05 M phosphate buffer (pH 7.0), and 3.9 ml of water was placed in a vial with a screw cap and then placed in a dark oven at 40 °C. To 0.1 ml of this solution were added 9.7 ml of 75% ethanol and 0.1 ml of 30% ammonium thiocyanate. Precisely 3 min after the addition of 0.1 ml of 0.02 M ferrous chloride in 3.5% HCl to the reaction mixture, the absorbance of red color was measured at 500 nm every 24 h until one day after the absorbance of control reached its maximum. Butylated hydroxy toluene (BHT) and  $\alpha$ -tocopherol were used as positive controls, while a mixture without a plant sample was used as the negative control.

#### (b) Thiobarbituric acid (TBA) method

The method of Ottolenghi (26) was followed. Two milliliters of 20% trichloroacetic acid and 2 ml of 0.67% 2-thiobarbituric acid were added to 1 ml of sample solution, prepared with the FTC method. The mixture was placed in a boiling water bath and after cooling was centrifuged at 3000 rpm for 20 min. Absorbance of supernatant was measured at 552 nm. Antioxidant activity was based on the absorbance on the final day of the FTC method.

### Free radical scavenging assay

The scavenging activity of DPPH free radicals by different plant extracts was determined according to the method reported by Gyamfi (27). Fifty microliters of the plant extract in methanol, yielding 100 µg/ml in each reaction, was mixed with 1 ml of 0.1 mM DPPH in methanol solution and 450 µl of 50 mM Tris-HCl buffer (pH 7.4). Methanol (50 µl) only was used as the experimental control. After 30 min of incubation at room temperature, the reduction in the number of DPPH free radicals was measured, reading the absorbance at 517 nm. BHT and α-tocopherol were used as controls. The percent inhibition was calculated from the following equation:

$$\% \text{ Inhibition} = \left[ \frac{\text{Absorbance of control} - \text{Absorbance of test sample}}{\text{Absorbance of control}} \right] \times 100$$

### Phytochemical analysis

Phytochemical analysis of the major phytoconstituents of the plant extracts was undertaken using standard qualitative methods (color tests and/or TLC) as described earlier (24).

### Total phenolic compound analysis

The amounts of phenolics in the selected Indian medicinal plant extracts were determined with Folin-Ciocalteu reagent using the method of Spanos and Wrolstad (28), as modified by Lister and Wilson (29). To 50 ml of each sample (3 replicates), 2.5 ml of 10% dilution of Folin-Ciocalteu reagent and 2 ml of Na<sub>2</sub>CO<sub>3</sub> (7.5%, w/v) were added and the resulting mixture was incubated at 45 °C for 15 min. The absorbance of all samples was measured at 765 nm using a Spectronic 20D<sup>+</sup>. Results were expressed as milligrams of gallic acid equivalent per gram of dry weight (mg GAE/g dw).

## Results and Discussion

Reactive oxygen species (ROS), from both endogenous and exogenous sources, may be involved in the etiologies of such diverse human diseases as arteriosclerosis, ischemic injury, cancer, and neurodegenerative diseases, as well as in processes like inflammation and ageing (5,30,31). There is evidence that indigenous antioxidants may be useful in preventing the deleterious consequences of oxidative stress and there is increasing interest in the protective biochemical functions of natural antioxidants contained in spices,

herbs, and medicinal plants (32,33). Our attention has been focused, in particular, on the parts of 12 commonly used Indian medicinal plants.

The antioxidant activities of the plant extracts were measured by the FTC method and compared with the TBA method at a concentration of 0.02% in methanolic solutions. The plant extracts tested showed low absorbance values, which indicated a high level of antioxidant activity. None of the plant extracts showed absorbance values greater than the negative controls (without plant extracts) at the end point of both methods, indicating the presence of antioxidant activity. However, all the plant extracts exhibited strong antioxidant activity as determined by both the FTC and TBA methods, surpassing the activity of the standard commercial antioxidants, alpha-tocopherol, and butylated hydroxy toluene (Figures 1-4).

The extracts of *Delonix regia* (flowers), *Terminalia bellerica* (fruits), *Terminalia chebula* (fruits), *Ocimum sanctum* (leaves), and *Camellia sinensis* (leaves) and some other plants exerted good antioxidant activity by various methods. The antioxidant activity detected with the TBA method was higher than that detected with the FTC method. This might suggest that the amount of peroxide in the initial stage of lipid peroxidation was less than the amount of peroxide in the secondary stage. Furthermore, the secondary product was much more stable for a period of time.

Free radicals have been implicated in many disease conditions, the important ones being superoxide radicals, hydroxy radicals, peroxy radicals, and single oxygen. Herbal drugs containing free radical scavengers are gaining importance in treating such diseases. Many plant extracts exhibit efficient antioxidant properties due to their phytoconstituents, including phenolics (34). In the present experiment, methanolic extracts of 12 plants were evaluated for their free radical scavenging activity using the DPPH radical assay. Reduction of DPPH radicals can be observed by the decrease in absorbance at 516 nm. Different plant extracts reduced DPPH radicals significantly. Values of percent decolorization of DPPH radicals are shown in Table 1. Of the 12 plants tested, 7 plants, namely *Mangifera indica*, *Punica granatum*, *Terminalia bellerica*, *Terminalia chebula*, *Cichorium intybus*, *Ocimum sanctum*, and *Camellia sinensis* showed more than 70% decolorization. The activity of *Trigonella foenum graecum* (57%) was at par when compared with




Figure 1. Antioxidant properties of plant extracts determined with the FTC method.




Figure 2. Antioxidant properties of plant extracts determined with the FTC method.

the commercial antioxidants  $\alpha$ -tocopherol (58%) and butylated hydroxy toluene (49%), while the extract of *Allium sativum* showed negligible activity.

Some variations in the extent of extract antioxidant activity were observed for each type of assay used in this study. The extracts of *Terminalia bellerica*, *Terminalia chebula*, and *Mangifera indica* had good DPPH radical

scavenging activity, but low lipid peroxidation, while the extracts of *Camellia sinensis*, *Cichorium intybus*, and *Ocimum sanctum* had relatively high antioxidant potential according to the FTC method. These differences may be due to their different antioxidant mechanisms; however, in some of the above plant extracts, the active constituents are known, such as gallic acid (*T. chebula*)




Figure 3. Antioxidant activities of plant extracts determined with the TBA method.




Figure 4. Antioxidant activities of plant extracts determined with the TBA method.

and catechins (*Camellia sinensis*). Our findings on the antioxidant activity of medicinal plants correlated with the reports of others (19-23).

A fair correlation between total phenolic content and antioxidant activity was observed in 9 of the 12 plants, whereas no direct relationship could be detected in *Piper cubeba*, *Lawsonia inermis*, and *Trigonella foenum-graecum* extracts. These observations clearly indicated a close linkage between phenolics and antioxidant activity;

however, due to the phytochemical diversity in the antioxidant phytochemicals, the above variation is expected.

The phytochemical analysis of the crude extracts indicated the presence of major phytochemicals, including phenolics, alkaloids, glycosides, flavonoids, and tannins (Table 1), which may have been responsible for the observed antioxidant activity. Our results further support the view that some traditionally used Indian

Table 1. Free radical scavenging activity of Indian medicinal plants determined with the DPPH method, and total phenolic contents.

S. No.	Name of the plant/ Family	Common Name	Part used	% Decolorization $\pm$ SD	*Phytocompounds detected	**Total phenolic contents $\pm$ SD
1.	<i>Allium sativum</i> L. (Alliaceae)	Lesan	Bulb	7.97 $\pm$ 1.09	P, T	28.66 $\pm$ 2.52
2.	<i>Camellia sinensis</i> L. (Theaceae)	Tea	Leaves	69.95 $\pm$ 3.62	P, G, S	163.33 $\pm$ 7.37
3.	<i>Cichorium intybus</i> L. (Compositae)	Chicory	Roots	73.34 $\pm$ 4.87	A, P, F, G, T	76.83 $\pm$ 8.75
4.	<i>Delonix regia</i> Gamble. (Leguminosae)	Gulmohar	Flowers	71.93 $\pm$ 3.86	A, P, F, G	169.67 $\pm$ 11.23
5.	<i>Lawsonia inermis</i> L. (Lythraceae)	Heena	Leaves	67.67 $\pm$ 5.48	A, P, G, S	38.67 $\pm$ 4.51
6.	<i>Mangifera indica</i> L. (Anacardiaceae)	Aam	Leaves	78.14 $\pm$ 3.63	F, G	135.00 $\pm$ 9.54
7.	<i>Ocimum sanctum</i> L. (Labiatae)	Tulsi	Leaves	73.70 $\pm$ 5.87	P, G, S	80.00 $\pm$ 8.18
8.	<i>Piper cubeba</i> L. (Piperaceae)	Kabab chini	Seeds	66.20 $\pm$ 3.20	P, F	42.83 $\pm$ 3.75
9.	<i>Punica granatum</i> L. (Punicaceae)	Anar	Rind	75.50 $\pm$ 3.90	A, P, F, G	122.00 $\pm$ 6.24
10.	<i>Terminalia bellerica</i> Roxb. (Combretaceae)	Bahera	Fruit	76.29 $\pm$ 3.06	P, F, G, S, T	111.67 $\pm$ 19.29
11.	<i>Terminalia chebula</i> Retz. (Combretaceae)	Harir	Fruit	85.36 $\pm$ 2.44	P, G, S	166.33 $\pm$ 18.01
12.	<i>Trigonella foenum-graecum</i> L. (Leguminosae)	Methi	Leaves	57.45 $\pm$ 2.44	A, F, G	74.33 $\pm$ 5.13
13.	BHT	----	----	49.28 $\pm$ 2.10		
14.	$\alpha$ -tocopherol	----	----	58.16 $\pm$ 4.13		

\*Phytocompounds Key: A, Alkaloids; P, Phenols; F, Flavonoids; G, Glycosides, S, Saponins; T, Tannins

\*\*mg/g of dry plant extracts

medicinal plants are promising sources of potential antioxidants. Further study will be aimed at isolating and identifying the substances responsible for the antioxidant activity of plant extracts, which may be further exploited in herbal formulations.

### Acknowledgments

We are grateful to Dr. S. Farooq, Director of The Himalaya Drug Company, New Delhi, for providing some of the plant materials and the Chairman of The Department of Agricultural Microbiology for his keen

interest in this work. We are also thankful to UGC, New Delhi, for financial support in the form of major research project No. F.3-58/ 2002 (SR-II).

### Corresponding author:

Iqbal AHMAD

Department of Agricultural Microbiology,

Aligarh Muslim University,

Aligarh, 202002 - INDIA

E-mail: iqbalahmad8@yahoo.co.in



## References

1. Cerutti AA. Oxidant stress and carcinogenesis. *Eur J Clin Invest* 21: 1-11, 1991.
2. Harman D. Free radical theory of aging, increase the functional life span. *Annals of the New York Academy of Sciences* 717: 1-15, 1994.
3. Ames B. Micronutrients prevent cancer and delay aging. *Toxicol Letters*, 102, 5-18, 1998.
4. Finkel T, Holbrook NJ. Oxidant, oxidative stress and biology of ageing. *Nature* 408: 239-247, 2000.
5. Halliwell B, Gutteridge JMC. Free radical in biology and medicine, 3<sup>rd</sup> Edition. Oxford University Press, London Chapter 3. 1998.
6. Devasagayam TPA, Tilak JC, Bloor KK et al. Review: Free radicals and antioxidants in human health: *Curr Stat Fut Prosp. JAPI*, 52: 794-804, 2004.
7. Dejana M, Arouma OI, Bianchi M et al. Inhibition of peroxynitrite dependent DNA base modification and tyrosin nitration by the extra virgin olive oil derived antioxidant hydroxytyrosol. *Free Radical Biol Med* 26: 762-769, 1999.
8. Lee KG, Shibamoto T. Antioxidant properties of the Aroma compounds isolated from soyabean and mung beans. *J Agri Food Chem* 48: 4290-4293, 2000.
9. Wang SY, Jiao H. Correlation of antioxidant capacities to oxygen radical scavenging enzyme activities in blackberry. *J Agr Food Chem* 48: 5672-5676, 2000.
10. Kikuzaki H, Usuguchi J, Nakatani N. Constituents of Zingiberaceae I. Diarylheptanoid from the rhizomes of ginger (*Zingiber officinale* Roscoe). *Chem Phar Bull* 39: 120, 1991.
11. Jitoe A, Masuda T, Tengah IGP et al. Antioxidant activity of tropical ginger extracts and analysis of the contained curcuminoids. *J Agr Food Chem* 8: 1337, 1992.
12. Kikuzaki H, Nakatani N. Antioxidant effects of some ginger constituents. *J Food Sci* 58: 1407-1410, 1993.
13. Lee SE, Hwang HJ, Ha JS et al. Screening of medicinal plant extracts for antioxidant activity. *Life Sci* 73: 167-179, 2003.
14. Prior RL. Fruit and vegetables in the prevention of cellular oxidative damage. *Am J Clin Nutr* 78: 570S-578S, 2003.
15. Cai Y, Luo Q, Sun M et al. Antioxidant activity and phenolic compounds of 112 traditional Chinese medicinal plants associated with anticancer. *Life Sci*. 74: 2157-2184, 2004.
16. Kaur C, Kapoor HC. Anti-oxidant activity and total phenolic content of some Asian vegetables. *Int J Food Sci Technol*. 37: 153-161, 2002.
17. Ivanova D, Gerova D, Chervenkov T et al. Polyphenols and antioxidant capacity of Bulgarian medicinal plants. *J Ethnopharmacol* 95: 145-150, 2005.
18. Landbo AK, Meyer AS. Ascorbic acid improves the antioxidant activity of European grape juices to inhibit lipid peroxidation of human LDL *in vitro*. *Int J Food Sci* 36: 727-736, 2001.
19. Saleem A, Ahotupa M, Pihlaja K. Total phenolics concentration and antioxidant potential of extracts of medicinal plants of Pakistan. *Z. Natureforsch.* 56: 973-978, 2001.
20. Bajpai M, Pande A, Tewari SK et al. Phenolic contents and antioxidant activity of some food and medicinal plants. *Int J Food Sci Nutr* 56: 287-291, 2005.
21. Sultana S, Khan N, Sharma S et al. Modulation of biochemical parameters by *Hemidesmus indicus* in cumene hydroperoxide-induced murine skin: possible role in protection against free radicals-induced cutaneous oxidative stress and tumor promotion. *J. Ethnopharmacol* 85: 33-41, 2003.
22. Russo A, Izzo AA, Cardile V et al. Indian medicinal plants as antiradical and DNA cleavage protectors. *Phytomedicine* 8: 125-132, 2001.
23. Ricci D, Giamperi L, Bucchini A et al. Antioxidant activity of *Punica granatum* fruits. *Fitoterapia* 77: 310-312, 2006.
24. Ahmad I, Beg AZ. Antimicrobial and phytochemical studies on 45 Indian medicinal plants against multi-drug resistant human pathogens. *J Ethnopharmacol* 74: 113- 123, 2001.
25. Aqil F, Ahmad I. Broad-spectrum antibacterial and antifungal properties of certain Indian medicinal plants. *World J Microbiol Biotechnol* 19: 653-657, 2003.
26. Ottolenghi A. Interaction of ascorbic acid and mitochondria lipids. *Arch Biochem Biophys*, 79, 355, 1959.
27. Gyamfi MA, Yonamine M, Aniya Y. Free radical scavenging action of medicinal herbs from Ghana *Thonningia sanguinea* on experimentally induced liver injuries. *Gen Pharmacol* 32: 661-667, 1999.
28. Spanos GA, Wrolstad RE. Influence of processing and storage on the phenolic composition of Thompson seedless grape juice. *J Agri Food Chem* 38: 1565-1571, 1990.
29. Lister E, Wilson P. Measurement of total phenolics and ABTS assay for antioxidant activity (personal communication). Crop Research Institute, Lincoln, New Zealand. 2001.
30. Good PF, Werner P, Hsu A et al. Evaluation of neuronal oxidative damage in Alzheimer's disease. *Am J Pathol* 149: 21-28, 1996.
31. Gassen M, Youdim MB. The potential role of iron chelators in the treatment of Parkinson's disease and related neurological disorders. *Pharmacol Toxicol* 80: 159-166, 1997.
32. Osawa T, Katsuzaki H, Kumon H et al. Protective role of phenolics antioxidants in plant against oxidative damage. In: *Frontiers of active oxygen species in biology and medicine* (Asada K, Yoshikawa T eds.), pp 333-336, 1994.
33. Noda Y, Anzai Kmori A, Kohono M et al. Hydroxyl and superoxide anion radical scavenging activities of natural source antioxidants using the computerized JES-FR30 ESR spectrometer system. *Bioch Mol Biol Inter* 42: 35-44, 1997.
34. Larson RA. The antioxidant of higher plants. *Phytochem* 27: 969-978, 1988.