

1-1-2010

Helminth parasites of the eastern spadefoot toad, *Pelobates syriacus* (Pelobatidae), from Turkey

HİKMET S. YILDIRIMHAN

CHARLES R. BURSEY

Follow this and additional works at: <https://journals.tubitak.gov.tr/zoology>



Part of the [Zoology Commons](#)

Recommended Citation

YILDIRIMHAN, HİKMET S. and BURSEY, CHARLES R. (2010) "Helminth parasites of the eastern spadefoot toad, *Pelobates syriacus* (Pelobatidae), from Turkey," *Turkish Journal of Zoology*. Vol. 34: No. 3, Article 5. <https://doi.org/10.3906/zoo-0810-2>

Available at: <https://journals.tubitak.gov.tr/zoology/vol34/iss3/5>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Zoology by an authorized editor of TÜBİTAK Academic Journals. For more information, please contact academic.publications@tubitak.gov.tr.

Helminth parasites of the eastern spadefoot toad, *Pelobates syriacus* (Pelobatidae), from Turkey

Hikmet S. YILDIRIMHAN^{1,*}, Charles R. BURSEY²

¹Uludağ University, Science and Literature Faculty, Department of Biology, 16059, Bursa - TURKEY

²Department of Biology, Pennsylvania State University, Shenango Campus, Sharon, Pennsylvania 16146 - USA

Received: 07.10.2008

Abstract: Ninety-one eastern spadefoot toads, *Pelobates syriacus*, were collected from 3 localities in Turkey between 1993 and 2003 and examined for helminths. One species of Monogenea (*Polystoma* sp.) and 3 species of Nematoda (*Aplectana brumpti*, *Oxysomatium brevicaudatum*, *Skrjabinelazia taurica*) were found. *Pelobates syriacus* represents a new host record for *Polystoma* sp. and *S. taurica*.

Key words: Monogenea, Nematoda, eastern spadefoot toads, *Pelobates syriacus*, Turkey

Türkiye'den toplanan toprak kurbağası (*Pelobates syriacus*)'nın (Pelobatidae) helmint parazitleri

Özet: 1993-2003 yılları arasında Türkiye'den 3 değişik yerden 91 toprak kurbağası helmintleri belirlenmek üzere toplanmıştır. İnceleme sonucunda 4 helmint türüne rastlanmıştır. Bunlardan biri Monogenea (*Polystoma* sp.), 3'ü (*Aplectana brumpti*, *Oxysomatium brevicaudatum*, *Skrjabinelazia taurica*) Nematoda'ya aittir. *Pelobates syriacus*, *Polystoma* sp. ve *S. taurica* için yeni konak kayıdır.

Anahtar sözcükler: Monogen, Nematoda, toprak kurbağası, *Pelobates syriacus*, Türkiye

Introduction

The eastern spadefoot toad, *Pelobates syriacus* Boettger, 1889, a fossorial species from Israel, Syria, and Turkey to Transcaucasica, lives in self-constructed burrows in loose and soft soil at elevations up to 1600 m, except during the breeding periods. In Turkey, it is found in suitable habitats in Anatolia and Turkish Thrace (Baran and Atatür, 1986). To our knowledge, there are 2 published reports of helminths in *P. syriacus*: Schad et al. (1960)

reported an occurrence of *Aplectana brumpti* and Yıldırımhan et al. (1997a) found *Oxysomatium brevicaudatum*. The purpose of this paper is to present a formal list of helminth species harbored by *P. syriacus*.

Materials and methods

Ninety -one eastern spadefoot toads, *Pelobates syriacus*, (31 female, 60 male; mean snout-vent length range [SVL] = 51-74 mm) were collected by hand

* E-mail: yhikmet@uludag.edu.tr

between 1993-2003 from 3 localities in Turkey: Bursa, Osmangazi district, (40°12'N, 29°00'E; elevation 120 m; N = 25); Konya province, Seydişehir district (37°17'N, 32°04'E; elevation 1090 m; N = 42); Edirne province, Büyükdöllük Marsh (41°45'N, 26°35'E; elevation 53 m; N = 24).

The toads were examined within 1 week of capture. They were over-anaesthetized with sodium pentobarbital and the body cavity was opened by an incision from vent to throat. The abdominal cavity, stomach, intestine, heart, lungs, liver, urinary bladder, and mouth were examined separately for helminths using a dissecting microscope. Helminths were killed in hot saline solution; nematodes were fixed in 70% ethanol and mounted in glycerol; monogenans were fixed in 70% ethanol, stained with iron-carmin (per Georgiev et al., 1986), cleared in clove oil, and mounted in Entellan® for examination with a compound microscope. Identifications are based upon the reference keys of Ryzhikov et al. (1980) and Prudhoe and Bray (1982). Helminth voucher specimens were deposited in the helminth collection of Uludağ University Museum of Zoology, Bursa, Turkey; Toad specimens were deposited in the Department of Biology, Uludağ University, Bursa, Turkey. Anuran nomenclature follows Frost (2008).

Results

Pelobates syriacus Boettger, 1889

Ninety-one *Pelobates syriacus* from 3 localities in Turkey were utilized for this study: Bursa, Osmangazi district (12 female, 13 male, SVL = 65.1 ± 4.3 mm, range 58-74 mm) collected in June 1993; Edirne, Büyükdöllük Marsh (3 females, 21 males, SVL = 58.7 ± 4.11 mm, range 61-73 mm) collected in May 2000; Konya, Seydişehir county (16 females, 26 males, SVL = 57.3 ± 4.26 mm, range 51-64 mm) collected in July 2000 (N = 19), August 2002 (N = 7), June 2003 (N = 16).

Polystoma sp.

Prevalence, mean intensity, and range: Hosts infected, 11 of 91 (12%, 1.70 ± 1.01, 1-4).

Temporal distribution: Konya-Seydişehir county: July 2000, 5 hosts with 1, 1, 1, 2, 2; August 2002, 1 host with 2; June 2003, 5 hosts with 1, 1, 1, 3, 4 respectively;

Site of infection: Urinary bladder.

Geographic range: The genus *Polystoma* has a cosmopolitan distribution (Diengdoh and Tandon, 1991).

Specimens deposited: Uludağ University Museum of Zoology

Remarks: *P. syriacus* represents a new host record for *Polystoma* sp. Three species of *Polystoma*, *P. macrocnemis* Biserkov, Yildirimhan and Ugurtas, 2001 in the Uludağ frog, *Rana macrocnemis*, *P. skrjabini* Buchvarov, 1984 in the European treefrog, *Hyla arborea*, and *P. viridis* Euzet, Combes and Batchvarov, 1974 in the European green toad, *Pseudepidalea viridis* (formerly *Bufo viridis*), have been reported from Turkey (Yildirimhan, 1999; Biserkov et al., 2001; Düşen and Öz, 2004). Euzet et al. (1974) assumed that each species of anuran harbored a distinct species of *Polystoma*. *Polystoma macrocnemis* and *P. viridis* are so closely related to the type species *P. integerrimum* (Frölich, 1791) with respect to both geographical range and morphometric characteristics that separation is difficult (Biserkov et al., 2001). It should also be noted that the brown frogs (*Rana temporaria* group) in Turkey are recognized as 2 species, *R. camerani* and *R. macrocnemis*, by some authors, e.g., Duellman (1999) or as the same species under the name *R. macrocnemis* by other authors, e.g., Anderson (1985), Baran and Atatür (1986). Thus more work on the species of *Polystoma* in Turkey is necessary before our specimens can be properly evaluated either as a new species or a previously described one.

Gallien (1935) studied the general biology of *Polystoma integerrimum* and reported that the adults lay eggs only in the spring, which give rise to larvae which attach themselves to the gills of tadpoles and remain attached through tadpole metamorphosis at which time the larvae migrate through the digestive tract to the urinary bladder.

Aplectana brumpti Travassos, 1931

(Syn: *Aplectana miranda* Ivanitzky, 1940; *Aplectana corti* Lopez-Neyru, 1947; *Aplectana ivanitzkyi* Morkov, Khonyakina, and Grivoreva 1972).

Prevalence, mean intensity, and range: Hosts infected, 25 of 91 (27%, 74.6 ± 61.66, 2-184).

Temporal distribution: Edirne-Büyükdöllük Marsh: May 2000, 10 hosts with 2, 40, 48, 102, 115, 118, 126, 126, 132, 146, respectively; Konya-Seydişehir county: July 2000, 6 hosts with 14, 20, 21, 32, 36, 48, respectively; August 2002, 3 hosts with 8, 14, 16, respectively; June 2003, 6 hosts with 4, 33, 142, 166, 172, 184, respectively;

Site of infection: Intestine.

Additional Turkish hosts: *Pseudepidalea viridis* (reported as *Bufo viridis* Schad et al., 1960); *Pelobates syriacus*, (Schad et al., 1960).

Type host and type locality: *Pseudepidalea viridis* (reported as *Bufo viridis*), Corsica, Spain (Travassos, 1931).

Other reported hosts: Common European toad, *Bufo bufo* (Fernández et al., 1987); natterjack toad, *Epidalea calamita* (reported as *Bufo calamita* Fernández et al., 1987); marsh frog, *Pelophylax ridibundus* (reported as *Rana ridibunda* Ivanitzky, 1940); *Pseudepidalea viridis* (reported as *Bufo viridis* Travassos, 1931; Lopez-Neyra, 1947; Kozák, 1969; Kozłowska, 1960; Frandsen, 1974; Baker, 1980a); European common frog, *Rana temporaria* (Ivanitzky, 1940); grass snake, *Natrix natrix* (Sharpilo, 1976); dice snake, *Natrix tessellata* (Sharpilo, 1976).

Geographic range: Western Europe (Baker, 1980a).

Specimens deposited: Uludağ University Museum of Zoology

Remarks: This is the second report of *A. brumpi* in *P. syriacus* from Turkey. Females produce eggs that larvate in utero before release to the environment where hatching occurs; the final host becomes infected orally (Anderson, 2000).

***Oxysomatium brevicaudatum* (Zeder, 1800) Railliet and Henry, 1916**

(Syn: *Oxysoma* Schneider, 1866 (preoccupied); *Neoxysomatium* Ballesteros-Marsquez, 1945; *Fusaria brevicaudata* Zeder, 1800; *Oxysoma contortum* Linstow, 1906; *Oxysomatium longespiculum* Railliet and Henry, 1916).

Prevalence, mean intensity and range: Hosts infected, 28 of 91 (31%, 18.3 ± 10.86, 3-45).

Temporal distribution: Edirne-Büyükdöllük Marsh: May 2000, 9 hosts with 14, 15, 22, 28, 30, 32, 35, 37, 45, respectively; Bursa-Osmangazi district May 1993, 19 hosts with 3, 6, 7, 7, 8, 8, 9, 10, 11, 12, 14, 14, 15, 16, 18, 20, 21, 24, 32, respectively;

Site of infection: Intestine.

Additional Turkish hosts: *Bufo bufo* (Yıldırımhan et al., 1997a); square-marked toad, *Amietophrynus regularis* (reported as *Bufo regularis*, Schad et al., 1960); *Pseudepidalea viridis* (reported as *Bufo viridis* Schad et al., 1960); *Pelobates syriacus* (Yıldırımhan et al., 1997a); *Rana macrocnemis* (Schad et al., 1960); *Rana ridibunda* (Schad et al., 1960; Yıldırımhan et al., 2005; Sağlam and Arıkan, 2006); *Natrix natrix* (Schad et al., 1960);

Type host and type locality: No type host, Germany

Other reported hosts: Midwife toad, *Alytes obstetricans* (Chabaud and Campana-Rouget, 1955; Baker, 1980b); *Amietophrynus regularis* (reported as *Bufo regularis* Baker, 1980b); European fire-bellied toad, *Bombina bombina* (Baker, 1980b); *Bufo bufo* (Kozłowska, 1960; Vojtkova et al., 1963; Kozak, 1969, 1973; Frandsen, 1974; Prokopic and Krivanec, 1975; Vojtkova, 1976; Baker, 1980b; Soriano et al., 1996; Galli et al., 2001; Shimalov and Shimalov, 2001; reported as *B. vulgaris*, Andre, 1912; Baylis, 1928); *Hyla arborea* (Kozłowska, 1960; Vojtkova, 1976; Baker, 1980b; Galeano et al., 1990); common spadefoot toad, *Pelobates fuscus* (Baker, 1980b); *Pelophylax ridibundus* (reported as *Rana ridibunda* Baker, 1980b; Kirin and Buchvarov, 2002; reported as *Rana esculenta* Kozłowska, 1960; Vojtkova et al., 1963; Prokopic and Krivanec, 1975; Vojtkova, 1976; Baker, 1980b); *Pseudepidalea viridis* (reported as *Bufo viridis*, Kozłowska, 1960; Prokopic and Krivanec, 1975; Vojtkova, 1976; Baker, 1980b); moor frog, *Rana arvalis* (Kozak, 1973; Prokopic and Krivanec, 1975; Vojtkova, 1976; Baker, 1980b; Cedhagen, 1988; reported as *Rana terrestris* Kozłowska, 1960); agile frog, *Rana dalmatina* (Baker, 1980b); Balkan stream frog, *Rana graeaca* (Hristovski, 1974; Baker, 1980b); Iberian frog, *Rana iberica* (Navarro et al, 1988); *Rana macrocnemis* (Baker, 1980b); *Rana temporaria* (Andre, 1913; Kozłowska, 1960; Vojtkova et al., 1963; Hristovski and Lees, 1973; Kozak, 1973; Prokopic and Krivanec, 1975; Vojtkova, 1976; Sattmann, 1986;

Navarro et al, 1988; Kirin and Buchvarov, 2002); smooth newt, *Lissotriton vulgaris* (reported as *Trituris vulgaris* Barus and Groschaft, 1962; Barus et al., 1963; Vojtkova, 1976; Baker, 1980b); Alpine newt, *Mesotriton alpestris* (reported as *Trituris alpestris* Kozłowska, 1960; Barus and Groschaft, 1962; Barus et al., 1963; Vojtkova, 1976; Baker, 1980b); Alpine salamander, *Salamandra atra* (Baker, 1980b); European fire salamander, *Salamandra salamandra* (Grabda and Grabda, 1953; Barus et al., 1963; Prokopic and Krivanec, 1975; Baker, 1980b; Bertman, 1986; reported as *S. maculosa* Andre, 1912; Vojtkova, 1976); Monte Albo cave salamander, *Speleomantes flavus* (Ricci, 1987); northern crested newt, *Trituris cristatus* (Barus et al., 1963; Frandsen, 1974; Vojtkova, 1976; Baker, 1980b); slow worm, *Anguis fragilis* (Zeder, 1800; Baylis, 1928; Sharpilo, 1976; Baker, 1980b; Moravec, 1963; Lewin, 1990; Shimalov et al., 2000; Borkovcova and Kopriva, 2004); sand lizard, *Lacerta agilis* (Lewin, 1992a); European legless lizard, *Pseudopus apodus* (reported as *Ophisaurus apodus* Baker, 1980b); *Natrix natrix* (Lukasiak, 1939; Baker, 1980b; Lewin, 1992b; Shimalov and Shimalov, 2000); *Natrix tessellata* (Sharpilo, 1976); northern viper, *Vipera berus* (Sharpilo, 1976; Shimalov and Shimalov, 2000).

Geographic range: Western Europe (Baker, 1987).

Specimens deposited: Uludağ University Museum of Zoology

Remarks: This is the second report of *O. brevicaudatum* in *P. syriacus*. The life history of *O. brevicaudatum* has apparently not been studied. It is assigned to the subfamily Cosmocercinae in which 2 pathways for infection are known, namely infection orally or skin penetration (Anderson, 2000).

***Skrjabinelazia taurica* Sypliaxov, 1930**

(Syn: *Salobrella* Freitas, 1940).

Prevalence, mean intensity, and range: Hosts infected, 1 of 91 (1%, 3).

Temporal distribution: Edirne-Büyükdöllük Marsh: May 2000, 1 host with 3 respectively;

Site of infection: Intestine.

Type host and type locality: Crimean wall lizard, *Podarcis tauricus* (reported as *Lacerta taurica*), Crimea, USSR (Sypliaxov, 1930).

Additional Turkish hosts: *Podarcis tauricus* (reported as *Lacerta taurica* Schad et al., 1960); European green lizard, *Lacerta viridis* (Schad et al., 1960).

Other reported hosts: *Podarcis tauricus* (reported as *Lacerta taurica* Sharpilo, 1976).

Geographic range: Turkey, Ukraine (Baker, 1987).

Specimens deposited: Uludağ University Museum of Zoology

Remarks: Species of *Skrjabinelazia* have previously been reported only from lizards belonging to the families Gekkonidae or Lacertidae (Lhermitte, 2008). In some species, females produce 2 types of eggs, namely membranous eggs in which larvae develop and hatch in utero, and thick-shelled eggs, which are released (Lhermitte, 2008). Given the number of *S. taurica* found in this study, the infection reported here may represent an incidental infection acquired through some nontypical behavior of the host. However, *P. syriacus* would represent a new host record for *S. taurica*.

Discussion

In all, 2400 helminths were collected from 50 (55%) of the 91 toads examined. Four helminth species were found: 1 monogenean (N = 19) and 3 nematodes (N = 2381). No individual host contained more than 3 species of helminths; of the infected toads: 36 (72%) harbored 1 species of helminth, 13 (26%) harbored 2 species, and 1 (2%) harbored 3 species. The collection sites yielded different communities of helminths: toads collected in Bursa harbored only *O. brevicaudatum*; toads from Edirne harbored *A. brumpti*, *O. brevicaudatum*, and *S. taurica*; toads from Konya harbored *Polystoma* sp. and *A. brumpti*. The reasons for the lack of a helminth species at a particular location are currently unknown.

There were 1.30 ± 0.07 ($X \pm 1$ SE) (range 1-3) helminth species per infected toad and 48.00 ± 8.72 (range 1-184) helminth individuals per infected toad. Aho (1990) compiled distributional patterns for anurans in general and reported the mean (\pm SE) total number of helminth species per host species as 3.54 ± 0.24 (range 0-9). Thus the infection rates for *P. syriacus* are lower than these seen for anurans in

Table. Reported helminths of Turkish anurans.

	<i>Bombina orientalis</i>	<i>Bufo bufo</i>	<i>Pseudis viridis</i>	<i>Hyla arborea</i>	<i>Pelobates syriacus</i>	<i>Pelodytes caucasicus</i>	<i>Rana camerani</i>	<i>Rana holzschuhi</i>	<i>Rana macrocnemis</i>	<i>Pelophryllax rufibundus</i>
Monogenea										
<i>Polystoma</i> sp.	-	-	-	-	6	-	8	-	10	-
<i>Polystoma macrocnemis</i>	-	-	-	-	-	-	-	-	9,11	-
<i>Polystoma skrjabini</i>	-	-	-	5	-	-	-	-	-	-
<i>Polystoma viridis</i>	-	-	4	-	-	-	-	-	-	-
Digenea										
<i>Bucephalus polymorphus</i>	-	-	-	-	-	-	-	-	-	16
<i>Candidotrema loossi</i>	-	-	-	-	-	-	-	-	-	13
<i>Diplodiscus</i> sp.	-	-	-	-	-	-	-	-	-	15
<i>Diplodiscus subclavatus</i>	-	-	-	-	-	-	-	-	-	12,13,14,16,18
<i>Gorgodera</i> sp.	-	-	-	-	-	-	-	-	-	15
<i>Gorgodera cygnoides</i>	-	-	-	-	-	7	8	-	-	13,14,18
<i>Gorgoderina</i> sp.	-	-	-	-	-	-	-	-	-	15
<i>Gorgoderina vitelliloba</i>	-	-	-	-	-	-	8	9	9,10	13,14,17,18
<i>Haematoloechus breviansa</i>	-	-	-	-	-	-	-	-	-	13,14,18
<i>Haematoloechus variegatus</i>	-	-	-	-	-	-	-	-	-	13,17
<i>Haplometra cylindracea</i>	-	-	-	-	-	-	8,19	9	9,19	-
<i>Opisthioglyphe ranae</i>	1	-	-	-	-	-	-	-	-	13,14,18
<i>Opisthioglyphe rastellus</i>	-	-	-	-	-	-	8,19	9	9	-
<i>Plagiorchis</i> sp.	-	-	-	-	-	-	-	-	-	15
<i>Pleurogenes claviger</i>	-	-	-	-	-	-	-	-	9,10	12, 16
<i>Pleurogenoides medians</i>	-	-	-	5	-	-	8,19	-	9	12,13,14,17,18
<i>Pleurogenoides stromi</i>	-	-	-	-	-	-	-	-	-	14
<i>Prostocus confusus</i>	-	-	-	-	-	-	-	-	-	13,14,18
<i>Brachycoelium salamandrae</i>	-	-	-	-	-	-	-	-	-	18
<i>Rauschiella</i> sp.	-	-	-	-	-	-	-	-	-	13
<i>Encyclometra colubrimurorum</i> (metacercariae)	-	-	-	5	-	-	-	-	-	18
<i>Codonocephalus urmigerus</i> (metacercariae)	-	-	-	-	-	-	-	-	-	13,14,18
Cestoda										
<i>Nematotaenia dispar</i>	-	-	4	-	-	-	8	-	-	-
<i>Proteocephalus</i> sp. (juvenile)	-	-	4	5	-	-	-	-	-	-
Nematoda										
<i>Agfa tauricus</i>	-	-	-	-	-	7	-	-	-	-
<i>Aplectana brumpti</i>	-	-	3	-	3,6	7	-	-	-	-
<i>Cosmocerca</i> sp.	1	-	-	-	-	-	-	-	10	2,13,15
<i>Cosmocerca ornata</i>	-	-	3	-	-	7	8,19	9	3,9,19	3,14
<i>Cosmocerca commutata</i>	-	-	3,4	5	-	-	-	-	-	18
<i>Cosmocercoides</i> sp.	-	-	4	-	-	-	-	-	-	17
<i>Oswaldocruzia</i> sp.	-	2	3	-	-	-	-	-	-	13
<i>Oswaldocruzia filiformis</i>	-	-	4	-	-	7	8	-	3,9,10	14,17
<i>Neoxysomatium</i> sp.	-	-	-	-	-	-	-	-	-	18
<i>Oxysomatium brevicaudatum</i>	-	2	3,4	-	2,6	-	-	-	3	3,14,17
<i>Rhabdias</i> sp.	-	-	3	-	-	-	-	-	3	-
<i>Rhabdias bufonis</i>	1	2	4	-	-	7	8	-	9	2,13,16,17,18
<i>Skrjabinelazia taurica</i>	-	-	-	-	6	-	-	-	-	-
<i>Eustrongylides</i> sp.	-	-	-	-	-	-	-	-	-	18
<i>Eustrongylides excisus</i> (larvae)	-	-	-	-	-	-	-	-	-	14,17
<i>Abbreviata</i> sp.	-	-	-	-	-	-	-	-	-	18
Acanthocephala										
<i>Acantocephalus</i> sp.	-	-	-	-	-	-	-	-	-	15
<i>Acantocephalus ranae</i>	1	-	4	5	-	-	8	-	9,10,19	2,13,14,16,17,18
<i>Centrorhynchus</i> sp.	-	-	-	-	-	-	-	-	-	14
<i>Pseudoacanthocephala caucasicus</i>	-	-	-	-	-	7	-	-	-	-

1. Yildirimhan et al., 2001; 2. Yildirimhan, Oğuz and Uğurtaş, 1997a; 3. Schäd, 1960; 4. Yildirimhan, 1999; 5. Düşen and Öz, 2004; 6. this study; 7. Yildirimhan, 2009; 8. Yildirimhan et al, 2006a; 9. Yildirimhan et al, 2006b; 10. Yildirimhan, Uğurtaş and Altunel, 1997b; 11. Biserkov et al. 2001; 12. Oğuz et al., 1994; 13. Yildirimhan et al., 1996; 14. Yildirimhan et al., 2005; 15. Saygi and Başbüyük, 1990; 16. Kir et al., 2001; 17. Sağlam and Arıkan, 2006; 18. Düşen, 2006; 19. Düşen, 2007.

general. Whether the more terrestrial habitat of *P. syriacus* is responsible for this difference will require additional work.

Fourteen species of anurans occur in Turkey (Bombinatoridae: *Bombina bombina* (Linnaeus, 1758); Bufonidae: *Bufo bufo* (Linnaeus, 1758), *Pseudepidalea viridis* (Laurenti, 1768) formerly *Bufo viridis* (Laurenti, 1768); Hylidae: *Hyla arborea* (Linnaeus, 1758), *Hyla savignyi* (Audoin, 1827); Pelobatidae: *Pelobates fuscus* (Laurenti, 1768), *P. syriacus* (Boettger, 1889); Pelodytidae: *Pelodytes caucasicus* (Boulenger, 1896); Ranidae: *Rana camerani* Boulenger, 1886, *R. dalmatina* Bonaparte, 1840, *R. holtzi* F. Werner, 1898, *R. macrocnemis* Boulenger, 1885, *Pelophylax bedriagae* (Camerano, 1882) previously *Rana levantina* Schneider and Sinsch, 1992, *P. ridibundus* (Pallas, 1771) (Baran and Atatür 1998). To our knowledge, Turkish population of *Hyla savignyi*, *Pelobates fuscus*, *Pelophylax bedriagae*, and *Rana dalmatina* have not been examined for helminths.

The Table presents reported helminths from Turkish anurans. It is of interest to note that the nematodes commonly infecting Turkish anurans,

namely, *Aplectana brumpti*, *Cosmocerca ornata*, *Oswaldocruzia filiformis*, *Oxysomatium brevicaudatum*, and *Rhabdias bufonis*, directly infect the host (Anderson, 2000). Thus for nematodes, habitat is more important than diet in determining rates of infection. Acanthocephalans require at least 2 hosts in the life cycle; arthropods are the usual intermediate hosts in which the infect stage develops and when eaten by an appropriate final host develops to maturity in the digestive tract (Nickol, 1985). Species of *Polystoma* infect directly (Gallien, 1935). The digeneans generally utilize a molluscan first intermediate host from which cercariae leave and penetrate a frog host directly or a variety of invertebrate hosts, which are then eaten by the final host (Smyth and Smyth, 1980). Presence of the helminths in *P. syriacus* in some areas but not others is most likely the result of patchy distribution on the part of a particular helminth species as well as the discontinuous distribution patterns of the host. The conclusion that we draw is that generalist helminths may vary within a particular host over time and space, but within its population of hosts, the helminth species is persistent.

References

- Aho, J.M. 1990. Helminth communities of amphibians and reptiles: comparative approaches to understanding patterns and processes. Pages 157-195 in G.W. Esch, A.O. Bush, and J.M. Aho (eds). *Parasite Communities: Patterns and Processes*. Chapman and Hall, New York, U.S.A.
- Anderson, R.C. 2000. *Nematode Parasites of Vertebrates. Their Development and Transmission*. 2nd edition. CABI Publishing, Wallingford, Oxon, U.K. 650 pp.
- Anderson, S.C. 1985. Amphibians. Pp. 987-990 in E. Yarshater (ed.). *Encyclopaedia Iranica*, Vol, 1. London: Routledge and Kegan Paul.
- André, E. 1912. Recherches parasitologiques sur les amphibiens de la suisse. *Revue Suisse de Zoologie* 20: 471-485.
- André, E. 1913. Recherches parasitologiques sur les amphibiens de la suisse. II. *Revue Suisse de Zoologie* 21: 179-200.
- Baker, M.R. 1980a. Revision of Old World species of the genus *Aplectana* Railliet & Henry, 1916 (Nematoda, Cosmocercidae). *Museum National d'Histoire Naturelle* 2: 955-998.
- Baker, M.R. 1980b. A revision of the genus *Oxysomatium* Railliet & Henry, 1916 (Nematoda, Cosmocercidae). *Bulletin Museum National d'Histoire Naturelle Paris* 2: 707-718.
- Baker, M.R. 1987. Synopsis of the Nematoda parasitic in amphibians and reptiles. Memorial University of Newfoundland, Occasional Papers in Biology 11: 1-325.
- Baran, I. and Atatür, M.K. 1986. A taxonomic survey of the mountain frogs of Anatolia. *Amphibia-Reptilia* 7: 115-133.
- Baran, I. and Atatür, M.K. 1998. *Turkish Herpetofauna (Amphibians and Reptiles)*. Republic of Turkey, Ministry of Environment, Ankara. 214 pp.
- Barus, V. and Groschaft, J. 1962. Helminthofauna colku *Triturus alpestris* (Laurenti, 1768) a *Triturus vulgaris* L. ze Sumavske Oblasti. *Zoologicke Listy* 11: 253-264.
- Barus, V., Groschaft, J. and Otcenasek, M. 1963. Helminthofauna ocasatych obozivelniku z uzemi Ceskoslovenska. *Ceskoslovenska Parasitologie* 10: 43-59.
- Baylis, H.A. 1928. Records of some parasitic worms from British vertebrates. *Annals and Magazine of Natural History (Series 10)* 1: 329-343.
- Bertman, M. 1986. *Brachycoelium salamandrae* (Frolich, 1789) (Trematoda, Brachycoeliidae) u Salamandry plamistej—*Salamandra salamandra* L. *Wiadomosci Parazytologiczne* 32: 173-176.

- Biserkov, V.Y., Yildirimhan, H.S., Buchvarov, G. and Uğurtaş, I.H. 2001. *Polystoma macrocnemis* n. sp. (Monogenea: Polystomatidae) from the Iranian longlegged wood frog *Rana macrocnemis* (Ranidae) in Turkey. *Systematic Parasitology* 48: 61-66.
- Borkovcova, M. and Kopriva, J.K. 2004. Parasitic helminths of reptiles (Reptilia) in South Moravia (Czech Republic) *Parasitology Research On line* 25 Nov 2004 2 p. Print version 2005, (5: 77-78).
- Cedhagen, T. 1988. Endoparasites in some Swedish amphibians. *Acta Parasitologica Polonica* 33: 107-113.
- Chabaud, A.G. and Campana-Rouget, Y. 1955. Helminthes de la Region de Banyuls I. Nematodes parasites d'amphibiens. *Vie et Milieu* 6: 83-92.
- Diengdoh, C.R. and Tandon, V. 1991. A new species of *Polystoma* (Monogenea) parasitic in rhacophorid amphibians in Meghalaya, India. *Helminthologia* 28: 173-178.
- Duellman, W.E. 1999. Patterns of distribution of amphibians. A global perspective. Johns Hopkins University Press, Baltimore, Maryland. 633 pp.
- Düşen, S. and Öz, M. 2004. Helminth parasites of the tree frog, *Hyla arborea* (Anura: Hylidae) from southwest Turkey. *Comparative Parasitology* 71: 258-261.
- Düşen, S. and Öz, M. 2006. Parasitic Helminths of the Marsh Frog, *Rana ridibunda* Pallas, 1771 (Anura: Ranidae), from Antalya Province, south-west Turkey. *Comparative Parasitology*, 73: 121-129.
- Düşen, S. 2007. Helminths of the Two Mountain Frogs, Banded Frog, *Rana camerani* Boulenger, 1886 and Uludağ Frog *Rana macrocnemis* Boulenger, 1885 (Anura: Ranidae), Collected from Antalya Province. *Acta Parasitologica Turcica* 31: 84-88.
- Euzet, L., Combes, C. and Batchvarov, G. 1974. Sur un nouveau Polystomataidae Européen, parasite de l'amphibien *Bufo viridis* Laur. *Vie et Milieu* 24: 129-140.
- Fernandez, J.P., Raga, J.A. Carbonell, E. and Babio, C.R. 1987. Sobre algunos nematodos parasitos encontrados en anfibios del genero *Bufo*. *Boletin de la Real Sociedad Espanola de Historia Natural Seccion Biologie* 83: 93-100.
- Frandsen, F. 1974. A study of Danish amphibian parasite fauna. *Acta Parasitologica Polonica* 22: 49-66.
- Frost, D.R. 2008. Amphibian Species of the World: an Online Reference. Version 5.2 (15 July, 2008). Electronic Database accessible at <http://research.amnh.org/herpetology/amphibia/index.php>. American Museum of Natural History, New York, USA.
- Galeano, M., Navarro, P. and Lluch, J. 1990. Helminthofauna de *Hyla* spp (Amphibia, Hylidae) en algunas localidades espanolas. *Miscellanea Zoologica* 14: 1-6.
- Galli, P., Crosa, G., Gentili, A. and Santagostino, M. 2001. New geographical records of parasitic nematodes from *Bufo bufo* in Italy. *Parassitologia* 43: 147-149.
- Gallien, L. 1935. Recherches expérimentales sur le dimorphisme évolutif et la biologie de *Polystmum integerrimum* Fröl. *Travaux de la Station Zoologique de Wimereux* 12: 1-181.
- Georgiev, B.B., Biserkov, V.Y. and Genov, T. 1986. *In toto* staining method for cestodes with iron acetocarmine. *Helminthologica* 23: 279-281.
- Grabda, E. and Grabda, J. 1953. Contribution a la connaissance de la faune parasitaire de salamandre *Salamandra salamandra*(L.) des environs de Bielsko (Pologne). *Fragmenta Faunistica Mussei Zoologici Polonici* 6: 243-247.
- Hristovski, N.D. 1974. Helminthofauna na *Rana graeca* Boulenger od Bitolsko (Makedonija-Jugoslavija). *Acta Parasitologica Iugoslavica* 6: 3-5.
- Hristovski, N.D. and Lees, E. 1973. The helminth fauna of *Rana temporaria* in relation to that of Europe generally. *Acta Parasitologica Iugoslavica* 4: 93-98.
- Ivanitzky, S.V. 1940 [On the helminth fauna of vertebrates in the Ukraine (Cestoda, Nematoda and Acanthocephala)]. *Sbornik Trudov Kharkovskogo Veterinarnogo Instituta* 19: 129-155. In Russian
- Kır, İ., Yildırım, M.Z., Becer, A. and İkiz, R. 2001. The feeding habits and parasites of the lake frogs (*Rana ridibunda* Palas 1771; Anura: Ranidae) of lake Eğirdir. *Acta Parasitologica Turcica* 25: 83-87.
- Kirin, D. and Buchvarov, G. 2002. Biodiversity and trematode assemblages in *Rana ridibunda* Pallas. From the district of Trojan town. *Experimental Pathology and Parasitology*. 5: 7-12.
- Kozák, A. 1969. Die nematodenfauna der Frösche der Theissebene. *Helminthologica* 10: 285-295.
- Kozák, A. 1973. Die Trematodenfauna des darpathengebietes der CSSR. *Biologia (Bratislava)* 28: 335-350.
- Kozłowska, J. 1960. On the nematodes of amphibians of Poland, mainly from the environment of Lodz. *Acta Parasitologica Polonica* 8: 215-230.
- Lewin, J. 1990. Parasitic worms in a slowworm (*Anguis fragilis* L.) population from the Bieszczady Mountains (Poland). *Acta Parasitologica Polonica* 35: 207-215.
- Lewin, J. 1992a. Parasites of the sand lizard (*Lacerta agilis* L.) in Poland. *Acta Parasitologica* 37: 19-24.
- Lewin, J. 1992b. Parasites of *Lacerta vivipara* Jacquin, 1787 in Poland. *Acta Parasitologica* 37: 79-82.
- Lhermitte, N., Bain, O. and Virga, A. 2008. *Skrjabinelazia rizzoi* n. sp. (Nematoda: Seuratoidea) from a Sicilian lacertid, with comments on specific and biological diversity in the genus. *Parasite* 15: 45-52.
- Lopez-Neyra, C.R. 1947. Helminths of the Vertebrados Ibéricos. Consejo Superior de Investigaciones Científicas, Granada. 1212 pp.

- Lukasiak, J. 1939. Badania nad fauna helmintologiczna Polski. Fragmenta Faunistica Musei Zoologici Polonici 4: 93-106.
- Moravec, F. 1963. Príspevek k poznani helmintofauny nasich plaz. Spisy Prirodovedcke Fakulty University v Brne 446: 353-396.
- Navarro, P., Izquierdo, S. Perez-Soler, R. Hornero, M.J. and Lluch, J. 1988. Contribucion al conocimiento de la helmintofauna de los herpetos ibericos. VIII. Nematoda Ascaridida Skrjabin et Schultz, 1940 de *Rana* spp. Revista Ibérica de Parasitología 48: 167-174.
- Nickol, B.B. 1985. Epizootiology. Pages 307-346 in Biology of the Acanthocephala D.W.T. Crompton and B. B. Nickol eds. Cambridge University Press, Cambridge, UK.
- Oğuz, M.C., Altunel, F.N. and Uğurtaş, I.H.. 1994. [An investigation of the species of platyhelminthes and *Acanthocephalus ranae* (Schränk 1788, Echinorhynchida, Acanthocephala) of marsh frogs (*Rana ridibunda* Pallas, 1771) with were collected from the Bursa and Edirne Regions.] Turkish Journal of Zoology 18: 47-51.
- Prokopic, J. and Krivanec, K. 1975. Helminths of amphibians, their interaction and host-parasite relationships. Acta Scientiarum Naturalium Brno 9: 1-48.
- Prudhoe, S. and Bray, R.A. 1982. Platyhelminth parasites of the amphibia. British Museum (Natural History), Oxford University Press, Oxford, UK, 217 pp. + 4 microfiche.
- Ricci, M. 1987. Parassiti del gen. *Speleomantes* (Amphibia Urodela: Plathelminthes) in Italia. Rivista di Parasitologia 4: 5-25.
- Ryzhikov, K.M., Sharpilo, V.P. and Shevechenko, N.N. 1980. [Helminths of amphibians of the fauna of the USSR]. Izdatel'stov Nauka, Moscow, Russia. 278 pp. In Russian.
- Saglam, N. and Arıkan, H. 2006. Endohelminth fauna of the marsh frog *Rana ridibunda* from Lake Hazar, Turkey. Diseases of Aquatic Organisms 72: 253-260.
- Sattmann, V.H. 1986. Über die Helminthenfauna von *Triturus alpestris* Laurenti 1768 and *Rana temporaria* L. aus Almtümpeln in Oberösterreich. (Amphibia, Plathelminthes und Nemathelminthes). Annalen des naturhistorischen Museums in Wien, Serie B. 87: 193-196.
- Saygi, G. and Başbüyük, H.H. 1990. *Rana ridibunda*'ların Bağırsak ve İdrar Keselerinde Bulduğumuz Parazitler. Türkiye Parazitoloji Dergisi 15: 105-118.
- Schad, G.A., Kuntz, R.E. and Wells, W.H. 1960. Nematode parasites from Turkish vertebrates. An annotated list. Canadian Journal of Zoology 38: 949-963.
- Sharpilo, V.P. 1976. [Parasitic worms of the reptilian fauna of the USSR]. Izdat. 'Naukova Dumka', Kiev, USSR. 287 pp. In Russian.
- Shimalov, V.V. and Shimalov, V.T. 2001. Helminth fauna of toads in Belorussian Polesie. Parasitology Research 87:84.
- Shimalov, V.V., Shimalov, V.T. and Shimalov, A.V. 2000. Helminth fauna of lizards (Reptilia, Sauria) in the southern part of Belarus. Parasitology Research 86: 343.
- Smyth, J. D. and Smyth, M. M. 1980. Frogs as host-parasite systems. I. An introduction to parasitology through the parasites of *Rana temporaria*, *R. esculenta* and *R. pipiens*. Macmillan Press, Ltd., London, U.K. 112 pp.
- Soriano, F., Navarro, P. and Lluch, J. 1996. Contribucion al conocimiento de la helmintofauna de los herpetos ibericos IX. Helminths de *Bufo bufo* (Linnaeus, 1758) (Amphibia, Bufonidae). Boletín de la Real Sociedad Española de Historia Natural Sección Biología 92: 89-93.
- Syplixov, O. 1930. Sur un nouveau nematode des reptiles: *Skrjabinelazia taurica* n. g., n. sp. Annales de Parasitologie Humaine et Comparee 8: 615-618.
- Travassos, L. 1931. Pesquisas helmintologicas realizadas em Hamburgo. IX. Ensaio monographico da familia Cosmocercidae Trav., 1925 (Nematoda). Memorias do Instituto Oswaldo Cruz 25: 237-298.
- Vojtková, L. 1976. [Nematoda of amphibians of CSSR I. Adult nematodes]. Folia Facultatis Scientiarum Naturalium Universitatis Purkynianae Brunrunensis Biologia 17: 5-80. In Czech.
- Vojtková, L., Moravec, F. and Nabelkova, L. 1963. [On the recent stage of investigation of the amphibian helminth fauna of CSSR]. Casopis Narodniho Musea. Oddil Prirodovendny 9: 121-131. In Czech.
- Yıldırımhan, H.S. 1999. [Researches on parasitic helminths of *Bufo viridis* Laurenti, 1768 (Anura: Amphibia)]. Turkish Journal of Zoology 23: 177-195. In Turkish.
- Yıldırımhan, H.S., Aydoğdu, A. Uğurtaş, I.H. and Altunel, F.N. 2001. [Helminth fauna of *Bombina bombina* (Linnaeus, 1761), fire-bellied toad, from Sakarya and Edirne, Turkey]. Acta Parasitologica Turcica 25: 308-311 In Turkish.
- Yıldırımhan, H.S., Karadeniz, E., Gürkan, E and Koyun, M. 2005. [Metazoon parasites of the marsh frog *Rana ridibunda* Pallas, 1771 (Anura) collected from different regions in Turkey.] Acta Parasitologica Turcica 29: 135-139. In Turkish.
- Yıldırımhan, H.S., Oğuz, M.C. and Uğurtaş, I.H. 1997a. [An investigation of the nematodes of some tailless frogs (*Rana ridibunda*, *Bufo bufo*, *Pelobates syriacus*) collected from the Bursa regions]. Hacettepe Fen ve Mühendislik Bilimleri Dergisi 18: 45-58. In Turkish.
- Yıldırımhan, H.S., Uğurtaş, I.H. and Altunel, F.N. 1996. [An investigation of helminths of *Rana ridibunda* Pallas, 1771 (marsh frogs)]. Türkiye Parazitoloji Dergisi 20: 113-130. In Turkish.
- Yıldırımhan, H.S., Uğurtaş, I.H. and Altunel, F.N. 1997b. [An investigation on parasitic helminths of *Rana macrocnemis* Boulenger, 1885 (Uludag frog)]. Turkish Journal of Zoology 21: 467-473. In Turkish.
- Yıldırımhan, H.S., Goldberg S.R. and Burse, C.R. 2006 a. Helminth Parasites of the Banded Frog *Rana camerani* (Ranidae) from Turkey. Comparative Parasitology. 73: 222-236.

Yildirimhan, H.S., Bursey C.R. and Goldberg, S.R. 2006b. Helminth Parasites of the Taurus Frog, *Rana holtzi*, and the Uludag Frog, *Rana macrocnemis*, with Remarks on the Helminth Community of Turkish Anurans. *Comparative Parasitology* 73: 237-248.

Yildirimhan, H.S., Bursey C.R. and Goldberg, S.R. 2009. Helminth Parasites of the Caucasian frog, *Pelodytes caucasicus*, from Turkey. *Comparative Parasitology* 76: 247-257.

Zeder, J.G.H. 1800. Erster Nachtrag zur Naturgeschichte der Eingeweidewürmer mit Aufässen und Anmerkungen herausgegeben. Leipzig, Germany. 320 pp.