

1-1-2015

Approximation properties of Szász type operators based on Charlier polynomials

ARUN KAJLA

PURSHOTTAM NARAIN AGRAWAL

Follow this and additional works at: <https://dctubitak.researchcommons.org/math>



Part of the [Mathematics Commons](#)

Recommended Citation

KAJLA, ARUN and AGRAWAL, PURSHOTTAM NARAIN (2015) "Approximation properties of Szász type operators based on Charlier polynomials," *Turkish Journal of Mathematics*: Vol. 39: No. 6, Article 17.

<https://doi.org/10.3906/mat-1502-80>

Available at: <https://dctubitak.researchcommons.org/math/vol39/iss6/17>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Mathematics by an authorized editor of TÜBİTAK Academic Journals.

Approximation properties of Szász type operators based on Charlier polynomials

Arun KAJLA*, Purshottam Narain AGRAWAL

Department of Mathematics, Indian Institute of Technology Roorkee, Roorkee, India

Received: 28.02.2015

Accepted/Published Online: 12.07.2015

Printed: 30.11.2015

Abstract: In the present paper, we study some approximation properties of the Szász type operators involving Charlier polynomials introduced by Varma and Taşdelen in 2012. First, we establish approximation in a Lipschitz type space and weighted approximation theorems for these operators. Then we obtain the error in the approximation of functions having derivatives of bounded variation.

Key words: Szász operator, Charlier polynomials, modulus of continuity, bounded variation

1. Introduction

In 1950, Szász [20] introduced the following linear positive operators:

$$S_n(f; x) = e^{-nx} \sum_{k=0}^{\infty} \frac{(nx)^k}{k!} f\left(\frac{k}{n}\right), \quad (1.1)$$

where $x \in [0, \infty)$ and $f(x)$ is a continuous function on $[0, \infty)$ whenever the above sum converges uniformly. Many researchers have studied approximation properties of these operators and modified Szász operators by involving different types of polynomials. Jakimovski and Leviatan [13] defined a generalization of Szász operators including the Appell polynomials and gave the approximation properties of these operators. In [21], Varma et al. considered the generalization of Szász operators involving Brenke type polynomials and studied convergence properties by using the Korovkin type theorem and the order of convergence with the help of classical method.

Recently, Altomare et al. [3] introduced a new kind of generalization of Szász–Mirakjan–Kantorovich operators and obtained the rate of convergence by means of suitable moduli of smoothness. Several researchers also defined different types of generalizations of these operators and studied their approximation properties; we refer the reader to those papers (cf. [5, 4, 7, 8, 15, 17]).

In [22], Varma and Taşdelen constituted a link between orthogonal polynomials and the positive linear operators. They considered Szász type operators including Charlier polynomials. These polynomials [11] have the generating functions of the form

$$e^t \left(1 - \frac{t}{a}\right)^u = \sum_{k=0}^{\infty} C_k^{(a)}(u) \frac{t^k}{k!}, \quad |t| < a, \quad (1.2)$$

*Correspondence: rachitkajla47@gmail.com

2010 AMS Mathematics Subject Classification: 26A15, 40A35, 41A25, 41A36.

where $C_k^{(a)}(u) = \sum_{r=0}^k \binom{k}{r} (-u)_r \left(\frac{1}{a}\right)^r$ and $(m)_0 = 1, (m)_j = m(m+1)\cdots(m+j-1)$ for $j \geq 1$.

For $C_\gamma[0, \infty) := \{f \in C[0, \infty) : |f(t)| \leq Me^{\gamma t} \text{ for some } \gamma > 0, M > 0 \text{ and } t \in [0, \infty)\}$, Varma and Taşdelen [22] defined the following Szász type operators involving Charlier polynomials

$$\mathcal{L}_n(f; x, a) = e^{-1} \left(1 - \frac{1}{a}\right)^{(a-1)nx} \sum_{k=0}^{\infty} \frac{C_k^{(a)}(-(a-1)nx)}{k!} f\left(\frac{k}{n}\right), \quad (1.3)$$

where $a > 1$ and $x \in [0, \infty)$. For the special case $a \rightarrow \infty$ and $x - \frac{1}{n}$ instead of x , these operators reduce to the operators (1.1). They studied uniform convergence of these operators by applying the Korovkin theorem on compact subsets of $[0, \infty)$ and the order of approximation by using the classical modulus of continuity.

This paper is structured as follows. In Section 2, we present some moment estimates and a result needed to study approximation of functions with derivatives of bounded variation. In Section 3, we discuss the main results of the paper wherein we establish approximation in a Lipschitz type space and weighted approximation theorems for the operators \mathcal{L}_n . Lastly, we obtain the rate of convergence for the functions having a derivative of bounded variation on every finite subinterval of $[0, \infty)$, for these operators.

2. Preliminaries

Let $e_i(x) = x^i, i = 0, 1, 2, \dots$

Lemma 1 [22] *For the operators $\mathcal{L}_n(f; x, a)$, we get*

$$(i) \quad \mathcal{L}_n(e_0(t); x, a) = 1;$$

$$(ii) \quad \mathcal{L}_n(e_1(t); x, a) = x + \frac{1}{n};$$

$$(iii) \quad \mathcal{L}_n(e_2(t); x, a) = x^2 + \frac{x}{n} \left(3 + \frac{1}{a-1}\right) + \frac{2}{n^2}.$$

Lemma 2 *For the operators $\mathcal{L}_n(f; x, a)$, we have*

$$(i) \quad \mathcal{L}_n(e_3(t); x, a) = x^3 + \frac{x^2}{n} \left(6 + \frac{3}{a-1}\right) + \frac{2x}{n^2} \left(\frac{1}{(a-1)^2} + \frac{3}{a-1} + 5\right) + \frac{5}{n^3};$$

$$(ii) \quad \mathcal{L}_n(e_4(t); x, a) = x^4 + \frac{x^3}{n} \left(10 + \frac{6}{a-1}\right) + \frac{x^2}{n^2} \left(31 + \frac{30}{a-1} + \frac{11}{(a-1)^2}\right)$$

$$+ \frac{x}{n^3} \left(67 + \frac{31}{a-1} + \frac{20}{(a-1)^2} + \frac{6}{(a-1)^3}\right) + \frac{15}{n^4}.$$

Proof From the generating function of the Charlier polynomials given by (1.2) we have

$$\sum_{k=0}^{\infty} \frac{k^3 C_k^{(a)}(- (a - 1)nx)}{k!} = e\left(1 - \frac{1}{a}\right)^{-(a-1)nx} \left[n^3 x^3 + x^2 n^2 \left(6 + \frac{3}{a-1}\right) + 2xn \left(\frac{1}{(a-1)^2} + \frac{3}{a-1} + 5\right) + 5 \right]$$

$$\sum_{k=0}^{\infty} \frac{k^4 C_k^{(a)}(- (a - 1)nx)}{k!} = e\left(1 - \frac{1}{a}\right)^{-(a-1)nx} \left[n^4 x^4 + x^3 n^3 \left(10 + \frac{6}{a-1}\right) + x^2 n^2 \left(31 + \frac{30}{a-1} + \frac{11}{(a-1)^2}\right) + nx \left(67 + \frac{31}{a-1} + \frac{20}{(a-1)^2} + \frac{6}{(a-1)^3}\right) + 15 \right],$$

from which the lemma is immediate. □

Let $e_i^x(t) = (t - x)^i, i = 0, 1, 2 \dots$

Lemma 3 For the operators $\mathcal{L}_n(f; x, a)$, we have

(i) $\mathcal{L}_n(e_1^x(t); x, a) = \frac{1}{n};$

(ii) $\mathcal{L}_n(e_2^x(t); x, a) = \frac{ax}{n(a-1)} + \frac{2}{n^2};$

(iii) $\mathcal{L}_n(e_4^x(t); x, a) = \frac{x}{n^3} \left(17 + \frac{49}{(a-1)} - \frac{20}{(a-1)^2} + \frac{6}{(a-1)^3}\right) + \frac{x^2}{n^2} \left(19 - \frac{46}{(a-1)} + \frac{3}{(a-1)^2}\right) + \frac{15}{n^4}.$

Proof Using Lemmas 1 and 2, the proof of this lemma easily follows. Hence, the details are omitted. □

To study the rate of convergence of functions having a derivative of bounded variation, let us rewrite the operators (1.3) as

$$\mathcal{L}_n(f; x, a) = \int_0^{\infty} f(w) \frac{\partial}{\partial w} \{ \mathcal{K}_n(x, w, a) \} dw, \tag{2.1}$$

where

$$\mathcal{K}_n(x, w, a) = \begin{cases} \sum_{k \leq nw} e^{-1} \left(1 - \frac{1}{a}\right)^{(a-1)nx} \frac{C_k^{(a)}(- (a - 1)nx)}{k!}, & \text{if } 0 < w < \infty, \\ 0, & \text{if } w = 0. \end{cases}$$

From Lemma 3, for $x \in (0, \infty)$ and sufficiently large n , we have

$$\mathcal{L}_n(|e_1^x(t)|; x, a) \leq (\mathcal{L}_n(e_2^x(t); x, a))^{1/2} \leq \sqrt{\frac{\lambda(a)x}{n}}, \tag{2.2}$$

where $\lambda(a)$ is some positive constant depending on a .

We also get, for $r \geq 2$ and fixed $x \in [0, \infty)$,

$$\mathcal{L}_n(e_{2r}^x(t); x, a) = O(n^{-r}); \quad n \rightarrow \infty. \tag{2.3}$$

Lemma 4 For all $x \in (0, \infty)$ and sufficiently large n , we have

$$(i) \vartheta_{n,a}(x, t) = \int_0^t \frac{\partial}{\partial w} \{ \mathcal{K}_n(x, w, a) \} dw \leq \frac{1}{(x-t)^2} \frac{\lambda(a)x}{n}, \quad 0 \leq t < x;$$

$$(ii) 1 - \vartheta_{n,a}(x, z) = \int_z^\infty \frac{\partial}{\partial w} \{ \mathcal{K}_n(x, w, a) \} dw \leq \frac{1}{(z-x)^2} \frac{\lambda(a)x}{n}, \quad x < z < \infty,$$

where $\lambda(a)$ is a constant as described in (2.2).

Proof First we prove (i).

$$\begin{aligned} \vartheta_{n,a}(x, t) &= \int_0^t \frac{\partial}{\partial w} \{ \mathcal{K}_n(x, w, a) \} dw \leq \int_0^t \left(\frac{x-w}{x-t} \right)^2 \frac{\partial}{\partial w} \{ \mathcal{K}_n(x, w, a) \} dw \\ &\leq \frac{1}{(x-t)^2} \mathcal{L}_n((w-x)^2; x, a) \\ &\leq \frac{1}{(x-t)^2} \frac{\lambda(a)x}{n}. \end{aligned}$$

The proof of (ii) is similar, and hence it is omitted. □

Theorem 1 [22] Let $f \in C[0, \infty) \cap E$. Then

$$\lim_{n \rightarrow \infty} \mathcal{L}_n(f; x, a) = f(x),$$

and the operators given by (1.3) converge uniformly in each compact subset of $[0, \infty)$ where

$$E := \{ f : [0, \infty) \rightarrow \mathbb{R}, |f(x)| \leq Me^{Ax}, A \in \mathbb{R} \quad \text{and} \quad M \in \mathbb{R}^+ \}.$$

In what follows, let $\tilde{C}_B[0, \infty)$ be the space of all real valued bounded and uniformly continuous functions f on $[0, \infty)$, endowed with the norm $\|f\|_{\tilde{C}_B[0, \infty)} = \sup_{x \in [0, \infty)} |f(x)|$.

3. Main results

3.1. Degree of approximation

Let $a_1, a_2 > 0$ be fixed. We consider the following Lipschitz type space (see [18]):

$$Lip_M^{(a_1, a_2)}(r) := \left\{ f \in \tilde{C}_B[0, \infty) : |f(t) - f(x)| \leq M \frac{|t-x|^r}{(t+a_1x^2+a_2x)^{\frac{r}{2}}}; x, t \in (0, \infty) \right\},$$

where M is a positive constant and $0 < r \leq 1$.

Theorem 2 Let $f \in Lip_M^{(a_1, a_2)}(r)$ and $r \in (0, 1]$. Then, for all $x \in (0, \infty)$, we have

$$| \mathcal{L}_n(f; x, a) - f(x) | \leq M \left(\frac{\zeta_{n,a}(x)}{a_1x^2 + a_2x} \right)^{\frac{r}{2}},$$

where $\zeta_{n,a}(x) = \mathcal{L}_n(e_x^x(t); x, a)$.

Proof Applying the Hölder's inequality with $p = \frac{2}{r}$ and $q = \frac{2}{2-r}$, we find that

$$\begin{aligned} |\mathcal{L}_n(f; x, a) - f(x)| &\leq \left\{ e^{-1} \left(1 - \frac{1}{a}\right)^{(a-1)nx} \sum_{k=0}^{\infty} \frac{C_k^{(a)}(- (a-1)nx)}{k!} \left| f\left(\frac{k}{n}\right) - f(x) \right|^{\frac{2}{r}} \right\}^{r/2} \\ &\leq M \left\{ e^{-1} \left(1 - \frac{1}{a}\right)^{(a-1)nx} \sum_{k=0}^{\infty} \frac{C_k^{(a)}(- (a-1)nx)}{k!} \frac{\left(\frac{k}{n} - x\right)^2}{\left(\frac{k}{n} + a_1x^2 + a_2x\right)} \right\}^{r/2}. \end{aligned}$$

Since $f \in Lip_M^{(a_1, a_2)}(r)$ and $\frac{1}{\sqrt{\frac{k}{n} + a_1x^2 + a_2x}} < \frac{1}{\sqrt{a_1x^2 + a_2x}}, \forall x \in (0, \infty)$, we have

$$\begin{aligned} |\mathcal{L}_n(f; x, a) - f(x)| &\leq \frac{M}{(a_1x^2 + a_2x)^{\frac{r}{2}}} \left\{ e^{-1} \left(1 - \frac{1}{a}\right)^{(a-1)nx} \sum_{k=0}^{\infty} \frac{C_k^{(a)}(- (a-1)nx)}{k!} \left(\frac{k}{n} - x\right)^2 \right\}^{r/2} \\ &\leq M \left(\frac{\zeta_{n,a}(x)}{a_1x^2 + a_2x} \right)^{\frac{r}{2}}. \end{aligned}$$

This completes the proof of the theorem. □

Next, we obtain a local direct estimate for the operators defined in (1.3) using the Lipschitz-type maximal function of order r introduced by Lenze [16] as

$$\tilde{\omega}_r(f, x) = \sup_{t \neq x, t \in [0, \infty)} \frac{|f(t) - f(x)|}{|t - x|^r}, \quad x \in [0, \infty) \text{ and } r \in (0, 1]. \tag{3.1}$$

Theorem 3 Let $f \in \tilde{C}_B[0, \infty)$ and $0 < r \leq 1$. Then, for all $x \in [0, \infty)$, we have

$$|\mathcal{L}_n(f; x, a) - f(x)| \leq \tilde{\omega}_r(f, x) (\zeta_{n,a}(x))^{\frac{r}{2}}.$$

Proof From equation (3.1), we have

$$|\mathcal{L}_n(f; x, a) - f(x)| \leq \tilde{\omega}_r(f, x) \mathcal{L}_n(|t - x|^r; x, a).$$

Applying Hölder's inequality with $p = \frac{2}{r}$ and $q = \frac{2}{2-r}$, we get

$$|\mathcal{L}_n(f; x, a) - f(x)| \leq \tilde{\omega}_r(f, x) (\mathcal{L}_n((t - x)^2; x, a))^{\frac{r}{2}} \leq \tilde{\omega}_r(f, x) (\zeta_{n,a}(x))^{\frac{r}{2}}.$$

Thus, the proof is completed. □

3.2. Weighted approximation

Let $B_\varphi[0, \infty)$ be the space of all real valued functions on $[0, \infty)$ satisfying the condition $|f(x)| \leq M_f \varphi(x)$, where M_f is a positive constant depending only on f and $\varphi(x) = 1 + x^2$ is a weight function. Let

$C_\varphi[0, \infty)$ be the space of all continuous functions in $B_\varphi[0, \infty)$ with the norm $\|f\|_\varphi := \sup_{x \in [0, \infty)} \frac{|f(x)|}{\varphi(x)}$ and

$$C_\varphi^*[0, \infty) := \left\{ f \in C_\varphi[0, \infty) : \lim_{x \rightarrow \infty} \frac{|f(x)|}{\varphi(x)} \text{ is finite} \right\}.$$

The usual modulus of continuity of f on $[0, b]$ is defined as

$$\omega_b(f, \delta) = \sup_{|t-x| \leq \delta} \sup_{x, t \in [0, b]} |f(t) - f(x)|.$$

Theorem 4 *Let $f \in C_\varphi[0, \infty)$. Then we have*

$$\|\mathcal{L}_n(f; \cdot, a) - f\|_{C[0, b]} \leq 4M_f(1 + b^2)\zeta_{n, a}(b) + 2\omega_{b+1}(f, \sqrt{\zeta_{n, a}(b)}),$$

where $\zeta_{n, a}(b) = \frac{ab}{n(a-1)} + \frac{2}{n^2}$.

Proof Let $x \in [0, b]$ and $t > b + 1$. Then $t - x > 1$, and hence

$$\begin{aligned} |f(t) - f(x)| &\leq M_f(2 + t^2 + x^2) = M_f \{2 + 2x^2 + (t - x)^2 + 2x(t - x)\} \\ &\leq M_f(t - x)^2 (3 + 2x + 2x^2) \\ &\leq 4M_f(1 + x^2)(t - x)^2. \end{aligned} \tag{3.2}$$

For $x \in [0, b]$ and $t \in [0, b + 1]$ we have

$$|f(t) - f(x)| \leq \omega_{b+1}(f; |t - x|) \leq \left(1 + \frac{|t - x|}{\delta}\right) \omega_{b+1}(f; \delta). \tag{3.3}$$

Thus, from (3.2) and (3.3) for all $x \in [0, b]$ and $t \geq 0$, we have

$$|f(t) - f(x)| \leq 4M_f(1 + x^2)(t - x)^2 + \left(1 + \frac{|t - x|}{\delta}\right) \omega_{b+1}(f, \delta), \delta > 0.$$

Hence, applying Cauchy–Schwarz inequality, we get

$$\begin{aligned} |\mathcal{L}_n(f; x, a) - f(x)| &\leq 4M_f(1 + x^2)\mathcal{L}_n((t - x)^2; x, a) + \omega_{b+1}(f, \delta) \left(1 + \frac{1}{\delta}\mathcal{L}_n(|t - x|; x, a)\right) \\ &\leq 4M_f(1 + x^2)\zeta_{n, a}(x) + \omega_{b+1}(f, \delta) \left(1 + \frac{1}{\delta}\sqrt{\zeta_{n, a}(x)}\right) \\ &\leq 4M_f(1 + b^2)\zeta_{n, a}(b) + \omega_{b+1}(f, \delta) \left(1 + \frac{1}{\delta}\sqrt{\zeta_{n, a}(b)}\right). \end{aligned}$$

Choosing $\delta = \sqrt{\zeta_{n, a}(b)}$, we get the desired result. □

Theorem 5 *Let $f \in C_\varphi^*[0, \infty)$. Then we have*

$$\lim_{n \rightarrow \infty} \|\mathcal{L}_n(f; \cdot, a) - f\|_\varphi = 0. \tag{3.4}$$

Proof From [9], we know that it is sufficient to verify the following three equations:

$$\lim_{n \rightarrow \infty} \|\mathcal{L}_n(e_m; \cdot, a) - e_m\|_\varphi = 0, \quad m = 0, 1, 2. \tag{3.5}$$

Since $\mathcal{L}_n(e_0; x, a) = 1$, the condition in (3.5) holds true for $m = 0$.

By Lemma 1, we have

$$\|\mathcal{L}_n(e_1; \cdot, a) - e_1\|_\varphi = \sup_{x \geq 0} \frac{1}{1+x^2} \left| x + \frac{1}{n} - x \right| \leq \frac{1}{n}.$$

Thus, $\lim_{n \rightarrow \infty} \|\mathcal{L}_n(e_1; \cdot, a) - e_1\|_\varphi = 0$. Similarly, we get

$$\begin{aligned} \|\mathcal{L}_n(e_2; \cdot, a) - e_2\|_\varphi &= \sup_{x \geq 0} \frac{1}{1+x^2} \left| x^2 + \frac{x}{n} \left(3 + \frac{1}{a-1} \right) + \frac{2}{n^2} - x^2 \right| \\ &\leq \frac{1}{n} \left(3 + \frac{1}{(a-1)} \right) + \frac{2}{n^2}, \end{aligned}$$

which implies that $\lim_{n \rightarrow \infty} \|\mathcal{L}_n(e_2; \cdot, a) - e_2\|_\varphi = 0$. This completes the proof. □

Next we give a theorem to approximate all functions in $C_\varphi[0, \infty)$. This type of result is discussed in [10] for locally integrable functions.

Theorem 6 For each $f \in C_\varphi[0, \infty)$ and $\beta > 0$, we have

$$\lim_{n \rightarrow \infty} \sup_{x \in [0, \infty)} \frac{|\mathcal{L}_n(f; x, a) - f(x)|}{(1+x^2)^{1+\beta}} = 0.$$

Proof For any fixed $x_0 > 0$,

$$\begin{aligned} \sup_{x \in [0, \infty)} \frac{|\mathcal{L}_n(f; x, a) - f(x)|}{(1+x^2)^{1+\beta}} &\leq \sup_{x \leq x_0} \frac{|\mathcal{L}_n(f; x, a) - f(x)|}{(1+x^2)^{1+\beta}} + \sup_{x \geq x_0} \frac{|\mathcal{L}_n(f; x, a) - f(x)|}{(1+x^2)^{1+\beta}} \\ &\leq \|\mathcal{L}_n(f) - f\|_{C[0, x_0]} + \|f\|_\varphi \sup_{x \geq x_0} \frac{|\mathcal{L}_n(1+t^2; x, a)|}{(1+x^2)^{1+\beta}} \\ &\quad + \sup_{x \geq x_0} \frac{|f(x)|}{(1+x^2)^{1+\beta}}, \\ &=: I_1 + I_2 + I_3, \text{ say.} \end{aligned} \tag{3.6}$$

Since $|f(x)| \leq \|f\|_\varphi(1+x^2)$, we have

$$I_3 = \sup_{x \geq x_0} \frac{|f(x)|}{(1+x^2)^{1+\beta}} \leq \sup_{x \geq x_0} \frac{\|f\|_\varphi}{(1+x^2)^\beta} \leq \frac{\|f\|_\varphi}{(1+x_0^2)^\beta}.$$

Let $\epsilon > 0$ be arbitrary. In view of Theorem 1, there exists $n_1 \in \mathbb{N}$ such that

$$|\mathcal{L}_n(1+t^2; x, a) - (1+x^2)| < \frac{\epsilon}{3\|f\|_\varphi}, \quad \forall n \geq n_1,$$

or

$$\mathcal{L}_n(1+t^2; x, a) < (1+x^2) + \frac{\epsilon}{3\|f\|_\varphi}, \forall n \geq n_1.$$

Hence,

$$\begin{aligned} \|f\|_\varphi \frac{|\mathcal{L}_n(1+t^2; x, a)|}{(1+x^2)^{1+\beta}} &< \frac{1}{(1+x^2)^{1+\beta}} \|f\|_\varphi \left((1+x^2) + \frac{\epsilon}{3\|f\|_\varphi} \right), \forall n \geq n_1 \\ &< \frac{\|f\|_\varphi}{(1+x^2)^\beta} + \frac{\epsilon}{3}, \forall n \geq n_1. \end{aligned} \quad (3.7)$$

$$\text{Hence, } \|f\|_\varphi \sup_{x \geq x_0} \frac{|\mathcal{L}_n(1+t^2; x, a)|}{(1+x^2)^{1+\beta}} < \frac{\|f\|_\varphi}{(1+x_0^2)^\beta} + \frac{\epsilon}{3}, \quad \forall n \geq n_1.$$

$$\text{Thus, } I_2 + I_3 < \frac{2\|f\|_\varphi}{(1+x_0^2)^\beta} + \frac{\epsilon}{3}, \quad \forall n \geq n_1.$$

$$\text{Now let us choose } x_0 \text{ to be so large that } \frac{\|f\|_\varphi}{(1+x_0^2)^\beta} < \frac{\epsilon}{6}.$$

Then,

$$I_2 + I_3 < \frac{2\epsilon}{3}, \quad \forall n \geq n_1. \quad (3.8)$$

By Theorem 4, there exists $n_2 \in \mathbb{N}$ such that

$$I_1 = \|\mathcal{L}_n(f) - f\|_{C[0, x_0]} < \frac{\epsilon}{3}, \quad \forall n \geq n_2. \quad (3.9)$$

Let $n_0 = \max(n_1, n_2)$. Then, combining (3.6)–(3.9):

$$\sup_{x \in [0, \infty)} \frac{|\mathcal{L}_n(f; x, a) - f(x)|}{(1+x^2)^{1+\beta}} < \epsilon, \quad \forall n \geq n_0.$$

This completes the proof. □

3.3. Rate of approximation

In this section we obtain the estimate of the rate of convergence for the operators \mathcal{L}_n for functions with derivatives of bounded variation. In recent years, several researchers have obtained results in this direction for different sequences of linear positive operators. We refer the reader to some of the related papers (cf. [1, 2, 6, 12, 14, 19], etc.).

Let $f \in DBV_\gamma[0, \infty)$, $\gamma > 0$ be the class of all functions defined on $[0, \infty)$, having a derivative of bounded variation on every finite subinterval of $[0, \infty)$ and $|f(t)| \leq Mt^\gamma$, $\forall t > 0$.

It turns out that for $f \in DBV_\gamma[0, \infty)$, we can write

$$f(x) = \int_0^x g(t)dt + f(0),$$

where $g(t)$ is a function of bounded variation on each finite subinterval of $[0, \infty)$.

Theorem 7 Let $f \in DBV_\gamma[0, \infty)$. Then for every $x \in (0, \infty)$, $r \in \mathbb{N}(2r > \gamma)$ and sufficiently large n , we have

$$\begin{aligned} |\mathcal{L}_n(f; x, a) - f(x)| &\leq \left| \frac{f'(x+) - f'(x-)}{2} \right| \left\{ \frac{\lambda(a)x}{n} \right\}^{1/2} + \frac{1}{n} \left| \frac{f'(x+) + f'(x-)}{2} \right| \\ &+ \frac{x}{\sqrt{n}} \bigvee_{x-\frac{x}{\sqrt{n}}}^{x+\frac{x}{\sqrt{n}}} (f'_x) + \frac{\lambda(a)}{n} \sum_{k=1}^{[\sqrt{n}]} \bigvee_{x-\frac{x}{k}}^{x+\frac{x}{k}} (f'_x) + |f'(x+)| \left\{ \frac{\lambda(a)x}{n} \right\}^{1/2} \\ &+ |f(2x) - f(x) - xf'(x+)| \frac{\lambda(a)}{nx} + \frac{C(\gamma, x, r, a)}{n^{\gamma/2}} + |f(x)| \frac{\lambda(a)}{nx}, \end{aligned}$$

where

$$f'_x(t) = \begin{cases} f'(t) - f'(x+), & x < t < \infty, \\ 0, & t = x, \\ f'(t) - f'(x-), & 0 \leq t < x. \end{cases}$$

$\bigvee_c^d(f'_x)$ is the total variation of f'_x on $[c, d]$ and $C(\gamma, x, r, a)$ is a constant depending only on γ, x, r , and a .

Proof By the hypothesis, we may write

$$\begin{aligned} f'(t) &= \frac{1}{2} \left(f'(x+) + f'(x-) \right) + f'_x(t) + \frac{1}{2} \left(f'(x+) - f'(x-) \right) \operatorname{sgn}(t - x) \\ &+ \delta_x(t) \left(f'(t) - \frac{1}{2} \left(f'(x+) + f'(x-) \right) \right), \end{aligned} \tag{3.10}$$

where

$$\delta_x(t) = \begin{cases} 1, & t = x, \\ 0, & t \neq x. \end{cases}$$

From equations (2.1) and (3.10), we have

$$\begin{aligned} \mathcal{L}_n(f; x, a) - f(x) &= \int_0^\infty \frac{\partial}{\partial t} \{ \mathcal{K}_n(x, t, a) \} f(t) dt - f(x) = \int_0^\infty (f(t) - f(x)) \frac{\partial}{\partial t} \{ \mathcal{K}_n(x, t, a) \} dt \\ &= \int_0^x (f(t) - f(x)) \frac{\partial}{\partial t} \{ \mathcal{K}_n(x, t, a) \} dt + \int_x^\infty (f(t) - f(x)) \frac{\partial}{\partial t} \{ \mathcal{K}_n(x, t, a) \} dt \\ &= - \int_0^x \left(\int_t^x f'(u) du \right) \frac{\partial}{\partial t} \{ \mathcal{K}_n(x, t, a) \} dt + \int_x^\infty \left(\int_x^t f'(u) du \right) \frac{\partial}{\partial t} \{ \mathcal{K}_n(x, t, a) \} dt \\ &=: -F_1(n, x, a) + F_2(n, x, a), \text{ say.} \end{aligned}$$

By using equation (3.10), we get

$$\begin{aligned} F_1(n, x, a) &= \int_0^x \left\{ \int_t^x \left(\frac{1}{2} \left(f'(x+) + f'(x-) \right) + f'_x(u) + \frac{1}{2} \left(f'(x+) - f'(x-) \right) \operatorname{sgn}(u - x) \right. \right. \\ &\left. \left. + \delta_x(u) \left(f'(u) - \frac{1}{2} \left(f'(x+) + f'(x-) \right) \right) \right) du \right\} \frac{\partial}{\partial t} \{ \mathcal{K}_n(x, t, a) \} dt. \end{aligned}$$

Since $\int_x^t (\delta_x(u))du = 0$, we have

$$\begin{aligned}
 F_1(n, x, a) &= \frac{1}{2} \left(f'(x+) + f'(x-) \right) \int_0^x (x-t) \frac{\partial}{\partial t} \{ \mathcal{K}_n(x, t, a) \} dt + \int_0^x \left(\int_t^x f'_x(u) du \right) \frac{\partial}{\partial t} \{ \mathcal{K}_n(x, t, a) \} dt \\
 &\quad - \frac{1}{2} \left(f'(x+) - f'(x-) \right) \int_0^x |x-t| \frac{\partial}{\partial t} \{ \mathcal{K}_n(x, t, a) \} dt.
 \end{aligned} \tag{3.11}$$

Proceeding similarly, we find that

$$\begin{aligned}
 F_2(n, x, a) &= \frac{1}{2} \left(f'(x+) + f'(x-) \right) \int_x^\infty (t-x) \frac{\partial}{\partial t} \{ \mathcal{K}_n(x, t, a) \} dt + \int_x^\infty \left(\int_x^t f'_x(u) du \right) \frac{\partial}{\partial t} \{ \mathcal{K}_n(x, t, a) \} dt \\
 &\quad + \frac{1}{2} \left(f'(x+) - f'(x-) \right) \int_x^\infty |t-x| \frac{\partial}{\partial t} \{ \mathcal{K}_n(x, t, a) \} dt.
 \end{aligned} \tag{3.12}$$

Combining (3.11) and (3.12), we get

$$\begin{aligned}
 \mathcal{L}_n(f; x, a) - f(x) &= \frac{1}{2} \left(f'(x+) + f'(x-) \right) \int_0^\infty (t-x) \frac{\partial}{\partial t} \{ \mathcal{K}_n(x, t, a) \} dt \\
 &\quad + \frac{1}{2} \left(f'(x+) - f'(x-) \right) \int_0^\infty |t-x| \frac{\partial}{\partial t} \{ \mathcal{K}_n(x, t, a) \} dt \\
 &\quad - \int_0^x \left(\int_t^x f'_x(u) du \right) \frac{\partial}{\partial t} \{ \mathcal{K}_n(x, t, a) \} dt + \int_x^\infty \left(\int_x^t f'_x(u) du \right) \frac{\partial}{\partial t} \{ \mathcal{K}_n(x, t, a) \} dt.
 \end{aligned}$$

Hence,

$$\begin{aligned}
 |\mathcal{L}_n(f; x, a) - f(x)| &\leq \left| \frac{f'(x+) + f'(x-)}{2} \right| |\mathcal{L}_n(t-x; x, a)| + \left| \frac{f'(x+) - f'(x-)}{2} \right| |\mathcal{L}_n(|t-x|; x, a)| \\
 &\quad + \left| \int_0^x \left(\int_t^x f'_x(u) du \right) \frac{\partial}{\partial t} \{ \mathcal{K}_n(x, t, a) \} dt \right| + \left| \int_x^\infty \left(\int_x^t f'_x(u) du \right) \frac{\partial}{\partial t} \{ \mathcal{K}_n(x, t, a) \} dt \right|.
 \end{aligned} \tag{3.13}$$

On an application of Lemma 4 and integration by parts, we obtain

$$\int_0^x \left(\int_t^x f'_x(u) du \right) \frac{\partial}{\partial t} \{ \mathcal{K}_n(x, t, a) \} dt = \int_0^x \left(\int_t^x f'_x(u) du \right) \frac{\partial}{\partial t} \vartheta_{n,a}(x, t) dt = \int_0^x f'_x(t) \vartheta_{n,a}(x, t) dt.$$

Thus,

$$\begin{aligned}
 \left| \int_0^x \left(\int_t^x f'_x(u) du \right) \frac{\partial}{\partial t} \{ \mathcal{K}_n(x, t, a) \} dt \right| &\leq \int_0^x |f'_x(t)| \vartheta_{n,a}(x, t) dt \\
 &\leq \int_0^{x-\frac{x}{\sqrt{n}}} |f'_x(t)| \vartheta_{n,a}(x, t) dt + \int_{x-\frac{x}{\sqrt{n}}}^x |f'_x(t)| \vartheta_{n,a}(x, t) dt.
 \end{aligned}$$

Since $f'_x(x) = 0$ and $\vartheta_{n,a}(x, t) \leq 1$, we get

$$\begin{aligned} \int_{x-\frac{x}{\sqrt{n}}}^x |f'_x(t)|\vartheta_{n,a}(x, t)dt &= \int_{x-\frac{x}{\sqrt{n}}}^x |f'_x(t) - f'_x(x)|\vartheta_{n,a}(x, t)dt \leq \int_{x-\frac{x}{\sqrt{n}}}^x \bigvee_t^x(f'_x)dt \\ &\leq \bigvee_{x-\frac{x}{\sqrt{n}}}^x(f'_x) \int_{x-\frac{x}{\sqrt{n}}}^x dt = \frac{x}{\sqrt{n}} \bigvee_{x-\frac{x}{\sqrt{n}}}^x(f'_x). \end{aligned}$$

Similarly, applying Lemma 4 and putting $t = x - \frac{x}{u}$, we get

$$\begin{aligned} \int_0^{x-\frac{x}{\sqrt{n}}} |f'_x(t)|\vartheta_{n,a}(x, t)dt &\leq \frac{\lambda(a)x}{n} \int_0^{x-\frac{x}{\sqrt{n}}} |f'_x(t)|\frac{dt}{(x-t)^2} \\ &\leq \frac{\lambda(a)x}{n} \int_0^{x-\frac{x}{\sqrt{n}}} \bigvee_t^x(f'_x) \frac{dt}{(x-t)^2} \\ &= \frac{\lambda(a)}{n} \int_1^{\sqrt{n}} \bigvee_{x-\frac{x}{u}}^x(f'_x)du \leq \frac{\lambda(a)}{n} \sum_{k=1}^{[\sqrt{n}]} \bigvee_{x-\frac{x}{k}}^x(f'_x). \end{aligned}$$

Consequently,

$$\left| \int_0^x \left(\int_t^x f'_x(u)du \right) \frac{\partial}{\partial t} \{ \mathcal{K}_n(x, t, a) \} dt \right| \leq \frac{x}{\sqrt{n}} \bigvee_{x-\frac{x}{\sqrt{n}}}^x(f'_x) + \frac{\lambda(a)}{n} \sum_{k=1}^{[\sqrt{n}]} \bigvee_{x-\frac{x}{k}}^x(f'_x). \tag{3.14}$$

Also, we have

$$\begin{aligned} &\left| \int_x^\infty \left(\int_x^t f'_x(u)du \right) \frac{\partial}{\partial t} \{ \mathcal{K}_n(x, t, a) \} dt \right| \\ &\leq \left| \int_x^{2x} \left(\int_x^t f'_x(u)du \right) \frac{\partial}{\partial t} (1 - \vartheta_{n,a}(x, t))dt \right| \\ &+ \left| \int_{2x}^\infty \left(\int_x^t f'_x(u)du \right) \frac{\partial}{\partial t} \{ \mathcal{K}_n(x, t, a) \} dt \right| \\ &\leq \left| \int_{2x}^\infty (f(t) - f(x)) \frac{\partial}{\partial t} \{ \mathcal{K}_n(x, t, a) \} dt \right| + |f'(x+)| \left| \int_{2x}^\infty (t-x) \frac{\partial}{\partial t} \{ \mathcal{K}_n(x, t, a) \} dt \right| \\ &+ \left| \int_x^{2x} f'_x(u)du \right| |1 - \vartheta_{n,a}(x, 2x)| + \int_x^{2x} |f'_x(t)|(1 - \vartheta_{n,a}(x, t))dt. \end{aligned} \tag{3.15}$$

We may write

$$\begin{aligned} \int_x^{2x} |f'_x(t)|(1 - \vartheta_{n,a}(x, t))dt &= \int_x^{x+\frac{x}{\sqrt{n}}} |f'_x(t)|(1 - \vartheta_{n,a}(x, t))dt + \int_{x+\frac{x}{\sqrt{n}}}^{2x} |f'_x(t)|(1 - \vartheta_{n,a}(x, t))dt \\ &=: I_1 + I_2 \text{ (say)}. \end{aligned} \tag{3.16}$$

Since $f'_x(x) = 0$ and $(1 - \vartheta_{n,a}(x, t)) \leq 1$, we have

$$\begin{aligned}
 I_1 &= \int_x^{x+\frac{x}{\sqrt{n}}} |f'_x(t) - f'_x(x)|(1 - \vartheta_{n,a}(x, t))dt \\
 &\leq \int_x^{x+\frac{x}{\sqrt{n}}} \bigvee_x^{x+\frac{x}{\sqrt{n}}}(f'_x)dt \\
 &= \frac{x}{\sqrt{n}} \bigvee_x^{x+\frac{x}{\sqrt{n}}}(f'_x).
 \end{aligned} \tag{3.17}$$

Next, we estimate I_2 . Applying Lemma 4 and putting $t = x + \frac{x}{u}$ we have

$$\begin{aligned}
 I_2 &= \int_{x+\frac{x}{\sqrt{n}}}^{2x} |f'_x(t)|(1 - \vartheta_{n,a}(x, t))dt \\
 &\leq \frac{\lambda(a)x}{n} \int_{x+\frac{x}{\sqrt{n}}}^{2x} |f'_x(t) - f'_x(x)| \frac{dt}{(x-t)^2} \\
 &\leq \frac{\lambda(a)x}{n} \int_{x+\frac{x}{\sqrt{n}}}^{2x} \bigvee_x^t(f'_x) \frac{dt}{(x-t)^2} \\
 &= \frac{\lambda(a)}{n} \int_1^{\sqrt{n}} \bigvee_x^{x+\frac{x}{u}}(f'_x)du \\
 &\leq \frac{\lambda(a)}{n} \sum_{k=1}^{[\sqrt{n}]} \int_k^{k+1} \bigvee_x^{x+\frac{x}{u}}(f'_x)du \\
 &\leq \frac{\lambda(a)}{n} \sum_{k=1}^{[\sqrt{n}]} \bigvee_x^{x+\frac{x}{k}}(f'_x).
 \end{aligned} \tag{3.18}$$

Putting the values of I_1 and I_2 in (3.16), we get

$$\int_x^{2x} |f'_x(t)|(1 - \vartheta_{n,a}(x, t))dt \leq \frac{x}{\sqrt{n}} \bigvee_x^{x+\frac{x}{\sqrt{n}}}(f'_x) + \frac{\lambda(a)}{n} \sum_{k=1}^{[\sqrt{n}]} \bigvee_x^{x+\frac{x}{k}}(f'_x). \tag{3.19}$$

In view of (2.2), (3.15), (3.19), and Lemma 4, we get

$$\begin{aligned}
 \left| \int_x^\infty \left(\int_x^t f'_x(u)du \right) \frac{\partial}{\partial t} \{ \mathcal{K}_n(x, t, a) \} dt \right| &\leq M \int_{2x}^\infty t^\gamma \frac{\partial}{\partial t} \{ \mathcal{K}_n(x, t, a) \} dt + |f(x)| \int_{2x}^\infty \frac{\partial}{\partial t} \{ \mathcal{K}_n(x, t, a) \} dt \\
 &\quad + |f'(x+)| \left\{ \frac{\lambda(a)x}{n} \right\}^{1/2} + \frac{\lambda(a)}{nx} |f(2x) - f(x) - xf'(x+)| \\
 &\quad + \frac{x}{\sqrt{n}} \bigvee_x^{x+\frac{x}{\sqrt{n}}}(f'_x) + \frac{\lambda(a)}{n} \sum_{k=1}^{[\sqrt{n}]} \bigvee_x^{x+\frac{x}{k}}(f'_x).
 \end{aligned} \tag{3.20}$$

Since $t \leq 2(t-x)$ and $x \leq t-x$ when $t \geq 2x$, applying Hölder's inequality, (2.2), and (2.3), we obtain

$$\begin{aligned}
 M \int_{2x}^{\infty} t^{\gamma} \frac{\partial}{\partial t} \{ \mathcal{K}_n(x, t, a) \} dt + |f(x)| \int_{2x}^{\infty} \frac{\partial}{\partial t} \{ \mathcal{K}_n(x, t, a) \} dt \\
 \leq M 2^{\gamma} \int_{2x}^{\infty} (t-x)^{\gamma} \frac{\partial}{\partial t} \{ \mathcal{K}_n(x, t, a) \} dt + \frac{|f(x)|}{x^2} \int_0^{\infty} \frac{\partial}{\partial t} \{ \mathcal{K}_n(x, t, a) \} (t-x)^2 dt \\
 \leq M 2^{\gamma} \left(\int_0^{\infty} (t-x)^{2r} \frac{\partial}{\partial t} \{ \mathcal{K}_n(x, t, a) \} dt \right)^{\gamma/2r} + |f(x)| \frac{\lambda(a)}{nx} \\
 \leq \frac{C(\gamma, x, r, a)}{n^{\gamma/2}} + |f(x)| \frac{\lambda(a)}{nx}.
 \end{aligned} \tag{3.21}$$

Combining (3.13), (3.14), (3.20), and (3.21), we get the required result. \square

Acknowledgments

The authors are extremely thankful to the referees and the editor for a very careful reading of the manuscript and making valuable suggestions and comments that greatly improved our paper. The first author is thankful to the "University Grants Commission" of India for financial support to carry out this research work.

References

- [1] Acar T, Gupta V, Aral A. Rate of convergence for generalized Szász operators. *Bull Math Sci* 2011; 1: 99–113.
- [2] Agrawal PN, Gupta V, Kumar AS, Kajla A. Generalized Baskakov-Szász type operators. *Appl Math Comput* 2014; 236: 311–324.
- [3] Altomare F, Montano MC, Leonessa V. On a generalization of Szász-Mirakjan-Kantorovich operators. *Results Math* 2013; 63: 837–863.
- [4] Aral A. A generalization of Szász-Mirakyan operators based on q-integers. *Math Comput Modelling* 2008; 47: 1052–1062.
- [5] Atakut Ç, Ispir N. Approximation by modified Szász-Mirakjan operators on weighted spaces. *Proc Indian Acad Sci Math* 2002; 112: 571–578.
- [6] Cárdenas-Morales D, Gupta V. Two families of Bernstein-Durrmeyer type operators. *Appl Math Comput* 2014; 248: 342–353.
- [7] Ciupa A. On a generalized Favard-Szász type operator. *Research Seminar on Numerical and Statistical Calculus, Univ. Babeş Bolyai Cluj-Napoca, Preprint* 1994; 1: 33–38.
- [8] Finta Z, Govil NK, Gupta V. Some results on modified Szász-Mirakjan operators. *J Math Anal Appl* 2007; 327: 1284–1296.
- [9] Gadjiev AD. On P. P. Korovkin type theorems. *Math Zametki* 1976; 20: 781–786.
- [10] Gadjiev AD, Efendiyev RO, Ibikli E. On Korovkin type theorem in the space of locally integrable functions. *Czech Math J* 2003; 53: 45–53.
- [11] Ismail MEH. *Classical and Quantum Orthogonal Polynomials in One Variable*. Cambridge, UK: Cambridge University Press, 2005.
- [12] Ispir N. Rate of convergence of generalized rational type Baskakov operators. *Math Comput Modelling* 2007; 46: 625–631.
- [13] Jakimovski A, Leviatan D. Generalized Szász operators for the approximation in the infinite interval. *Mathematica (Cluj)* 1969; 34: 97–103.

- [14] Karsli H. Rate of convergence of new gamma type operators for functions with derivatives of bounded variation. *Math Comput Modelling* 2007; 45: 617–624.
- [15] Kasana HS, Prasad G, Agrawal PN, Sahai A. Modified Szász operators. In: *Proceedings of the Conference on Mathematical Analysis and Its Applications*, Kuwait. Oxford, UK: Pergamon Press, 1985, 29–42.
- [16] Lenze B. On Lipschitz type maximal functions and their smoothness spaces. *Nederl Akad Indag Math* 1988; 50: 53–63.
- [17] Mazhar SM, Totik V. Approximation by modified Szász operators. *Acta Sci Math* 1985; 49: 257–269.
- [18] Özarslan MA, Aktuğlu H. Local approximation for certain King type operators. *Filomat* 2013; 27: 173–181.
- [19] Özarslan MA, Duman O, Kaanoğlu C. Rates of convergence of certain King-type operators for functions with derivative of bounded variation. *Math Comput Modelling* 2010; 52: 334–345.
- [20] Szász O. Generalization of S. Bernstein's polynomials to the infinite interval. *J Res Nat Bur Standards* 1950; 45: 239–245.
- [21] Varma S, Sucu S, İçöz G. Generalization of Szász operators involving Brenke type polynomials. *Comput Math Appl* 2012; 64: 121–127.
- [22] Varma S, Taşdelen F. Szász type operators involving Charlier polynomials. *Math Comput Modelling* 2012; 56: 118–122.