

1-1-2006

New Record of a Dinoflagellate Species *Corythodinium tessellatum* (Stein) Loeblich Jr. & Loeblich III from Turkish Coastal Waters of the North-eastern Mediterranean Sea

SEYFETTİN TAŞ

ERDOĞAN OKUŞ

TUFAN KORAY

Follow this and additional works at: <https://journals.tubitak.gov.tr/botany>



Part of the [Botany Commons](#)

Recommended Citation

TAŞ, SEYFETTİN; OKUŞ, ERDOĞAN; and KORAY, TUFAN (2006) "New Record of a Dinoflagellate Species *Corythodinium tessellatum* (Stein) Loeblich Jr. & Loeblich III from Turkish Coastal Waters of the North-eastern Mediterranean Sea," *Turkish Journal of Botany*. Vol. 30: No. 1, Article 5. Available at: <https://journals.tubitak.gov.tr/botany/vol30/iss1/5>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Botany by an authorized editor of TÜBİTAK Academic Journals. For more information, please contact academic.publications@tubitak.gov.tr.

New Record of a Dinoflagellate Species *Corythodinium tessellatum* (Stein) Loeblich Jr. & Loeblich III from Turkish Coastal Waters of the North-eastern Mediterranean Sea

Seyfettin TAŞ, Erdoğan OKUŞ

İstanbul University, Institute of Marine Sciences and Management, 34470, Vefa, İstanbul-TURKEY

Tufan KORAY

Ege University, Faculty of Fisheries, Department of Hydrobiology, 35100, Bornova, İzmir-TURKEY

Received: 17.06.2005

Accepted: 20.12.2005

Abstract: The dinoflagellate genus *Corythodinium* (Stein) Loeblich Jr. & Loeblich III and one species belonging to this genus, *Corythodinium tessellatum* (Stein) Loeblich Jr. & Loeblich III, are reported for the first time from the coastal waters of Turkey (north-eastern Mediterranean). The morphological characteristics of this species are described and information about its ecological distribution is given.

Key Words: *Corythodinium tessellatum*, dinoflagellate, phytoplankton, north-eastern Mediterranean, Datça-Bozburun Peninsula.

Kuzeydoğu Akdeniz'in Türkiye Kıyısı Sularında Bir Dinoflagellat Türü Olan *Corythodinium tessellatum* (Stein) Loeblich Jr. & Loeblich III İçin Yeni Kayıt

Özet: Dinoflagellatlara ait *Corythodinium* (Stein) Loeblich Jr. & Loeblich III cinsi ve bu cinse ait *Corythodinium tessellatum* (Stein) Loeblich Jr. & Loeblich III türü Türkiye'nin kıyısı sularında (kuzeydoğu Akdeniz) ilk kez rapor edilmiştir. Bu türün morfolojik özellikleri tanımlanmış ve ekolojik dağılımı hakkında bilgiler verilmiştir.

Anahtar Sözcükler: *Corythodinium tessellatum*, dinoflagellat, fitoplankton, kuzeydoğu Akdeniz, Datça-Bozburun Yarımadası.

Introduction

Dinoflagellates are organisms including some of the most important primary producers in marine and fresh water biotopes (Schnepf & Elbrächter, 1992). The oligotrophic conditions in the Mediterranean Sea could favour the richness of dinoflagellates, typical organisms of oligotrophic waters (Gómez, 2003). Dinoflagellates are well represented in the oligotrophic waters of the Mediterranean Sea (Gómez & Abi Saab, 2003).

Corythodinium tessellatum (Stein) Loeblich Jr. & Loeblich III is a rare species of the dinoflagellate group. This species is distributed in warm temperate to tropical waters and most records are from the Atlantic Ocean. A total of about 25 species from this genus have been recorded from the world's oceans (Tomas, 1997) and 10 species from the Mediterranean Sea (Gómez, 2003). In

this study, the genus *Corythodinium* (Stein) Loeblich Jr. & Loeblich III and a member of this genus, *C. tessellatum*, are reported from the north-eastern Mediterranean coast of Turkey. This record will contribute to the microplankton checklist of Turkish seas prepared by Koray (2001).

Materials and Methods

The samples of phytoplankton were collected from the coastal waters of Datça-Bozburun Peninsula (north-eastern Mediterranean; lat. 36° 32'N-36° 49'N, long. 27° 20'E-28° 19'E). The study area and the location of the sampling point are shown in Figure 1. The phytoplankton samples were collected with Niskin bottles from different depths (0.5, 5, 10 and 20 m) and with a 55 µm mesh standard plankton net from surface water with horizontal




Figure 1. The location of the sampling area.

towing. The samples were preserved in 4% formaldehyde (final concentration). A light microscope was used for the identification of the species and photos were taken by Nikon Diaphot 300 phase-contrast inverted microscope with a video camera. For identification and taxonomy, Delgado & Fortuna (1991), Dodge (1985), and Tomas (1997) were used.

Results and Discussion

The species *Corythodinium tessellatum* was very rare in specimens in the surface samples. This species were found in the spring (May 2003) in water and net sampling (temperature 24.7 °C, salinity 39.2 psu). The taxonomy of this genus and species are given as follows (Fensome et al., 1998; Gómez, 2003):

Division: Dinoflagellata (Bütschli, 1885) Fensome et al., 1993

Subdivision: Dinokaryota Fensome et al., 1993

Class: Dinophyceae Pascher 1914

Subclass: Peridiniphycidae Fensome et al., 1993

Order: Peridinales Haeckel 1894

Family: Oxytoxaceae Lindemann 1928

Genus: *Corythodinium* (Stein) Loeblich Jr. & Loeblich III 1966

The members of this genus are armoured and small to large biconical to elongate cells with sculptured theca. The cingulum is anterior to median and prominently excavated and with narrow lists. The epitheca is shorter than the hypotheca, and the anterior cingular list is almost the same diameter as the posterior cingular list (Tomas, 1997).

Corythodinium tessellatum (Stein) Loeblich Jr. & Loeblich III. Studies in Tropical Oceanography, 1966.

In this species, the cells are medium sized, the cingulum is premedian, the hypotheca exceeds the epitheca and ends in a thick antapical spine. The broad hypotheca is characteristic with longitudinal striae connecting evenly spaced, offsets horizontal striae (Tomas, 1997). A row of large pores is situated on the anterior side of the ridge, with smaller pores on the posterior (Dodge & Saunders, 1985). This species is distributed in warm temperate to tropical waters and most records are from the Atlantic Ocean (Tomas, 1997). Belonging to this genus 10 species are reported from all regions of the Mediterranean Sea (Gómez, 2003). The cell size is given in the literature as length 50 µm and

width 30 µm (Dodge 1985; Dodge & Saunders 1985). In this study the total length of the cells is 59-60 µm and width is 29-30 µm (Figure 2).

When the biogeographical distribution of the species is considered, the occurrence of it in the Turkish coastal waters of the north-eastern Mediterranean Sea is normal. However, in previous studies (Kıdeyş et al., 1989; Eker & Kıdeyş, 2000; Polat et al., 2000; Polat & Işık, 2002) this species may have been neglected, because of its rare occurrence. On the other hand, studies in this area (Datça-Bozburun Peninsula) are especially scarce. In addition, with the increasing marine traffic in the region, especially ballast waters, may have caused the transportation of this species from one site to another.




Figure 2. Light microscopy photograph of *Corythodinium tessellatum*.

References

- Delgado M & Fortuna JM (1991). Atlas de fitoplancton del Mar Mediterraneo. *Sci Mar* 55: 1-133.
- Dodge JD & Saunders RD (1985). A partial revision of the genus *Oxytoxum* (Dinophyceae) with the aid of scanning electron microscopy. *Bot Mar* 28: 99-122.
- Dodge JD (1985). Atlas of dinoflagellates. A scanning electron microscope survey. Botany Department, Royal Holloway & Bedford Colleges, (University of London) Egham, Surrey. 119 p. London: Farrand Press.
- Eker E & Kıdeyş AE (2000). Weekly variations in phytoplankton structure of a harbour in Mersin Bay (north-eastern Mediterranean). *Turk J Bot* 24: 13-24.
- Fensome RA, MacRae RA & Williams GL (1998). Dinoflaj. Geological survey of Canada open file. No.3653.
- Gómez F & Abi Saab M (2003). Records of *Ceratoperidinium* Margalef (Dinophyceae) from the Mediterranean Sea. *Vie Milleu* 53: 43-46.
- Gómez F (2003). Checklist of Mediterranean free-living dinoflagellates. *Bot Mar* 46: 215-242.
- Kıdeyş AE, Ünsal M & Bingel F (1989). Seasonal changes in net phytoplankton of Erdemli, north-eastern Mediterranean. *Turk J Bot* 13: 45-54.
- Koray T (2001). A check list for phytoplankton of Turkish Seas. *Journal of Fisheries and Aquatic Sciences* 18: 1-27.
- Polat S & Işık O (2002). Phytoplankton distribution, diversity and nutrients at the north-eastern Mediterranean coast of Turkey (Karataş-Adana). *Turk J Bot* 26: 77-86.
- Polat S, Sarıhan, E & Koray, T (2000). Seasonal changes in the phytoplankton of the north-eastern Mediterranean (Bay of İskenderun). *Turk J Bot* 24: 1-12.
- Schnepf E & Elbrächter M (1992). Nutritional strategies in dinoflagellates. A review with emphasis on cell biological aspects. *Europ J Protistol* 28: 3-24.
- Tomas CR (1997). Identifying marine phytoplankton. p. 858. New York: Academic Press.