

# Turkish Journal of Botany

---

Volume 29 | Number 6

Article 7

---

1-1-2005

## A New Record for the Flora of Turkey: *Iberis saxatilis* L. (Brasssicaceae)

TUNCAY DİRMENCI

FATİH SATIL

GÜLENDAM TÜMEN

Follow this and additional works at: <https://journals.tubitak.gov.tr/botany>



Part of the [Botany Commons](#)

---

### Recommended Citation

DİRMENCI, TUNCAY; SATIL, FATİH; and TÜMEN, GÜLENDAM (2005) "A New Record for the Flora of Turkey: *Iberis saxatilis* L. (Brasssicaceae)," *Turkish Journal of Botany*: Vol. 29: No. 6, Article 7. Available at: <https://journals.tubitak.gov.tr/botany/vol29/iss6/7>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Botany by an authorized editor of TÜBİTAK Academic Journals. For more information, please contact [academic.publications@tubitak.gov.tr](mailto:academic.publications@tubitak.gov.tr).

## A New Record for the Flora of Turkey: *Iberis saxatilis* L. (*Brassicaceae*)

Tuncay DIRMENCI

Balıkesir University, Necatibey Education Faculty, Department of Biology Education, 10100 Balıkesir - TURKEY

Fatih SATIL, Güldemir TÜMEN

Balıkesir University, Faculty of Science and Arts, Department of Biology, 10100 Balıkesir - TURKEY

Received: 10.09.2004

Accepted: 06.10.2005

**Abstract:** *Iberis saxatilis* L. is reported for the first time (B1 Balıkesir) from Turkey. A description, including an illustration of the plant parts, and a distribution map are presented.

**Key Words:** *Brassicaceae*, *Iberis saxatilis*, new record, Kazdağ, Balıkesir

### Türkiye Florası İçin Yeni Bir Kayıt: *Iberis saxatilis* L. (*Brassicaceae*)

**Özet:** *Iberis saxatilis* L. türü Türkiye'den (B1 Balıkesir) ilk defa kaydedilmiştir. Bu makalede türün betimi, morfolojik çizimleri ve yayılış haritası verilmiştir.

**Anahtar Sözcükler:** *Brassicaceae*, *Iberis saxatilis*, yeni kayıt, Kazdağ, Balıkesir

### Introduction

Some *Iberis* L. specimens were collected during an expedition to Balıkesir province as part of the Flowering Plants of Kazdağ National Park Project (TÜBİTAK-YDABAG-103Y104). We were not able to identify them using the Flora of Turkey (Hedge, 1965; Yıldırımlı, 2000). They were similar to *Iberis saxatilis* L. (Table 1) according to Flora D'Italia (Anzalone et al., 1982), Flora Europaea (Da Silva & Franco, 1993), the Mountain Flora of Greece (Franzen, 1986) and Flora Hellenica (Tan, 2002). After thorough examinations of all the specimens and the relevant literature, it was decided that this species was a new record for Turkey. The number of *Iberis* species known from Turkey is now 9.

*Iberis saxatilis* L., Cent. Pl. 2:23 (1756) (Figures 1, 2).

Perennial, procumbent, diffuse shrub with straight, flowering stems 2.5-10 cm, glabrous to puberulent below, densely retrorsely puberulent above. Leaves 3-9 x 1-1.5 mm, semi-cylindrical on not flowering shoots and below on flowering stems; flat on upper flowering stems, entire, linear, acute, mucronulate, glabrous on both surfaces, ciliate on margins when young, soon glabrous. Inflorescence 7-12 x 9-12 mm, corymbose in flower,

shortly elongating in fruit. Sepals, 1.5-2.5 mm, green to purplish, glabrous, white membranous-margined. Petals 3.5-6.5 mm (inner 3.5-4.5 mm, outer 5-6.5 mm), white to purplish-white, gradually narrowed into a short claw. Fruiting pedicel to 6 mm. Silicula 5-8 x 4-6 mm, ovate, broadly winged from the base, lobes rounded.

Type: Origin of type material not known (LINN 825/3 photo!).

Material Examined: [Turkey] B1 Balıkesir: Iter trojanum, Mt. Ida in marmor, mont. Szus-Dagh, 17.7.1883, P. Sintenis (photo!). Kazdağ (Ida), Nanekırı, limestone, 1500 m, 19.05.2004, T. Dirmenci (2516a!) & F. Satılı (GAZI, HUB, Yıldırımlı) ibid 05.06.2004, T. Dirmenci (2523!) & F. Satılı (EGE), ibid Zığındere valley, limestone, 1600 m, 19.05.2004, T. Dirmenci (2516b!) & F. Satılı (ANK), Susuz dağ-Kuş tepe, 1500 m, 02.07.2004, T. Dirmenci (2872b!) & F. Satılı (Hb. T. Dirmenci). Fels mit Rutschalde, Dép. Hérault: Pégairolles-de-Buéges-St.-Guillème-le-Désert 14 km sw Pégairolles-de-Boéges, 450 m, 4.4.1982, Max Nydegger 30006 (GAZI!); Kalkfels, Vaucluse: Carpentras-Sault-de-Vaucluse, Gorges de la Nesque, Belv. 734 m. 3.4.1980, Max Nydegger (GAZI!).

Table 1. Comparison of the diagnostic characteristics of *I. saxatilis* in Kazdağ with *I. saxatilis* in other Flora.

Characters	Kazdağı	Flora Hellenica	Flora Europaea	Flora D'Italia
Plant height	2.5-10 cm	2-10 cm	Up to 15 cm	5-15 cm
Leaflets	3-9 x 1-1.5 mm semi-cylindrical linear at first ciliate but soon glabrous acute, mucronulate	(2-)5-15 x 0.3-1.5 mm semi-terete linear ciliate, otherwise glabrous subacute	up to 20 x 1.5 mm semi-cylindrical linear at first ciliate but soon glabrous acut, mucronulate	10-15 x 1 mm - linear
Inflorescence	7-12 x 9-12 mm corymbose in flower shortly elongating in fruit	10-22 mm racemose	- corymbose in flower elongating in fruit	racemose
Pedicel	2-4 mm	2-4 mm	-	-
Sepals	1.5-2.5 mm green to purplish with scarious margin	2-5-3 mm purplish with scarious margin	- -	2.5 mm
Petals	inner 3.5-4.5 mm, outer 5-6.5 mm white to purplish-white	inner 3-6 mm, outer 7-10 mm white	- -	6-7 mm white
Style	0.7-1.2 mm	0.5-1.2 mm	-	
Fruiting pedicel	6 mm	-	-	
Silicula	5-8 x 4-6 mm ovate broadly winged from the base	4.5-7 x 4-6 mm broadly ovate wing 0.1-0.3 mm broad below	5-8(-9) x 4.5-6(-7) mm obovate broadly winged from the base	7 x 6 mm -

The recommended IUCN Threat Category (IUCN, 2001) is Endangered (EN). It was collected from 3 localities in Kazdağ. The Nanekırı population is in good condition, but the other 2 populations are poor, and the distribution area is less than 10 km<sup>2</sup> in Turkey. Therefore, the recommended threat category for this species is Endangered.

**Ecology:** It grows on limestone at 1500-1600 m together with e.g. *Achillea fraasii* Sch.Bip. subsp. *troiana* Asch. & Heimerl, *Asperula sintenisii* Asch. ex Bornm., *Paeonia mascula* (L.) Mill. subsp. *arietina* (Anders) Cullen & Heywood, *Aristolochia pallida* Willd., *Saxifraga sancta* Gris., *Iris suaveolens* Boiss. & Reut., *Viola tricolor* L., *Cerasus prostrata* (Lab.) Ser. var. *prostrata*, *Astragalus heldreichii* Boiss., *Astragalus idae* Grossh., *Minuartia garckeana* (Asch. & Sint. & ex Boiss.) Mattf., *Linum boissieri* Asch. & Sint. & ex Boiss., *Tulipa sylvestris* L., and *Muscari bourgaei* Baker.

## Discussion

*Iberis* is represented by 20 species in Europe (Da Silva & Franco, 1993), by 6 species in Flora Hellenica (Tan, 2002), and by 8 species in the Flora of Turkey (Hedge, 1965; Yıldırımlı, 2000, 2001). According to Franzen (1986) and Tan (2002), *I. saxatilis* was collected from Kazdağ (Ida) by Sintenis in 1883. This specimen was not seen by Hedge (1965).

*I. saxatilis* is allied to *I. sempervirens* L. and *I. runemarkii* Greuter & Bardet (Syn: *Iberis arbuscula* Runemark), which are also woody at the base. It can be easily distinguished from *I. sempervirens* by its short flowering stems, 2.5-10 cm (not 15-30 cm), semi-cylindrical leaves (not oblong to oblong-spathulate), length and width of leaves, 3-9 x 1-1.5 mm (not 8-30 x 2-6 mm), outer petals to 6.5 mm (not 7-11 mm), and small fruit, 5-8 mm (not 8-10 mm). It differs from *I. runemarkii* in its semi-cylindrical leaves (not spathulate),




Figure 1. The plant parts of *Iberis saxatilis* L.: A-Habit, B-Leaves (a-in fruit period, b-in flowering period, c-in early flowering period), C-Flower, D-Calyx, E-Fruit.

length and width of leaves, 3-9 x 1-1.5 mm (not 10-30 x 2-5 mm), leaf margins entire (not 1-2 small teeth on each side), fruiting pedicel up to 6 mm (not 5-9 mm), sepals 1.5-2.5 mm (not c. 6 mm), outer petals to 6.5 mm (not up to 15 mm), and small fruit, 5-8 mm (not 8-10 mm).

#### Key to Related *Iberis* L. Species

1. Flowering stems 15-30 cm, leaves 8-30 x 2-6 mm, outer petals 7-15 mm
2. Leaves entire, sepals 2-3 mm, outer petals up to 11 mm      *I. sempervirens*

2. Leaves toothed at apex, sepals c. 6 mm, outer petals up to 15 mm      *I. runemarkii*

1. Flowering stems 2.5-10 cm, leaves 3-9 x 1-1.5 mm, sepals 1.5-2.5 mm, outer petals up to 6.5 mm.      *I. saxatilis*

*I. saxatilis* is widely distributed in Europe (Greece, Italy, Spain, Romania, France, Switzerland, Yugoslavia, Crimea and N. Africa). It was divided into 2 subspecies in Flora Europaea: *I. saxatilis* subsp. *saxatilis* and subsp. *cineraria* (Poiret) P.W.Ball & Heywood. *I. saxatilis* subsp.




Figure 2. Distribution map of (■) *Iberis saxatilis* L. in Turkey.

*saxatilis* is apparently the Turkish plant. According to Flora Europaea, it differs from subsp. *cineraria* in its stem up to 15 cm (not 40 cm), glabrous to puberulent (not velutinous), leaves up to 1.5 mm wide (not 2 mm), and silicula with a shallow notch (not acute).

#### Key to related subspecies of *Iberis saxatilis* L.

Flowering stems up to 10 cm (Turkish plants), leaves up to 1.5 mm wide, silicula with a shallow notch  
subsp. *saxatilis*

Flowering stems up to 40 cm, leaves up to 2 mm wide, notch of silicula acute  
subsp. *cineraria*

#### References

- Anzalone B, Becherer A, Ehrendorfer F, Merxmüller H, Metlesics H, Montelucci G, Rasetti F, Reichstein T & Segelberg I (1982). *Flora D'Italia* Vol. 1, Sandro Pigatti, Edagricole.
- Da Silva ARP & Franco JA (1993). *Iberis* L. In: Tutin TG, Heywood VH, Walters SM & Webb DA, (ed. 2). *Flora Europaea* (ed 2). Vol. 1, 390-393. Cambridge: Cambridge University Press.
- Franzen R (1986). *Iberis* L. In: Strid A. (ed). *Mountain Flora of Greece*. Vol. 1, 331-334. Cambridge: Cambridge University Press.
- Hedge IC (1965). *Iberis* L. In: Davis PH, (ed.). *Flora of Turkey and the East Aegean Islands*. Vol. 1, 309-312. Edinburgh: Edinburgh University Press.
- IUCN Species Survival Commission (2001). *IUCN red list categories and criteria*. Approved by the 51st meeting of the IUCN Council, Version 3.1 Switzerland: Gland.
- Tan K (2002). *Iberis* L. In: Strid A & Tan K. (ed.). *Flora Hellenica*. Vol. 2, 265-268. Koeltz Scientific Books.
- Yıldırımlı Ş (2000). *Iberis* L. In: Güner A, Özhatay N, Ekim T & Başer KHC. *Flora of Turkey and the East Aegean Islands* (Supplement II). Vol. 11, 31. Edinburgh: Edinburgh University Press.
- Yıldırımlı Ş (2001). The chorology of the Turkish species of *Brassicaceae*, *Buddlejaceae* and *Buxaceae* families. *Ot Sist Bot Derg.* 8:141-171.

#### Acknowledgements

We would like to thank Prof. Dr. Bayram Yıldız for the useful discussion, the curators of LD and G, who sent us photos of *I. saxatilis* and *I. runemarkii*, and the curators of HUB and GAZI, who gave us permission to examine the specimens.

This research was funded by TÜBİTAK-YDABAG (103Y104).