

1-1-2015

## Real hypersurfaces in complex two-plane Grassmannians whose shape operator is recurrent for the generalized Tanaka-Webster connection

JUAN DE DIOS PEREZ

YOUNG JIN SUH

CHANGHWA WOO

Follow this and additional works at: <https://journals.tubitak.gov.tr/math>



Part of the [Mathematics Commons](#)

---

### Recommended Citation

PEREZ, JUAN DE DIOS; SUH, YOUNG JIN; and WOO, CHANGHWA (2015) "Real hypersurfaces in complex two-plane Grassmannians whose shape operator is recurrent for the generalized Tanaka-Webster connection," *Turkish Journal of Mathematics*: Vol. 39: No. 3, Article 2. <https://doi.org/10.3906/mat-1403-74>

Available at: <https://journals.tubitak.gov.tr/math/vol39/iss3/2>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Mathematics by an authorized editor of TÜBİTAK Academic Journals. For more information, please contact [academic.publications@tubitak.gov.tr](mailto:academic.publications@tubitak.gov.tr).

## Real hypersurfaces in complex two-plane Grassmannians whose shape operator is recurrent for the generalized Tanaka–Webster connection

Juan de Dios PÉREZ<sup>1,\*</sup>, Young Jin SUH<sup>2</sup>, Changhwa WOO<sup>2</sup>

<sup>1</sup>Department of Geometry and Topology, University of Granada, Granada, Spain

<sup>2</sup>Department of Mathematics, Kyungpook National University, Daegu, Republic of Korea

Received: 26.03.2014

Accepted/Published Online: 15.01.2015

Printed: 29.05.2015

**Abstract:** We prove the non-existence of Hopf real hypersurfaces in complex two-plane Grassmannians whose shape operator  $A$  is generalized Tanaka–Webster recurrent if the principal curvature of the structure vector field is not equal to  $\text{trace}(A)$ .

**Key words:** Real hypersurfaces, complex two-plane Grassmannians, Hopf hypersurface, generalized Tanaka–Webster connection, recurrent shape operator

### 1. Introduction

The generalized Tanaka–Webster connection (from now on,  $g$ -Tanaka–Webster connection) for contact metric manifolds was introduced by Tanno ([13]) as a generalization of the connection defined by Tanaka in [12] and, independently, by Webster in [14]. This connection coincides with the Tanaka–Webster connection if the associated CR-structure is integrable. The Tanaka–Webster connection is defined as a canonical affine connection on a non-degenerate, pseudo-Hermitian CR-manifold. A real hypersurface  $M$  in a Kähler manifold has an (integrable) CR-structure associated with the almost contact structure  $(\phi, \xi, \eta, g)$  induced on  $M$  by the Kähler structure, but, in general, this CR-structure is not guaranteed to be pseudo-Hermitian. Cho [4] and Tanno [13] defined the  $g$ -Tanaka–Webster connection for a real hypersurface of a Kähler manifold by

$$\hat{\nabla}_X^{(k)} Y = \nabla_X Y + g(\phi AX, Y)\xi - \eta(Y)\phi AX - k\eta(X)\phi Y \quad (1.1)$$

for any  $X, Y$  tangent to  $M$ , where  $\nabla$  denotes the Levi-Civita connection on  $M$ ,  $A$  is the shape operator on  $M$  and  $k$  is a non-zero real number. In particular, if the real hypersurface satisfies  $A\phi + \phi A = 2k\phi$ , then the  $g$ -Tanaka–Webster connection  $\hat{\nabla}^{(k)}$  coincides with the Tanaka–Webster connection (see [4]).

Now let us denote by  $G_2(\mathbb{C}\mathbb{C}^{m+2})$  the set of all complex 2-dimensional linear subspaces in  $\mathbb{C}\mathbb{C}^{m+2}$ . This Riemannian symmetric space has a remarkable geometric structure. It is known to be the unique compact irreducible Riemannian symmetric space equipped with both a Kähler structure  $J$  and a quaternionic Kähler structure  $\mathfrak{J}$  not containing  $J$  (see Berndt and Suh [2]). In other words,  $G_2(\mathbb{C}\mathbb{C}^{m+2})$  is the unique compact, irreducible Kähler, quaternionic Kähler manifold, which is not a hyper-Kähler manifold.

\*Correspondence: jdperez@ugr.es

2010 AMS Mathematics Subject Classification: Primary 53C40; Secondary 53C15.

The first author is partially supported by MCT-FEDER Grant MTM2010-18099, the second author by Grant Proj. No. NRF-2012-R1A2A2A-01043023, and the third author by Fostering Core Leaders No. NRF-2013H1A8A1004325.

Let  $M$  be a real hypersurface in  $G_2(\mathbb{C}\mathbb{C}^{m+2})$  and  $N$  a local normal unit vector field on  $M$ . Let also  $A$  be the shape operator of  $M$  associated to  $N$ . The almost contact structure vector field  $\xi = -JN$  is said to be a Reeb vector field. Moreover, if  $\{J_1, J_2, J_3\}$  is a local basis of  $\mathfrak{J}$ , we define  $\xi_i = -J_i N$ ,  $i = 1, 2, 3$ . We will call  $\mathbb{D}^\perp = \text{Span}\{\xi_1, \xi_2, \xi_3\}$ . Its orthogonal complement in  $TM$  will be denoted by  $\mathbb{D}$ .

Berndt and Suh, [2] proved that for a connected hypersurface  $M$  in  $G_2(\mathbb{C}\mathbb{C}^{m+2})$ ,  $m \geq 3$ , both  $\text{Span}\{\xi\}$  and  $\mathbb{D}^\perp$  are invariant under the shape operator  $A$  if and only if either (A)  $M$  is an open part of a tube around a totally geodesic  $G_2(\mathbb{C}\mathbb{C}^{m+1})$  in  $G_2(\mathbb{C}\mathbb{C}^{m+2})$ , or (B)  $m$  is even, say  $m = 2n$  and  $M$  is an open part of a tube around a totally geodesic  $\mathbb{H}P^n$  in  $G_2(\mathbb{C}\mathbb{C}^{m+2})$ . Both types of real hypersurfaces have constant principal curvatures.

The Reeb vector field  $\xi$  is said to be Hopf if it is invariant under the shape operator  $A$ . The 1-dimensional foliation of  $M$  by the integral manifolds of the Reeb vector field  $\xi$  is said to be a Hopf foliation of  $M$ . We say that  $M$  is a Hopf hypersurface in  $G_2(\mathbb{C}\mathbb{C}^{m+2})$  if and only if the Hopf foliation of  $M$  is totally geodesic. This is equivalent to the fact that the Reeb vector field is Hopf.

If the shape operator  $A$  of  $M$  satisfies  $(\nabla_X A)Y = 0$  for any vector fields  $X, Y$  tangent to  $M$ , the shape operator is said to be parallel with respect to the Levi-Civita connection. Suh [10] proved the non-existence of real hypersurfaces in  $G_2(\mathbb{C}\mathbb{C}^{m+2})$  with parallel shape operator with respect to the Levi-Civita connection.

On the other hand, Kobayashi and Nomizu [7] introduced the notion of recurrent tensor field of type (r,s) on a manifold  $M$  with a linear connection  $\nabla$ . A non-zero tensor field  $K$  of type (r,s) on  $M$  is said to be recurrent if there exists a 1-form  $\omega$  on  $M$  such that  $\nabla K = K \otimes \omega$ .

Suh [11] proved the non-existence of real hypersurfaces in  $G_2(\mathbb{C}\mathbb{C}^{m+2})$  with recurrent shape operator with respect to the Levi-Civita connection if  $\mathbb{D}$  (respectively,  $\mathbb{D}^\perp$ ) is invariant by the shape operator. Kim et al. [6] showed that this last condition is superfluous.

Jeong et al. [5] considered real hypersurfaces in  $G_2(\mathbb{C}\mathbb{C}^{m+2})$  whose shape operator is parallel with respect to the g-Tanaka–Webster connection, that is,  $(\hat{\nabla}_X^{(k)} A)Y = 0$  for any  $X, Y$  tangent to  $M$  and proved that there do not exist Hopf real hypersurfaces in  $G_2(\mathbb{C}\mathbb{C}^{m+2})$ ,  $m \geq 3$ , with parallel shape operator with respect to the g-Tanaka–Webster connection  $\hat{\nabla}^{(k)}$  if  $\alpha = g(A\xi, \xi) \neq 2k$ .

This paper is devoted to the study of real hypersurfaces in complex two-plane Grassmannians whose shape operator is recurrent with respect to the g-Tanaka–Webster connection  $\hat{\nabla}^{(k)}$ . That is, there exists a 1-form  $\omega$  on  $M$  such that  $(\hat{\nabla}_X^{(k)} A)Y = \omega(X)AY$  for any  $X, Y$  tangent to  $M$ . We will call  $h = \text{trace}(A)$ . Notice that if  $\omega \equiv 0$ , the shape operator should be parallel with respect to the g-Tanaka–Webster connection. Thus we will suppose that the 1-form  $\omega$  does not vanish. We will prove the following

**Main Theorem** *There do not exist Hopf real hypersurfaces in  $G_2(\mathbb{C}\mathbb{C}^{m+2})$ ,  $m \geq 3$ , whose shape operator is recurrent with respect to the g-Tanaka–Webster connection if  $\alpha = g(A\xi, \xi) \neq h$ , where  $h = \text{Tr}A$ .*

## 2. Preliminaries

For the study of the Riemannian geometry of  $G_2(\mathbb{C}\mathbb{C}^{m+2})$  see [1]. All the notations we will use from now on are those in [2] and [3]. We will suppose that the metric  $g$  of  $G_2(\mathbb{C}\mathbb{C}^{m+2})$  is normalized for the maximal sectional curvature of the manifold to be eight. Then the Riemannian curvature tensor  $\bar{R}$  of  $G_2(\mathbb{C}\mathbb{C}^{m+2})$  is locally given

by

$$\begin{aligned} \bar{R}(X, Y)Z &= g(Y, Z)X - g(X, Z)Y + g(JY, Z)JX - g(JX, Z)JY - 2g(JX, Y)JZ \\ &+ \sum_{\nu=1}^3 \{g(J_\nu Y, Z)J_\nu X - g(J_\nu X, Z)J_\nu Y - 2g(J_\nu X, Y)J_\nu Z\} \\ &+ \sum_{\nu=1}^3 \{g(J_\nu JY, Z)J_\nu JX - g(J_\nu JX, Z)J_\nu JY\}, \end{aligned} \tag{2.1}$$

where  $J_1, J_2, J_3$  is any canonical local basis of  $\mathfrak{J}$ .

Let  $M$  be a real hypersurface of  $G_2(\mathbb{C}\mathbb{C}^{m+2})$ , that is, a submanifold of  $G_2(\mathbb{C}\mathbb{C}^{m+2})$  with real codimension one. The induced Riemannian metric on  $M$  will also be denoted by  $g$ , and  $\nabla$  denotes the Riemannian connection of  $(M, g)$ . Let  $N$  be a local unit normal field of  $M$  and  $A$  the shape operator of  $M$  with respect to  $N$ . The Kähler structure  $J$  of  $G_2(\mathbb{C}\mathbb{C}^{m+2})$  induces on  $M$  an almost contact metric structure  $(\phi, \xi, \eta, g)$ . Furthermore, let  $J_1, J_2, J_3$  be a canonical local basis of  $\mathfrak{J}$ . Then each  $J_\nu$  induces an almost contact metric structure  $(\phi_\nu, \xi_\nu, \eta_\nu, g)$  on  $M$ .

Since  $\mathfrak{J}$  is parallel with respect to the Riemannian connection  $\bar{\nabla}$  of  $(G_2(\mathbb{C}\mathbb{C}^{m+2}), g)$ , for any canonical local basis  $J_1, J_2, J_3$  of  $\mathfrak{J}$  there exist three local 1-forms  $q_1, q_2, q_3$  such that

$$\bar{\nabla}_X J_\nu = q_{\nu+2}(X)J_{\nu+1} - q_{\nu+1}(X)J_{\nu+2} \tag{2.2}$$

for any  $X$  tangent to  $G_2(\mathbb{C}\mathbb{C}^{m+2})$ , where subindices are taken modulo 3.

From the expression of the curvature tensor of  $G_2(\mathbb{C}\mathbb{C}^{m+2})$  the Gauss equation is given by

$$\begin{aligned} R(X, Y)Z &= g(Y, Z)X - g(X, Z)Y + g(\phi Y, Z)\phi X - g(\phi X, Z)\phi Y - 2g(\phi X, Y)\phi Z \\ &+ \sum_{\nu=1}^3 \{g(\phi_\nu Y, Z)\phi_\nu X - g(\phi_\nu X, Z)\phi_\nu Y - 2g(\phi_\nu X, Y)\phi_\nu Z\} \\ &+ \sum_{\nu=1}^3 \{g(\phi_\nu \phi Y, Z)\phi_\nu \phi X - g(\phi_\nu \phi X, Z)\phi_\nu \phi Y\} \\ &- \sum_{\nu=1}^3 \{\eta(Y)\eta_\nu(Z)\phi_\nu \phi X - \eta(X)\eta_\nu(Z)\phi_\nu \phi Y\} \\ &- \sum_{\nu=1}^3 \{\eta(X)g(\phi_\nu \phi Y, Z) - \eta(Y)g(\phi_\nu \phi X, Z)\}\xi_\nu \\ &+ g(AY, Z)ZX - g(AX, Z)AY \end{aligned} \tag{2.3}$$

for any  $X, Y, Z$  tangent to  $M$ . The Codazzi equation is also given by

$$\begin{aligned}
 (\nabla_X A)Y - (\nabla_Y A)X &= \eta(X)\phi Y - \eta(Y)\phi X - 2g(\phi X, Y)\xi \\
 &+ \sum_{\nu=1}^3 \{ \eta_\nu(X)\phi_\nu Y - \eta_\nu(Y)\phi_\nu X - 2g(\phi_\nu X, Y)\xi_\nu \} \\
 &+ \sum_{\nu=1}^3 \{ \eta_\nu(\phi X)\phi_\nu \phi Y - \eta_\nu(\phi Y)\phi_\nu \phi X \} \\
 &+ \sum_{\nu=1}^3 \{ \eta(X)\eta_\nu(\phi Y) - \eta(Y)\eta_\nu(\phi X) \} \xi_\nu
 \end{aligned}
 \tag{2.4}$$

for any  $X, Y$  tangent to  $M$ . The structures of  $G_2(\mathbb{C}\mathbb{C}^{m+2})$  give the following

$$(\nabla_X \phi)Y = \eta(Y)AX - g(AX, Y)\xi, \tag{2.5}$$

$$\nabla_X \xi = \phi AX, \tag{2.6}$$

$$\nabla_X \xi_\nu = q_{\nu+2}(X)\xi_{\nu+1} - q_{\nu+1}(X)\xi_{\nu+2} + \phi_\nu AX, \tag{2.7}$$

$$(\nabla_X \phi_\nu)Y = -q_{\nu+1}(X)\phi_{\nu+2}Y + q_{\nu+2}(X)\phi_{\nu+1}Y + \eta_\nu(Y)AX - g(AX, Y)\xi_\nu. \tag{2.8}$$

A real hypersurface of type (A) has three (if  $r = \frac{\pi}{2\sqrt{8}}$ ) or four (otherwise) distinct principal curvatures  $\alpha = \sqrt{8} \cot(\sqrt{8}r)$ ,  $\beta = \sqrt{2} \cot(\sqrt{2}r)$ ,  $\lambda = -\sqrt{2} \tan(\sqrt{2}r)$ ,  $\mu = 0$ , for some radius  $r \in (0, \frac{\pi}{\sqrt{8}})$ , with corresponding multiplicities  $m(\alpha) = 1$ ,  $m(\beta) = 2$ ,  $m(\lambda) = m(\mu) = 2m - 2$ . The corresponding eigenspaces can be seen in [2].

A real hypersurface of type (B) has five distinct principal curvatures  $\alpha = -2 \tan(2r)$ ,  $\beta = 2 \cot(2r)$ ,  $\gamma = 0$ ,  $\lambda = \cot(r)$ ,  $\mu = -\tan(r)$ , for some  $r \in (0, \frac{\pi}{4})$ , with corresponding multiplicities  $m(\alpha) = 1$ ,  $m(\beta) = 3 = m(\gamma)$ ,  $m(\lambda) = 4m - 4 = m(\mu)$ . For the corresponding eigenspaces see [2].

In the following we will need the following Proposition, [2],

**Proposition 2.1** *Let  $M$  be a Hopf real hypersurface in  $G_2(\mathbb{C}\mathbb{C}^{m+2})$ ,  $m \geq 3$ , such that  $A\xi = \alpha\xi$ . Then  $Y(\alpha) = \xi(\alpha)\eta(Y) - 4 \sum_{\nu=1}^3 \eta_\nu(\xi)\eta_\nu(\phi Y)$  for any  $Y$  tangent to  $M$ .*

and the following Theorem, [8],

**Theorem 2.2** *Let  $M$  be a connected orientable Hopf real hypersurface in  $G_2(\mathbb{C}\mathbb{C}^{m+2})$ ,  $m \geq 3$ . Then the Reeb vector field  $\xi$  belongs to the distribution  $\mathbb{D}$  if and only if  $m$  is locally congruent to an open part of a tube around a totally geodesic  $\mathbb{H}P^n$  in  $G_2(\mathbb{C}\mathbb{C}^{m+2})$ , where  $m = 2n$ .*

### 3. Proof of main theorem

As we suppose that  $(\hat{\nabla}_X^{(k)} A)Y = \omega(X)AY$  for any  $X, Y$  tangent to  $M$ , from (1.1) we get

$$\begin{aligned}
 (\nabla_X A)Y &= -g(\phi AX, AY)\xi + \eta(AY)\phi AX + k\eta(X)\phi AY + g(\phi AX, Y)A\xi \\
 &- \eta(Y)A\phi AX - k\eta(X)A\phi Y + \omega(X)AY
 \end{aligned}
 \tag{3.1}$$

for any  $X, Y$  tangent to  $M$ . As  $A\xi = \alpha\xi$ , taking  $Y = \xi$  in (3.1) we obtain  $\nabla_X \alpha\xi = \alpha\phi AX + \alpha\omega(X)\xi$ . That is,  $X(\alpha)\xi + \alpha\phi AX = \alpha\omega(X)\xi + \alpha\phi AX$ . Thus

$$X(\alpha) = \alpha\omega(X) \tag{3.2}$$

for any  $X$  tangent to  $M$ .

**Proposition 3.1** *Let  $M$  be a Hopf real hypersurface in  $G_2(\mathbb{C}\mathbb{C}^{m+2})$ ,  $m \geq 3$ , whose shape operator is recurrent with respect to the  $g$ -Tanaka–Webster connection. If  $\alpha \neq h$ , either  $\xi \in \mathbb{D}$  or  $\xi \in \mathbb{D}^\perp$ .*

**Proof** From [9] we know that if  $\alpha = 0$ , a Hopf real hypersurface in  $G_2(\mathbb{C}\mathbb{C}^{m+2})$  satisfies either  $\xi \in \mathbb{D}$  or  $\xi \in \mathbb{D}^\perp$ . Therefore we suppose that  $\alpha \neq 0$ .

We can write  $\xi = \eta(X_0)X_0 + \eta(\xi_1)\xi_1$  for a certain  $X_0 \in \mathbb{D}$ . If  $\eta(X_0) = 0$  (respectively,  $\eta(\xi_1) = 0$ ),  $\xi \in \mathbb{D}^\perp$  (respectively,  $\xi \in \mathbb{D}$ ). Therefore we suppose  $\eta(X_0)\eta(\xi_1) \neq 0$ . From (3.1) and the Codazzi equation we have

$$\begin{aligned} & \eta(X)\phi Y - \eta(Y)\phi X - 2g(\phi X, Y)\xi + \sum_{\nu=1}^3 \{ \eta_\nu(X)\phi_\nu Y - \eta_\nu(Y)\phi_\nu X - 2g(\phi_\nu(X, Y)\xi_\nu \} \\ & + \sum_{\nu=1}^3 \{ \eta_\nu(\phi X)\phi_\nu \phi Y - \eta_\nu(\phi Y)\phi_\nu \phi X \} + \sum_{\nu=1}^3 \{ \eta(X)\eta_\nu(\phi Y) - \eta(Y)\eta_\nu(\phi X) \} \xi_\nu \\ & = -2g(A\phi AX, Y)\xi - \eta(AX)\phi AY + \eta(A Y)\phi AX - k\eta(Y)\phi AX + k\eta(X)\phi AY \\ & + g((\phi A + A\phi)X, Y)A\xi + \eta(X)A\phi AY - \eta(Y)A\phi AX \\ & + k\eta(Y)A\phi X - k\eta(X)A\phi Y + \omega(X)AY - \omega(Y)AX \end{aligned} \tag{3.3}$$

for any  $X, Y$  tangent to  $M$ . Taking  $X = \xi$  in (3.3) we get

$$\begin{aligned} & \phi Y + \eta_1(\xi)\phi_1 Y - \sum_{\nu=1}^3 \eta_\nu(Y)\phi_\nu \xi - 2 \sum_{\nu=1}^3 g(\phi_\nu \xi, Y)\xi_\nu + \sum_{\nu=1}^3 \eta_\nu(\phi Y)\xi_\nu \\ & = -\alpha\phi AY + k\phi AY + A\phi AY - kA\phi Y + \omega(\xi)AY - \alpha\omega(Y)\xi \end{aligned} \tag{3.4}$$

for any  $Y$  tangent to  $M$ . Taking the scalar product of (3.4) and  $\xi$  we obtain

$$-4\eta_1(\xi)g(\phi\xi_1, Y) = \alpha\omega(\xi)\eta(Y) - \alpha\omega(Y) \tag{3.5}$$

for any  $Y$  tangent to  $M$ . As  $\alpha \neq 0$ , from (3.2) and (3.5)

$$grad(\alpha) = \alpha\omega(\xi)\xi + 4\eta_1(\xi)\phi_1\xi. \tag{3.6}$$

Let  $\{E_i\}_{i=1, \dots, 4m-1}$  be an orthonormal basis of eigenvectors of  $M$ , and suppose  $AE_i = \lambda_i E_i$ ,  $i = 1, \dots, 4m - 1$ . From (3.1) we have

$$\begin{aligned} \sum_{i=1}^{4m-1} g((\nabla_X A)E_i, E_i) &= - \sum_{i=1}^{4m-1} g(\phi AX, AE_i)g(\xi, E_i) \\ &\quad + \sum_{i=1}^{4m-1} \eta(AE_i)g(\phi AX, E_i) + k\eta(X) \sum_{i=1}^{4m-1} g(\phi AE_i, E_i) \\ &\quad + \sum_{i=1}^{4m-1} g(\phi AX, E_i)g(A\xi, E_i) - \sum_{i=1}^{4m-1} g(\xi, E_i)g(A\phi AX, E_i) \\ &\quad - k\eta(X) \sum_{i=1}^{4m-1} g(A\phi E_i, E_i) + \omega(X) \sum_{i=1}^{4m-1} g(AE_i, E_i) \\ &= h\omega(X), \end{aligned} \tag{3.7}$$

because the other terms are clearly null. This yields  $\sum_{i=1}^{4m-1} g(\nabla_X \lambda_i E_i - A\nabla_X E_i, E_i) = \sum_{i=1}^{4m-1} X(\lambda_i) = h\omega(X)$ . Thus

$$X(h) = h\omega(X) \tag{3.8}$$

for any  $X$  tangent to  $M$ . Moreover,  $\sum_{i=1}^{4m-1} g((\nabla_{E_i} A)Y, E_i) = g(A\phi AY, \xi) + \omega(AY) = \omega(AY)$ . Thus from the Codazzi equation

$$\begin{aligned} \omega(AY) &= \sum_{i=1}^{4m-1} g((\nabla_Y A)E_i + \eta(E_i)\phi Y - \eta(Y)\phi E_i - 2g(\phi E_i, Y)\xi \\ &\quad + \sum_{\nu=1}^3 \{\eta_\nu(E_i)\phi_\nu Y - \eta_\nu(Y)\phi_\nu E_i - 2g(\phi_\nu E_i, Y)\xi_\nu\} \\ &\quad + \sum_{\nu=1}^3 \{\eta_\nu(\phi E_i)\phi_\nu \phi Y - \eta_\nu(\phi Y)\phi_\nu \phi E_i\} \\ &\quad + \sum_{\nu=1}^3 \{\eta(E_i)\eta_\nu(\phi Y) - \eta(Y)\eta_\nu(\phi E_i)\}\xi_\nu, E_i \\ &= \sum_{i=1}^{4m-1} g(\nabla_Y \lambda_i E_i, E_i) - \sum_{\nu=1}^3 g(\phi_\nu \xi, \phi_\nu \phi Y) \\ &\quad - \sum_{\nu=1}^3 \eta_\nu(\phi Y) \sum_{i=1}^{4m-1} g(\phi_\nu \phi E_i, E_i) + \eta_1(\phi Y)\eta(\xi_1) \\ &= \sum_{i=1}^{4m-1} Y(\lambda_i) + 2\eta_1(\xi)\eta_1(\phi Y) - \sum_{\nu=1}^3 \eta_\nu(\phi Y)trace(\phi_\nu \phi), \end{aligned} \tag{3.9}$$

for any  $Y$  tangent to  $M$ . As for any  $\nu = 1, 2, 3$ ,  $trace(\phi_\nu \phi) = trace(\phi \phi_\nu) = 2\eta_\nu(\xi)$ , see for example [9], from (3.9) we obtain

$$\omega(AY) = Y(h) = h\omega(Y) \tag{3.10}$$

for any  $Y$  tangent to  $M$ . Taking  $Y = \xi$  in (3.10) we get  $\omega(A\xi) = h\omega(\xi) = \alpha\omega(\xi)$ . Thus  $(h - \alpha)\omega(\xi) = 0$ . As we suppose  $h \neq \alpha$  it follows

$$\omega(\xi) = 0. \tag{3.11}$$

(3.6) and (3.11) yield

$$\text{grad}(\alpha) = 4\eta(\xi_1)\phi_1\xi. \tag{3.12}$$

We know that for any  $X, Y$  tangent to  $M$   $g(\nabla_X \text{grad}(\alpha), Y) = g(\nabla_Y \text{grad}(\alpha), X)$ . This yields

$$\begin{aligned} &(g(\phi AX, \xi_1) + g(\xi, \nabla_X \xi_1))g(\phi_1\xi, Y) + \eta_1(\xi)g(\nabla_X \phi_1\xi, Y) \\ &= (g(\phi AY, \xi_1) + g(\xi, \nabla_Y \xi_1))g(\phi_1\xi, X) + \eta_1(\xi)g(\nabla_Y \phi_1\xi, X). \end{aligned} \tag{3.13}$$

Taking  $Y = \xi$  in (3.13) we obtain  $\eta_1(\xi)g(\nabla_X \phi_1\xi, \xi) = g(\xi, \nabla_X \xi_1)g(\phi_1\xi, X) + \eta_1(\xi)g((\nabla_X \phi_1)\xi, X)$  for any  $X$  tangent to  $M$ . This gives  $-\eta_1(\xi)g(\phi\xi_1, \phi AX) = g(\xi, q_3(\xi)\xi_2 - q_2(\xi)\xi_3 + \phi_1 A\xi)g(\phi_1\xi, X) + \eta_1(\xi)g(-q_2(\xi)\phi_3\xi + q_3(\xi)\phi_2\xi + \eta_1(\xi)A\xi, X)$  for any  $X$  tangent to  $M$ . As we suppose  $\eta_1(\xi) = \eta(\xi_1) \neq 0$  we get  $-g(A\xi_1, X) + \alpha\eta(\xi_1)\eta(X) = -q_2(\xi)g(\phi_3\xi, X) + q_3(\xi)g(\phi_2\xi, X) + \alpha\eta(\xi_1)\eta(X)$  for any  $X$  tangent to  $M$ . Therefore

$$A\xi_1 = q_2(\xi)\phi_3\xi - q_3(\xi)\phi_2\xi. \tag{3.14}$$

The scalar product of (3.14) and  $\xi$  yields  $\alpha\eta(\xi_1) = 0$ . As  $\alpha \neq 0$ ,  $\eta(\xi_1) = 0$  and we arrive at a contradiction.

This finishes the proof of our Proposition. □

Now, if  $\xi \in \mathbb{D}$ , from Theorem 2.2,  $M$  is locally congruent to a real hypersurface of type (B). Therefore consider the case  $\xi \in \mathbb{D}^\perp$ . We can write  $\xi = \xi_1$ .

**Proposition 3.2** *Let  $M$  be a Hopf real hypersurface in  $G_2(\mathbb{C}\mathbb{C}^{m+2})$ ,  $m \geq 3$ , whose shape operator is recurrent with respect to the  $g$ -Tanaka-Webster connection. If  $\xi \in \mathbb{D}^\perp$  and  $\alpha \neq 0$ ,  $M$  is locally congruent to a type (A) real hypersurface.*

**Proof** With our hypothesis and being  $\xi = \xi_1$ , (3.4) becomes

$$\begin{aligned} &\phi Y + \phi_1 Y - \eta_2(Y)\xi_3 + \eta_3(Y)\xi_2 \\ &= -\alpha\phi AY + k\phi AY + A\phi AY - kA\phi Y + \omega(\xi)AY - \alpha\omega(Y)\xi \end{aligned} \tag{3.15}$$

for any  $Y$  tangent to  $M$ .

The scalar product of (3.15) and  $\xi$ , bearing in mind that  $\alpha \neq 0$ , yields  $\omega(Y) = \eta(Y)\omega(\xi)$  for any  $Y$  tangent to  $M$ . From (3.2) we obtain

$$\text{grad}(\alpha) = \alpha\omega(\xi)\xi. \tag{3.16}$$

Therefore for any  $X$  tangent to  $M$   $\nabla_X \text{grad}(\alpha) = X(\alpha\omega(\xi))\xi + \alpha\omega(\xi)\phi AX = \omega(\xi)X(\alpha)\xi + \alpha X(\omega(\xi))\xi + \alpha\omega(\xi)\phi AX$ . If  $Y$  is orthogonal to  $\xi$ ,  $g(\nabla_X \text{grad}(\alpha), Y) = \alpha\omega(\xi)g(\phi AX, Y)$ . Moreover, if  $X$  is also orthogonal to  $\xi$ , as  $g(\nabla_X \text{grad}(\alpha), Y) = g(\nabla_Y \text{grad}(\alpha), X)$ , we get  $\alpha\omega(\xi)g(\phi AX, Y) = \alpha\omega(\xi)g(\phi AY, X)$ . Thus either  $\omega(\xi) = 0$  and  $\omega$  should vanish, which is impossible, or  $g((\phi A + A\phi)X, Y) = 0$  for any  $X, Y$  orthogonal to  $\xi$ . As we also have  $(\phi A + A\phi)\xi = 0$  we obtain

$$\phi A + A\phi = 0. \tag{3.17}$$



From (3.17) it is easy to see that  $A^2\phi = \phi A^2$ . Now from (3.1) we have  $(\nabla_\xi A)\xi_2 = k\phi A\xi_2 - kA\phi\xi_2 + \omega(\xi)A\xi_2 = 2kA\xi_3 + \omega(\xi)A\xi_2$ . If  $X \in \mathbb{D}$  we get  $g((\nabla_\xi A)\xi_2, X) = g(\xi_2, (\nabla_\xi A)X)$  and applying the Codazzi equation this is equal to  $g(\xi_2, \alpha\phi AX + \phi X - A\phi AX + \phi_1 X)$ . This yields

$$g(A^2\xi_3, X) = (2k - \alpha)g(A\xi_3, X) + \omega(\xi)g(A\xi_2, X) \tag{3.18}$$

for any  $X \in \mathbb{D}$ . Developing  $(\nabla_\xi A)\xi_3$  we have

$$-g(A^2\xi_2, X) = (\alpha - 2k)g(A\xi_2, X) + \omega(\xi)g(A\xi_3, X) \tag{3.19}$$

for any  $X \in \mathbb{D}$ . If in (3.18) we take  $\phi X$  instead of  $X$  we get

$$-g(A^2\xi_2, X) = (2k - \alpha)g(A\xi_2, X) - \omega(\xi)g(A\xi_3, X). \tag{3.20}$$

From (3.19) and (3.20)  $g(A^2\xi_2, X) = 0$ . Similarly  $g(A^2\xi_3, X) = 0$  for any  $X \in \mathbb{D}$ . Thus (3.18) and (3.19) become

$$\begin{aligned} \omega(\xi)g(A\xi_2, X) + (2k - \alpha)g(A\xi_3, X) &= 0 \\ -(2k - \alpha)g(A\xi_2, X) + \omega(\xi)g(A\xi_3, X) &= 0 \end{aligned} \tag{3.21}$$

The matrix of coefficients of this homogeneous linear system has as determinant  $(\omega(\xi))^2 + (2k - \alpha)^2$ , and as  $\omega(\xi) \neq 0$ , this determinant does not vanish. This yields  $g(A\xi_2, X) = g(A\xi_3, X) = 0$  for any  $X \in \mathbb{D}$ . Thus  $\mathbb{D}$  is  $A$ -invariant and  $M$  must be locally of type (A).  $\square$

From Propositions 3.1 and 3.2,  $M$  is locally congruent to a real hypersurface either of type (A) or of type (B).

Let  $M$  be a type (A) real hypersurface. Clearly  $(\nabla_\xi A)\xi = 0$ . For (3.1) to be satisfied we should have  $\sqrt{8}\cot(\sqrt{8}r)\omega(\xi)\xi = 0$ . Therefore  $\omega(\xi) = 0$ .

As  $(\nabla_{\xi_i} A)\xi = \sqrt{8}\cot(\sqrt{8}r)\phi A\xi_i - A\phi A\xi_i$ ,  $i = 2, 3$ , for (3.1) to be satisfied this must be equal to  $\sqrt{8}\cot(\sqrt{8}r)\phi A\xi_i - A\phi A\xi_i + \omega(\xi_i)A\xi$ . Thus  $\omega(\xi_i) = 0$ ,  $i = 1, 2$ .

The same occurs if we take any  $X \in T_\lambda$  or  $Y \in T_\mu$ . This means that for a real hypersurface of type (A) to satisfy our condition we should have  $\omega \equiv 0$ , which is impossible.

A similar reasoning applied to a real hypersurface of type (B) shows that these real hypersurfaces do not satisfy our condition and our Theorem is proved.

### References

- [1] Berndt J. Riemannian geometry of complex two-plane Grassmannians. *Rend Sem Mat Univ Politec Torino* 1997; 55: 19–83.
- [2] Berndt J, Suh YJ. Real hypersurfaces in complex two-plane Grassmannians. *Monatsh Math* 1999; 127: 1–14.
- [3] Berndt J, Suh YJ. Isometric flows on real hypersurfaces in complex two-plane Grassmannians. *Monatsh Math* 2002; 137: 87–98.
- [4] Cho JT. CR structures on real hypersurfaces of a complex space form. *Publ Math Debrecen* 1999; 54: 473–487.
- [5] Jeong I, Lee H, Suh YJ. Real hypersurfaces in complex two-plane Grassmannians with generalized Tanaka-Webster parallel shape operator. *Kodai Math J* 2011; 34: 352–366.

- [6] Kim S, Lee H, Yang HY. Real hypersurfaces in complex two-plane Grassmannians with recurrent shape operator. *Bull Malays Math Sci Soc* 2011; 34: 1–14.
- [7] Kobayashi S, Nomizu K. *Foundations of Differential Geometry I*. New York, NY, USA: John Wiley and Sons, Inc., 1963.
- [8] Lee H, Suh YJ. Real hypersurfaces of type B in complex two-plane Grassmannians related to the Reeb vector. *Bull Korean Math Soc* 2010; 47: 551–561.
- [9] Pérez JD, Suh YJ. The Ricci tensor of real hypersurfaces in complex two plane Grassmannians. *J Korean Math Soc* 2007; 44: 211–235.
- [10] Suh YJ. Real hypersurfaces in complex two-plane Grassmannians with parallel shape operator. *Bull Austral Math Soc* 2003; 62: 493–502.
- [11] Suh YJ. Recurrent real hypersurfaces in complex two-plane Grassmannians. *Acta Math Hungar* 2006; 112: 1309–1324.
- [12] Tanaka N. On non-degenerate real hypersurfaces, graded Lie algebras and Cartan connections. *Japan J Math* 1976; 20: 89–102.
- [13] Tanno S. Variational problems on contact Riemannian manifolds. *Trans AMS* 1989; 314: 349–379.
- [14] Webster SM. Pseudo-Hermitian structures on a real hypersurface. *J Diff Geom* 1978; 13: 25–41.