

1-1-2005

Chromosome Counts of Some Veronica L. (Scrophulariaceae) Species from Iran

SHAHRYAR SAEIDI MEHRVARZ

ARDASHIR KHARABIAN

Follow this and additional works at: <https://journals.tubitak.gov.tr/botany>



Part of the [Botany Commons](#)

Recommended Citation

MEHRVARZ, SHAHRYAR SAEIDI and KHARABIAN, ARDASHIR (2005) "Chromosome Counts of Some Veronica L. (Scrophulariaceae) Species from Iran," *Turkish Journal of Botany*. Vol. 29: No. 4, Article 2. Available at: <https://journals.tubitak.gov.tr/botany/vol29/iss4/2>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Botany by an authorized editor of TÜBİTAK Academic Journals. For more information, please contact academic.publications@tubitak.gov.tr.

Chromosome Counts of Some *Veronica* L. (*Scrophulariaceae*) Species from Iran

Shahryar SAEIDI MEHRVARZ

Department of Biology, Faculty of Science, University of Guilan, Rasht – IRAN
E-mail: saeidimz@guilan.ac.ir

Ardeshir KHARABIAN

Department of Agronomy and Plant Breeding, Faculty of Agricultural Science, Islamic Azad University of Rasht, Rasht – IRAN

Received: 02.09.2004

Accepted: 07.03.2005

Abstract: The chromosome numbers of 10 *Veronica* L. species belonging to sect. *Alsinebe* (Griseb.) Lehm. and sect. *Beccabunga* Griseb. from northern Iran are given. The chromosome numbers for 3 of the 10 studied taxa, i.e. *V. siaretensis* E. Lehm., *V. ceratocarpa* C.A.Mey. ($2n = 14$) and *V. hederifolia* L. ($2n = 36$) are presented for the first time. Our results are compared with previous records.

Key Words: *Veronica*, Chromosome number, Iran, mitosis.

Introduction

Sect. *Beccabunga* subsect. *anagalloides* Keller includes *V. anagallis-aquatica* L. and *V. anagalloides* Guss., which presents a taxonomical problem. Sect. *Alsinebe* (Griseb.) Lehm. is also an extremely confusing group and includes 32 species and several subspecies. Hybridisation occurs frequently in this section with 3 ploidy levels, i.e. diploids, tetraploids and hexaploids are known (Öztürk & Fischer, 1982). Some taxa examined have nearly worldwide distribution, while *V. francispetae* M.A.Fischer, *V. siaretensis* E.Lehm. and *V. ceratocarpa* C.A.Mey. are endemic to the Elburz mountain range in Iran (Fischer, 1987).

There are several reports on the cytotaxonomy of various species of *Veronica* (Fischer, 1967, 1973; Podlech & Dieteric, 1969; Ferakova, 1976; Fernandes-Casas, 1977; Aryavand, 1987; Ghaffari 1987). The present study and the previous reports of chromosome numbers in *Veronica* confirm the variation in the basic number.

Materials and Methods

The materials used for this study were collected from wild populations. Flower buds were fixed in Carnoy's

mixture in the field; the ovaries were then excised and stained in 2% aceto-orcein solution for 5 min. They were immersed in 0.075 M KCl at room temperature for 20 min; then the ovaries were treated with enzyme solution containing 5% cellulase and 5% pectinase and adjusted to pH 4.0 for about 20 min at 35 °C in order to dissolve the cell wall (Kurata & Omura, 1978). The ovaries were squashed in 45% acetic acid and then permanently mounted in Entellan to facilitate investigations of somatic mitosis. Permanent preparations were made and studied using a BH-2 Zeiss photomicroscope. Voucher specimens are deposited in the Herbarium of Guilan University.

Results

The chromosome numbers from 13 populations of the 10 species examined are given in Table 1.

V. anagalloides Guss subsp. *heureka* M.A.Fischer (Figure A)

This taxon occupies an area from Central Anatolia to Pakistan. It is distinguishable from *V. anagalloides* subsp. *anagalloides* by its wider and shorter leaves and its suborbicular (not elliptical) and larger capsule. Our counts of $2n = 36$ agree with that given by Podlech & Dieteric

Table 1. Chromosome numbers of 10 *Veronica* species from Iran.

Taxon	Pop. No.	Locality	2n
<i>V. anagalloides</i> Guss subsp. <i>heureka</i> M.A.Fischer	1	Guilan; Asalam, 50 km to Khalkhal, 1700 m, Saeidi 24077	36
<i>V. anagallis-aquatica</i> L. subsp. <i>michauxii</i> (Lam.) A.Jelen.	2	Mazandaran; SW Ramsar, Javaher-Deh, 1800 m, Saeidi 1505	36
<i>V. siaretensis</i> E.Lehm.	3	Gorgan; Ziarat village, 1000-1100 m, Saeidi & Kaviani 1248	14
<i>V. francispetae</i> M.A.Fischer	4	Guilan; near Langroud, Talesh-Mahleh, 30 m, Saeidi 24025	14
<i>V. capillipes</i> Neveski	5	Tehran; from Karaj to Chalus, near Gachsar, 1400 m, Saeidi 1301	28
<i>V. campylopoda</i> Boiss.	6	Gorgan; Bastam, Tash-Olia, Shavar, Saeidi & Kaviani 1311	42
<i>V. polita</i> Fr.	7	Guilan; 7 km from Asalam to Talesh, 500-600 m Saeidi 1315	14
	8	Mazandaran; near Abas-Abad, 1080 m, Saeidi & Asaadi 1452	14
<i>V. persica</i> Poir.	9	Guilan; Daylaman, Shah-Shahidan, 1500-2000 m, Jamzad & Asri 71766	28
	10	Mazandaran; Nowshahr, Kheyroud-Kenar forest, 200 m, Saeidi 1307	28
<i>V. hederifolia</i> L.	11	Tehran; from Karaj to Chalus, near Gachsar, 1400 m, Saeidi 1310	36
<i>V. ceratocarpa</i> C.A.Mey.	12	Mazandaran; Klardasht, 1300-1400 m, Saeidi 24021	14
	13	Guilan; Masuleh, 700-800 m, Saeidi 1197	14

(1969), who studied material from Afghanistan. Öztürk & Fischer (1982) have reported $2n = 18$ for this subspecies based on only one collection of Turkish material of *V. anagallis-aquatica* subsp. *lysimachioides*.

V. anagallis-aquatica L. subsp. *michauxii* (Lam.) A.Jelen. (Figure B)

Some taxa of the *V. anagallis-aquatica* group have an almost worldwide distribution, but its centre of diversity is south-west Asia (Turkey to Pakistan), where there are at least 9 subspecies or species. The tetraploid chromosome number of $2n = 36$ studied here confirms that reported by previous authors (Öztürk & Fischer, 1982) as well as by Aryavand (1987), who examined material from central Iran.

According to Öztürk & Fischer (1982), this species is possibly a hybrid between subsp. *michauxii* and subsp. *anagallis-aquatica*, although capsules with fertile seeds develop.

V. siaretensis E.Lehm. (Figure C)

This species is endemic to Iran, where we found it only in one locality, Gorgan. The chromosome number of $2n = 14$ is the first record for this species.

V. ceratocarpa C.A.Mey. (Figure D)

This taxon is distributed in Turkey, Iran and middle Asia with a wide distribution in the deciduous forests of Elburz mountain in northern Iran. There are no previous counts for this taxon, and so the chromosome number of $2n = 14$ is reported here for the first time.

V. francispetae M.A.Fischer (Figure E)

This species is endemic to northern Iran and grows beneath trees and shrubs. The diploid chromosome number of $2n = 14$ confirms the count reported by Fischer (1981).


V. capillipes Nevski (Figure F)

This species has a disjunct distribution in north-east and central Iran, Pakistan and middle Asia. Meiosis studies of this species by Aryavand (1987) resulted in a count of $n = 14$, which he reported in the abstract,

although in the main text this is incorrectly given as $2n = 14$. Our count of $2n = 28$ for this species agrees with the $n = 14$ reported by Aryavand (1978), who studied plant material from Isfahan province in Iran.

V. campylopoda Boiss. (Figure G)

This species is an Irano-Turanian element with a wide distribution in Iran, Turkey and middle Asia. A meiotic count of $n = 21$ was reported for this species by Aryavand (1987) and Ghaffari (1987). We found $2n = 42$ for this species, which agrees with the previously published counts.



Figures. A-K: Somatic cells of *Veronica* species. A. *V. anagalloides* subsp. *heureka* B. *V. anagallis-aquatica* L. subsp. *michauxii* C. *V. siaretensis* D. *V. ceratocarpa* E. *V. francispetae* F. *V. capillipes* G. *V. campylopoda* H. *V. polita* J. *V. persica* K. *V. hederifolia*. (Scale bar = 10 μ m).

V. polita Fr. (Figure H)

This species is widespread in Asia and Europe and extends into Africa and is also distributed in the USA (Fischer, 1981). The diploid chromosome number of $2n = 14$ coincides with the results of previous authors (Fernandes et al., 1977; Aryavand, 1987; Ghaffari, 1987).

V. persica Poir. (Figure J)

This species is widespread in Iran and throughout the world. Our count of $2n = 28$ agrees with those reported by Ferakova (1976) and Fernandes et al. (1977).

V. hederifolia L. (Figure K)

This species has a relatively wide distribution in Iran, south-west Asia, middle Asia, Turkey and Europe (Fischer, 1981) and can be found at elevations of up to 1000 m. The count of $2n = 36$ is the first record for this species from Iran, although an earlier chromosome count of $2n = 54$ was reported from central Iran by Aryavand (1987), which agrees with previous counts by Fernandes et al. (1977) and Fischer (1981). Based on Danish and Dutch material, Gadella & Kliphuis (1975, 1976) reported $2n = 36$ for this species.

Discussion

The reports so far suggest that each taxon representing a biological entity (subspecies or species) is characterised by one ploidy level, either diploid, tetraploid or hexaploid.

A meiotic count of $n = 14$ was reported for *V. capillipes* by Aryavand (1987). Ghaffari, in unpublished data (personal communication), repeated a mistake in the text of the paper, which gives $2n = 14$, and which was published in Persian by Aryavand (1987). Aryavand (1987) has indicated that the species is diploid, but our studies confirm that *V. capillipes* is a tetraploid species.

Two different cytotypes were known for *V. hederifolia*, with $2n = 54$ the more common. This cytotype is also recorded for *V. hederifolia* in Iran by Aryavand (1987). The other cytotype ($2n = 36$), corresponding to subsp. *lucorum* (Klett & Richter) Hartl. was not previously known in Iran, but according to the present study the cytotype of $2n = 36$ suggests that this subspecies may be present in Iran.

Fischer (1967) concluded that the 2 cytotypes are so different that they can be regarded as separate species, *Veronica sublobata* Fisch. (tetraploid) and *Veronica*

hederifolia L. (hexaploid), rather than as just a subspecies. Neither Nordenstam & Nilsson (1969), nor De Jongh & Kern (1971) accepted this conclusion. They agreed with Fischer that the 2 taxa could be recognised, but they disagreed with him about the species level for these taxa. The *Veronica hederifolia* complex represented by material from the Netherlands by 2 different habitat cytotypes differ morphologically and probably do not exchange genes (Gadella & Kliphuis, 1976). They suggested that the morphological characters can to a certain extent vary, but in all cases tetraploid and hexaploid plants can be identified by means of a combination of at least 3 or 4 such characters. Gadella & Kliphuis (1975, 1976) support Fischer's view that these taxa should be regarded as separate biological species. They agree that the 2 groups are distinguishable based on the colour of the anthers and the corolla, and the form and structure of the seeds. However, we did not find any comparable morphological differences between them in Iran except for the habit of the 2 cytotypes. We found that the tetraploid plants were usually more slender and smaller than the hexaploid plants.

Hybrids between tetraploid and hexaploid plants of *Veronica hederifolia* are unknown and, considering the subspecific rank to which the variants belong, it seems that this subject needs to be studied in more detail before the correct taxa can be assigned. If the view is held that the (apparently rare) intermediate transitional forms are so important that they blur the distinction between the 2 forms, then the tetraploid should be assigned to *Veronica hederifolia* L. subsp. *lucorum* (Klett & Richter) Hartl. and the hexaploid to *Veronica hederifolia* L. subsp. *hederifolia*.

Since these transitional specimens are perfectly fertile and may occur in large proportions in Asian and European populations, this is the main reason why we have decided to assign the subspecific level to these 2 taxa.

These deviating chromosome numbers demonstrate the need for further studies on the *Veronica hederifolia* complex.

Acknowledgements

We are grateful to Dr. Sh. Zarre, Tehran University, for his useful comments. The first author was supported by project number 65 of the University of Guilan Research Council.

References

- Aryavand A (1987). Chromosome numbers in some Iranian *Veronica* L. (*Scrophulariaceae*) species. *Tehran Univ J Sci* 16: 57-64.
- De Jongh SE & Kern JH (1971). De variabiliteit van *Veronica hederifolia* L. in Nederland. *Gorteria* 5 (7-10): 160-165.
- Ferakova V (1976). Index of chromosome numbers of Slovakian flora, part 5. *Acta Fac Rerum Nat Univ Comeninae Bot* 25: 1-18.
- Fernandes-Casas J (1977). In IOPB chromosome number reports LV. *Taxon* 26: 107-109.
- Fernandes A Queiros A & Santos MF (1977). Contribution à la connaissance cytotaxonomique de Spermatophyta de Portugal. *Scrophulariaceae*. *Bot Soc Brot* 51: 37-90.
- Fischer MA (1967). Beiträge zur Cytotaxonomie der *Veronica hederifolia* Gruppe (*Scrophulariaceae*). *Österr Bot Zeitschr* 114: 189-233.
- Fischer MA (1973). Notizen zur Systematik, Chromosomenzahl und Verbreitung einiger *Veronica* Sippen in Kärnten. *Carinthia* II 163/83: 379-388.
- Fischer MA (1981). *Veronica* L., In: Rechinger KH (ed.). *Flora Iranica* 147: 25-165. Graz: Akad. Druck. U. Verlagsanstalt.
- Fischer MA (1987). On the origin of *Veronica persica* (*Scrophulariaceae*). A contribution to the history of a neophytic weed. *Pl Syst Evol* 155: 105-132.
- Gadella TWJ & Kliphuis E (1975). Cytological observations in the *Veronica hederifolia* complex in Denmark. *Bot Tidsskr* 69: 245-247.
- Gadella TWJ & Kliphuis E (1976). Some critical remarks of *Veronica hederifolia* L. in the Netherlands. *Koninklijke Nederlandse Akademie van Wetenschappen (Amsterdam) ser. C* 79: 61-73.
- Ghaffari SM (1987). Chromosome counts of some Angiosperms from Iran II. *Iran J Bot* 3 (2): 183-188.
- Kurata N & Omura T (1978). Karyotype analysis in rice I. A new method for identifying all chromosome pairs. *Jap J Genetics* 53: 251-255.
- Nordenstam B & Nilsson Ö (1969). Taxonomy and distribution of *Veronica hederifolia* s. lat. (*Scrophulariaceae*) in Scandinavia. *Bot Not* 122: 233-247.
- Öztürk A & Fischer MA (1982). Karyosystematics of *Veronica* sect. *Beccabunga* (*Scrophulariaceae*) with special reference to the taxa in Turkey. *Pl Syst Evol* 140: 307-319.
- Podlech D & Dieteric A (1969). Chromosomen studen an Afghanischen Pflanzen. *Candollea* 24 (2): 185-243.