

1-1-2015

Generalized Heineken–Mohamed type groups

OREST ARTEMOVYCH

Follow this and additional works at: <https://journals.tubitak.gov.tr/math>



Part of the [Mathematics Commons](#)

Recommended Citation

ARTEMOVYCH, OREST (2015) "Generalized Heineken–Mohamed type groups," *Turkish Journal of Mathematics*: Vol. 39: No. 2, Article 12. <https://doi.org/10.3906/mat-1408-1>

Available at: <https://journals.tubitak.gov.tr/math/vol39/iss2/12>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Mathematics by an authorized editor of TÜBİTAK Academic Journals. For more information, please contact academic.publications@tubitak.gov.tr.

Generalized Heineken–Mohamed type groups

Orest ARTEMOVYCH*

Institute of Mathematics, Cracow University of Technology, Krakow, Poland

Received: 01.08.2014 • Accepted: 14.01.2015 • Published Online: 23.02.2015 • Printed: 20.03.2015

Abstract: We prove that a torsion group G with all subgroups subnormal is a nilpotent group or $G = N(A_1 \times \cdots \times A_n)$ is a product of a normal nilpotent subgroup N and p_i -subgroups A_i , where $A_i = A_1^{(i)} \cdots A_{m_i}^{(i)} \triangleleft G$, $A_j^{(i)}$ is a Heineken–Mohamed type group, and p_1, \dots, p_n are pairwise distinct primes ($n \geq 1$; $i = 1, \dots, n$; $j = 1, \dots, m_i$ and m_i are positive integers).

Key words: Nilpotent group, indecomposable group, Heineken–Mohamed type group

1. Introduction

The concept of an HM^* -group (i.e. a p -group G with the nilpotent derived subgroup G' and the divisible Černikov quotient group G/G') was introduced by Asar [2] in the class of infinite p -groups. Quasicyclic groups, Heineken–Mohamed type groups (i.e. nonnilpotent groups with nilpotent and subnormal proper subgroups) and Čarin groups (see [5, 10] and [22, Example 1.5.1]) are HM^* -groups. Recall that a group G is called *indecomposable* if any two proper subgroups generate a proper subgroup in G and is called *decomposable* otherwise. Heineken–Mohamed type groups are indecomposable p -groups with the normalizer condition (see examples in [15, 18, 24]). With indecomposable groups are connected barely transitive groups (see [21, Lemma 2.10]).

Casolo [11] proved that a torsion group G with all subgroups subnormal contains a normal nilpotent subgroup N such that G/N is an abelian divisible group of finite rank (see also [12]). We make this result more precise in the following:

Theorem 1.1 *Let G be a torsion group with all subgroups subnormal. Then G is a nilpotent group or $G = NA$ is a product of a normal nilpotent subgroup N and a nonnilpotent subgroup*

$$A = A_1 \times \cdots \times A_n \quad (n \geq 1),$$

where A_i is a p_i -group, $A_i = A_1^{(i)} \cdots A_{m_i}^{(i)} \triangleleft G$, $A_j^{(i)}$ is a Heineken–Mohamed type group, and p_1, \dots, p_n are pairwise distinct primes ($i = 1, \dots, n$; $j = 1, \dots, m_i$ and m_i are positive integers).

Recently Smith [28] established that a group G with all subgroups subnormal such that its torsion part is a π -group, where π is a finite set of primes, is nilpotent-by-(divisible Černikov).

*Correspondence: artemo@usk.pk.edu.pl

2010 AMS Mathematics Subject Classification: 20E22, 20F18, 20J06.

Throughout, p will always denote a prime, C_{p^∞} the quasicyclic p -group, and \mathbb{Z} the ring of integers. For a group G , $G', G'', \dots, G^{(n)}, \dots$ will indicate the terms of derived series of G and G^n the subgroup of G generated by the n th powers of all elements in G .

For notations see [19, 23, 27].

2. Preliminary results

In this part we collect some results concerning groups G with the locally nilpotent derived subgroup G' and the divisible Černikov quotient p -group G/G' .

Lemma 2.1 *Let G be a group with the locally nilpotent derived subgroup G' and the quotient group G/G' be a divisible Černikov p -group. Then the following hold:*

- (i) $G' = [G', G]$;
- (ii) each nontrivial homomorphic image of G has infinite exponent and, in particular, $G = G^m$ for each positive integer m ;
- (iii) G' contains no proper G -invariant subgroup of finite index;
- (iv) if G is a hypercentral group, then it is a divisible Černikov p -group.

Proof For the proof see, e.g., [2, 3]. □

Lemma 2.2 [20, Chapter XV, § 63] *Let G be a group with the normalizer condition. If H is a normal hypercentral subgroup of G and the quotient group G/H is cyclic, then G is a hypercentral group.*

Lemma 2.3 [13, Lemma 1.18] *If a hypercentral group $G = AB$ is a product of an abelian normal subgroup A and a divisible abelian subgroup B , then G is abelian.*

Lemma 2.4 [13, Proposition 1.7] *If a p -group G contains an abelian subgroup of finite index, then it is hypercentral.*

Lemma 2.5 *Let G be a nonhypercentral group with the normalizer condition. If G contains a normal hypercentral soluble subgroup A with the quasicyclic quotient group G/A , then every nonhypercentral subgroup is subnormal in G .*

Proof If H is a proper nonhypercentral subgroup of G , then, in view of Lemma 2.2, $G = AH$ and, by [17, Lemma],

$$G/A'(A \cap H)$$

is a hypercentral group. Therefore, it is abelian by Lemma 2.3 and $A'H \triangleleft G$. Since $A'H/A''(A' \cap H)$ is hypercentral, by the same argument we obtain that $A''H \triangleleft A'H$. By induction, H is subnormal in G . □

Lemma 2.6 *The following properties hold:*

- (i) Any indecomposable soluble torsion group G is a locally finite p -group.
- (ii) Let G be a group with the hypercentral derived subgroup G' and the quotient group G/G' be a divisible Černikov p -group. If G is indecomposable, then it is a locally finite p -group.
- (iii) Any indecomposable locally nilpotent group is a p -group.

Proof (i) If G is abelian and indecomposable, then it is a quasicyclic or cyclic p -group. Therefore, we suppose that G is nonabelian. Then G/G' is a divisible Černikov p -group. Assume that there is an integer k such that $G/G^{(k)}$ is a p -group and $G^{(k)}$ is not a p -group, where $1 \leq k \leq n-1$ and n is the derived length of G .

The group G is a soluble torsion group and so it is a locally finite group. Since G is soluble and indecomposable, it is countable. Then G has a local system consisting of its finite subgroups. Since the group G contains an element of order p and, by Hall's Theorem, it has a Hall p -subgroup and a Hall p' -subgroup, G is generated by them. Then G being indecomposable implies that G is a p -group.

(ii) We denote the derived subgroup G' by H . If H is not torsion, then the torsion part τH is normal in H and the quotient group $H_1 = H/\tau H$ is torsion-free hypercentral. Then H_1/H'_1 is not torsion (see, e.g., [27, Exercises 12.2.6]) and therefore without loss of generality we can assume that H is a torsion-free abelian group. Let q be a prime different from p . By [9, Lemma 2.3], there exists a proper submodule N of a $\mathbb{Z}[G/H]$ -module H such that H/N is torsion as a group and contains an element of order q . Then G/N is a decomposable group, a contradiction. Hence, G is a torsion group and this yields that G is a locally finite p -group.

(iii) It follows by the same argument as in (ii). □

If G is a barely transitive group with the locally nilpotent derived subgroup G' and the quotient group G/G' is a divisible Černikov p -group, then G is an indecomposable locally finite p -group by Lemma 2.3 and [7, Proposition 2.4]. In view of Lemma 2.6, it is a very interesting open question whether nontorsion (and therefore non-“barely transitive”) indecomposable soluble groups exist.

Lemma 2.7 *Let G be a group with the hypercentral derived subgroup G' and the divisible Černikov quotient p -group G/G' . Then the following hold:*

- (1) if the derived subgroup G' has no proper supplements in G , then G is a p -group;
- (2) if G is an indecomposable group, then every proper subgroup of it is hypercentral;
- (3) if G is a soluble group with the derived subgroup G' of finite exponent, then every finite subgroup is subnormal in G ;
- (4) if G is a soluble group with the normalizer condition, then
 - (i) G' is not supplemented nontrivially in G ;
 - (ii) $G/G' \cong \mathbb{C}_{p^\infty}$ if and only if G is an indecomposable group;
 - (iii) if G' is nilpotent of finite exponent, then all subgroups are subnormal in G .

Proof (1) We can prove by the same argument as in the proof of Lemma 2.6(ii).

(2) Since G/G' is a quasicyclic group and $G'H$ is proper in G for any proper subgroup of G , we conclude that $G'H = G'\langle a \rangle$ for some element $a \in G$. Hence, $G'H$ is a hypercentral group.

(3) If F is a finite subgroup of G , then $G'F$ is a proper subgroup of G and we deduce that $\overline{G'F}$ is a nilpotent subgroup of $\overline{G} = G/G''$ by [4, Lemma 3.8]. As a consequence, $G''F$ is subnormal in G . Repeating this process, we obtain that $G^{(n)}F$ is subnormal in G for any positive integer n .

(4) Let G be a soluble group with the normalizer condition. Since G' is hypercentral and G/G' is a p -group, G is a p -group. By [17, Lemma], $G = G'S$ for some proper subgroup S and it follows that the quotient group

$$\overline{G} = G/G''(G' \cap S)$$

is hypercentral, which leads to a contradiction. If, moreover, G' is of finite exponent, then every subgroup is subnormal in G by condition (3) and [2, Lemma 3.6]. □

Lemma 2.8 *Let G be a nonhypercentral group with the hypercentral derived subgroup G' and the divisible Černikov quotient p -group G/G' . Then the following hold:*

(1) *if all subgroups of G are subnormal, then*

$$G = G_1 \cdots G_n \cdot D \quad (n \geq 1),$$

where D is a nilpotent group, G_i is a Heineken–Mohamed type group, $D' \leq G' = G'_1 \cdots G'_n \leq D$ and $G'_i = [G'_i, G_i] = [G', G_i] = [G'_i, G]$;

(2) *if G/G' is a quasicyclic p -group, then*

- (a) *G satisfies the normalizer condition if and only if it is a minimal nonhypercentral group;*
- (b) *every proper subgroup is subnormal in G if and only if G is a Heineken–Mohamed type group.*

Proof (1) By definition,

$$\overline{G} = G/G' = \overline{T}_1 \times \cdots \times \overline{T}_s$$

is a direct product of quasicyclic p -subgroups \overline{T}_i ($i = 1, \dots, s$). If T_i is an inverse image of \overline{T}_i in G , then

$$G' \leq T_i \triangleleft G.$$

Since G is nonhypercentral, there exists an integer l ($1 \leq l \leq s$) such that T_1, \dots, T_l are nonhypercentral, while T_{l+1}, \dots, T_s are hypercentral groups. We denote the product $T_{l+1} \cdots T_s$ by D . Then D is hypercentral and, by [26, Theorem 2.7], it is a nilpotent group.

Let $T = T_i$ ($i = 1, \dots, l$). Obviously,

$$[G', T] \leq T' \leq G'.$$

Moreover, the quotient group $T/[G', T]$ (and respectively $T/[T', T]$) is nilpotent (and consequently abelian), and therefore we obtain that

$$T' = [T', T] = [T', G] = [G', T].$$

Since $G/(T'_1 \cdots T'_l)$ is a hypercentral group, we conclude that the derived subgroup $G' = T'_1 \cdots T'_l$.

(2) Let G/G' be a quasicyclic p -group.

(a) Suppose that G satisfies the normalizer condition. If it is decomposable, then $G = G'S$ for some $S < G$, and so the quotient group

$$G/G''(G' \cap S)$$

is abelian by [17, Lemma], a contradiction. Hence, G is an indecomposable group. By Lemma 2.7(2), G is a minimal nonhypercentral group. The converse is an immediate consequence of the fact that $G'K$ is proper in G for every proper subgroup K of G and therefore K is not equal to its normalizer $N_G(K)$.

(b) Assume that every proper subgroup is subnormal in G . Then G satisfies the normalizer condition. By [11, Theorem 1], the derived subgroup G' is nilpotent. If G is a hypercentral group, then, by [26, Theorem 2.7], it is nilpotent and, consequently, $G \cong \mathbb{C}_{p^\infty}$. Therefore, we assume that G is nonhypercentral and, by (a), it is a minimal nonhypercentral group. If we assume that G is decomposable (and so $G = G'S$ for some $S < G$), then, by [17, Lemma],

$$\bar{G} = G/(G' \cap S) = \bar{G}' \times \bar{S}$$

is a hypercentral group. However, then it is abelian, a contradiction. Hence, G is an indecomposable group and for any $K < G$ there exists $a \in G$ such that $G'K = G'\langle a \rangle < G$ and, by [1, Lemma 2.4], $G'K$ is a nilpotent group. Thus, G is a Heineken–Mohamed type group.

The converse is clear. □

Corollary 2.9 *Let G be a nonabelian indecomposable soluble group. If G' is a nilpotent group of finite (respectively infinite) exponent, then G is a Heineken–Mohamed type group (respectively a minimal nonhypercentral group).*

Proof By Lemma 2.6, G is a p -group for some prime p . Let B be a proper subgroup of G/G' and $E(G', B)$ be a subgroup of G , which is an extension of G' by B . If G' is a nilpotent group of finite (respectively infinite) exponent, then [4, Lemma 3.8] (respectively Lemma 2.4) yields that the quotient group $E(G', B)/G''$ is nilpotent (respectively hypercentral). By the Hall Theorem [14, Theorem 7] (respectively the Betten Theorem [8]), $E(G', B)$ is a nilpotent (respectively hypercentral) group. Hence, G is a minimal nonnilpotent (respectively nonhypercentral) group. Moreover, if the derived subgroup G' is of finite exponent, then, by Lemma 2.7, G is a Heineken–Mohamed type group. □

Belyaev and Kuzucuoğlu [6, Corollary 1] proved that a Heineken–Mohamed type group has a faithful barely transitive representation. Then we prove the following:

Corollary 2.10 *A locally finite barely transitive group G with almost locally p -soluble point stabilizer is a Heineken–Mohamed type group or $G \cong \mathbb{C}_{p^\infty}$.*

Proof By [21, Lemma 2.10], G is indecomposable and, by Proposition 3 of [16], the derived subgroup G' is nilpotent of finite exponent. If G is nonabelian, then it is a Heineken–Mohamed type group by Corollary 2.9. □

3. Proof of Theorem 1.1

Suppose that G is a nonnilpotent p -group. As it was proved in [11], a torsion group G with subnormal proper subgroups contains a nilpotent normal subgroup N such that G/N is a divisible abelian Černikov p -group. If

$N^p = N$, then, by [25, Lemma 1], a subgroup $N\langle a \rangle$ is nilpotent (and so abelian) for any $a \in G$, a contradiction. Hence, $N^p \neq N$.

Let k, l be nonnegative integers. By [2, Lemma 3.3],

$$\overline{G} = G/N^{p^k} = \overline{T} \overline{N}$$

is a product of a maximal normal HM^* -subgroup \overline{T} and a normal nilpotent subgroup \overline{N} . If we denote the inverse image of \overline{T} by T and $M = T \cap N$, then, by [2, Lemma 3.3],

$$T/M^{p^k} = \overline{T}_0 \overline{M},$$

where \overline{T}_0 is a maximal normal HM^* -subgroup of T/M^{p^k} . Let T_0 be an inverse image of \overline{T}_0 in a group T . If $l < k$, then

$$T_0/M^{p^k}T'_0 \text{ and } T_0/M^{p^l}T'_0$$

are divisible Černikov groups, $M^{p^k}T'_0 \leq M^{p^l}T'_0$, and so the section

$$M^{p^l}T'_0/M^{p^k}T'_0$$

is a finite group. There exists a subgroup D of T_0 such that D/T'_0 is the divisible part of T_0/T'_0 . Then $T = T_0M = DM$ and $D' = T'_0$. This yields that D is an HM^* -group. The conclusion follows from Lemma 2.8.

Acknowledgment

The author would like to thank the referee for helpful suggestions.

References

- [1] Arikan A, Asar AO. On periodic groups in which every proper subgroup is an NC -group. Turk J Math 1994; 18: 255–262.
- [2] Asar AO. On nonnilpotent p -groups and the normalizer condition. Turk J Math 1994; 18: 114–129.
- [3] Asar AO. A characterizations of \overline{NC} - p -groups. Turk J Math 1997; 21: 475–483.
- [4] Baumslag G. Wreath products and p -groups. P Cambridge Phil Soc 1959; 55: 224–231.
- [5] Belyaev VV. Locally finite groups with almost abelian proper subgroups. Siberian Math J 1983; 24: 11–17.
- [6] Belyaev VV, Kuzucuoğlu M. Barely transitive and Heineken Mohamed groups. J London Math Soc 1997; 55: 261–263.
- [7] Betin C, Kuzucuoğlu M. Description of barely transitive groups with soluble point stabilizer. Comm Algebra 2009; 37: 1901–1907.
- [8] Betten A. Hinreichende Kriterien für die Hyperzentralität einer Gruppen. Arch Math (Basel) 1960; 30: 471–480 (in German).
- [9] Bruno B, Phillips RE. A note on groups with nilpotent-by-finite proper subgroups. Arch Math (Basel) 1995; 65: 369–374.
- [10] Čarin VS. A remark on the minimal condition for subgroups. Dokl Akad Nauk SSSR (N.S.) 1949; 66: 575–576 (in Russian).

- [11] Casolo C. On the structure of groups with all subgroups subnormal. *J Group Theory* 2002; 5: 293–300.
- [12] Casolo C. Groups with many subnormal subgroups. *Note Mat* 2008; 28 (Suppl. 2): 1–149.
- [13] Černikov SN. Groups with Given Properties of a System of Subgroups. Moscow, USSR: Nauka, 1980 (in Russian).
- [14] Hall P. Some sufficient condition for a group to be nilpotent. *Illinois J Math* 1958; 2: 787–801.
- [15] Hartley B. A note on the normalizer condition. *P Cambridge Phil Soc* 1973; 74: 11–15.
- [16] Hartley B, Kuzucuoğlu M. Non-simplicity of locally finite barely transitive groups. *P Edinburgh Math Soc* 1997; 40: 489–490.
- [17] Heineken H. Normalizer condition and nilpotent normal subgroups. *J London Math Soc* 1971/72; 4: 458–460.
- [18] Heineken H, Mohamed IJ. A group with trivial centre satisfying the normalizer condition. *J Algebra* 1968; 10: 368–376.
- [19] Kegel OH, Wehrfritz BAF. Locally Finite Groups. Amsterdam, the Netherlands: North-Holland Publishing Co., 1973.
- [20] Kuroš AG. The Theory of Groups. Moscow, USSR: Nauka, 1967 (in Russian).
- [21] Kuzucuoğlu M. Barely transitive permutation groups, *Arch Math (Basel)* 1990; 55: 521–532.
- [22] Lennox JC, Robinson DJS. The Theory of Infinite Soluble Groups. Oxford, UK: Clarendon Press, 2004.
- [23] Lennox JC, Stonehewer SE. Subnormal Subgroups of Groups. Oxford, UK: Clarendon Press, 1987.
- [24] Menegazzo F. Groups of Heineken-Mohamed. *J Algebra* 1995; 171: 807–825.
- [25] Möhres W. Torsionsgruppen deren Untergruppen alle subnormal sind. *Geom Dedicata* 1989; 31: 237–244 (in German).
- [26] Möhres W. Hyperzentrale Torsionsgruppen deren Untergruppen alle subnormal sind. *Illinois J Math* 1991; 35: 147–157 (in German).
- [27] Robinson DJS. A Course in the Theory of Groups. New York, NY, USA: Springer-Verlag, 1980.
- [28] Smith H. Groups that involve finitely many primes and have all subgroups subnormal. *J Algebra* 2011; 347: 133–142.