

1-1-2014

## Gradient estimates for the porous medium type equation on smooth metric measure space

DENG YIHUA

Follow this and additional works at: <https://journals.tubitak.gov.tr/math>



Part of the [Mathematics Commons](#)

---

### Recommended Citation

YIHUA, DENG (2014) "Gradient estimates for the porous medium type equation on smooth metric measure space," *Turkish Journal of Mathematics*: Vol. 38: No. 6, Article 4. <https://doi.org/10.3906/mat-1310-24>

Available at: <https://journals.tubitak.gov.tr/math/vol38/iss6/4>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Mathematics by an authorized editor of TÜBİTAK Academic Journals. For more information, please contact [academic.publications@tubitak.gov.tr](mailto:academic.publications@tubitak.gov.tr).

## Gradient estimates for the porous medium type equation on smooth metric measure space

Deng YIHUA\*

Department of Mathematics and Computing Science Hengyang Normal University Hengyang, Hunan, P.R. China

Received: 13.10.2013 • Accepted: 19.03.2014 • Published Online: 24.10.2014 • Printed: 21.11.2014

**Abstract:** The porous medium equation arises in different applications to model diffusive phenomena. In this paper, we obtain several gradient estimates for some porous medium type equations on smooth metric measure space with N-Bakry-Emery Ricci tensor bounded from below. In particular, we improve and generalize some current gradient estimates for the porous medium equations.

**Key words:** Gradient estimates, porous medium type equation, smooth metric measure space, positive solution

### 1. Introduction

A smooth metric measure space is a triple,  $(M^n, g, e^{-f} dv)$ , where  $M^n$  is a complete  $n$ -dimensional Riemannian manifold with metric  $g$ ,  $f$  is a smooth real-valued function on  $M^n$ , and  $dv$  is the Riemannian volume density. Smooth metric measure spaces carry a natural analog of the Laplace-Beltrami operator  $\Delta$ , the  $f$ -Laplacian, which is also called drifting Laplacian or Witten-Laplacian, defined for a function  $u$  by  $\Delta_f u = \Delta u - g(\nabla f, \nabla u) = \Delta u - \langle \nabla f, \nabla u \rangle$ . The N-Bakry-Emery Ricci tensor is defined by  $Ric_f^N = Ric + Hess f - \frac{1}{N} df \otimes df$ . A natural question about smooth metric measure space is which of the results about the Ricci tensor and the Laplace-Beltrami operator can be extended to the N-Bakry-Emery Ricci tensor and the  $f$ -Laplacian. For example, in [15], Yang discussed the gradient estimates for the following parabolic equation,

$$\frac{\partial}{\partial t} u = \Delta u + au \log u + bu,$$

on Riemannian manifolds. In [8], Huang and Ma considered the gradient estimates for the following parabolic equation,

$$\frac{\partial}{\partial t} u = \Delta_f u + au \log u + bu,$$

on smooth metric measure spaces. Inspired by the discussions of the gradient estimate of the harmonic function and positive solutions to linear heat flow on Riemannian manifolds, the authors in [1–3] discussed the gradient estimates of the  $f$ -harmonic function and positive solutions to linear heat flow on smooth metric measure spaces.

\*Correspondence: dengchen4032@126.com

This research was supported by the Natural Science Foundation of Hunan province (No.14JJ2120), Scientific Research Fund of Hunan Provincial Education Department (No.14A020), and was partly supported by the Key Constructional Discipline of Hunan Province (No.[2011]76).

2010 *AMS Mathematics Subject Classification*: Primary: 35B45; Secondary: 58J05.

Motivated by the study of eigenvalue estimates for the Laplace-Beltrami operator on Riemannian manifolds, the authors in [4,5,11,12,14] studied the eigenvalue estimates for drifting Laplacians on smooth metric measure spaces.

In [10], Lu et al. obtained some gradient estimates for the following porous medium equation on Riemannian manifolds with Ricci curvature bounded from below:

$$u_t = \Delta u^m, \tag{1.1}$$

where  $m > 1$ . In [6], Huang and Li improved the results in [10]. In [7], Huang and Li studied the following porous medium type equation,

$$u_t = \Delta_f u^m, \tag{1.2}$$

on smooth metric measure space. Under the assumption that the  $N$ -dimensional Bakry-Emery Ricci curvature is bounded from below, Huang and Li obtained some gradient estimates that generalized the results in [6] and [10].

Inspired by [6] and [7], we make further discussions of gradient estimates for positive solutions of equation (1.2). For this purpose, we let

$$v = \frac{m}{m-1}u^{m-1}, \quad a = \frac{(n+N)(m-1)}{(n+N)(m-1)+2}, \quad b = \frac{n+N}{(n+N)(m-1)+2}. \tag{1.3}$$

**Theorem 1.1** *Let  $(M^n, g, e^{-f} dv)$  be a smooth metric measure space. Suppose that  $u$  is a positive solution to (1.2). If  $Ric_f^N(B_p(2R)) \geq -K$  and  $K \geq 0$ , then on the ball  $B_p(R)$  we have*

$$\frac{|\nabla v|^2}{v} - \alpha(t) \frac{v_t}{v} \leq \frac{1}{1+a(\alpha-1)} \left\{ \frac{CHa\alpha^2}{R^2} \left( \frac{bm^2\alpha^2}{2(\alpha-1)(1-a)} + 3 + \sqrt{KR} \right) + \frac{a\alpha^2}{t} \right\}, \tag{1.4}$$

where  $C$  is a constant depending only on  $n$  and

$$H = \sup_{B_p(2R) \times [0, T]} (m-1)v, \quad \alpha(t) = e^{2HKt}.$$

**Corollary 1.2** *Let  $(M^n, g, e^{-f} dv)$  be a smooth metric measure space with  $Ric_f^N \geq -K$  and  $K \geq 0$ . Suppose that  $(M^n, g)$  is a complete noncompact Riemannian manifold and  $u$  is a positive solution to (1.2), then*

$$\frac{|\nabla v|^2}{v} - \alpha(t) \frac{v_t}{v} \leq \frac{1}{1+a(\alpha-1)} \frac{a\alpha^2(t)}{t}, \tag{1.5}$$

where  $\alpha(t) = e^{2SKt}$ ,  $S = \sup_{M^n \times [0, T]} (m-1)v$ .

**Theorem 1.3** *Let  $(M^n, g, e^{-f} dv)$  be a smooth metric measure space. Suppose that  $u$  is a positive solution to (1.2). If  $Ric_f^N(B_p(2R)) \geq -K$  and  $K \geq 0$ , then on the ball  $B_p(R)$  we have*

$$\frac{|\nabla v|^2}{v} - \alpha(t) \frac{v_t}{v} - \varphi(t) \leq \frac{CHa\alpha^2(t)}{R^2} \left[ (3 + \sqrt{KR}) + \frac{bm^2\alpha^2(t)}{\tanh(HKt)} \right], \tag{1.6}$$

where  $C$  is a constant depending only on  $n$ ,  $\alpha(t) = 1 + \frac{2}{3}SKt$ ,  $S = \sup_{M^n \times [0, T]} (m-1)v$ , and

$$H = \sup_{B_p(2R) \times [0, T]} (m-1)v, \quad \varphi(t) = \frac{a}{t} + aSK + \frac{a}{3}(SK)^2t + \frac{\lambda}{t^2}, \quad \lambda \geq 0.$$

**Corollary 1.4** *Let  $(M^n, g, e^{-f} dv)$  be a smooth metric measure space with  $Ric_f^N \geq -K$  and  $K \geq 0$ . Suppose that  $(M^n, g)$  is a complete noncompact Riemannian manifold and  $u$  is a positive solution to (1.2), then*

$$\frac{|\nabla v|^2}{v} - \alpha(t) \frac{v_t}{v} - \varphi(t) \leq 0, \tag{1.7}$$

where  $\varphi(t)$  and  $\alpha(t)$  are defined in Theorem 1.3.

**Remark 1** *Obviously, Theorem 1.1 and Corollary 1.2 in this paper are better than Theorem 1.3 and Corollary 1.4 in [6], respectively.*

**Remark 2** *Theorem 1.3 and Corollary 1.4 in this paper are more general than Theorem 1.4 and Corollary 1.7 in [6], respectively.*

**2. Some lemmas**

To prove Theorem 1.1 and Theorem 1.3, we need some lemmas. Suppose that  $u$  is a positive solution to (1.2).

Let  $v = \frac{m}{m-1}u^{m-1}$ . Direct calculation shows that

$$v_t = mu^{m-2} \Delta_f u^m = (m-1)v \Delta_f v + |\nabla v|^2. \tag{2.1}$$

Since  $v \neq 0$ , then (2.1) is equivalent to

$$\frac{v_t}{v} = (m-1) \Delta_f v + \frac{|\nabla v|^2}{v}. \tag{2.2}$$

Let  $L = \partial_t - (m-1)v \Delta_f$  and  $F = \frac{|\nabla v|^2}{v} - \alpha \frac{v_t}{v} - \varphi$ . We have the following.

**Lemma 2.1** *Suppose that  $u$  is a positive solution to (1.2). Then*

$$L\left(\frac{v_t}{v}\right) = (m-1) \frac{v_t}{v} \Delta_f v + \frac{v_t}{v} \frac{|\nabla v|^2}{v} + 2m \nabla v \nabla \left(\frac{v_t}{v}\right), \tag{2.3}$$

$$\begin{aligned} L\left(\frac{|\nabla v|^2}{v}\right) &\leq 2(m-1) \frac{|\nabla v|^2}{v} \Delta_f v - \frac{2(m-1)}{N+n} |\Delta_f v|^2 \\ &\quad - 2(m-1) Ric_f^N(\nabla v, \nabla v) + 2m \nabla v \nabla \left(\frac{|\nabla v|^2}{v}\right) + \frac{|\nabla v|^4}{v^2}. \end{aligned} \tag{2.4}$$

**Proof** Direct calculation shows that

$$\nabla\left(\frac{v_t}{v}\right) = \frac{\nabla v_t}{v} - \frac{v_t \nabla v}{v^2}. \tag{2.5}$$

Therefore, we get

$$\Delta\left(\frac{v_t}{v}\right) = \frac{1}{v} \Delta v_t - \frac{v_t}{v^2} \Delta v - \frac{2}{v^2} \nabla v \nabla v_t + \frac{2v_t}{v^3} |\nabla v|^2, \tag{2.6}$$

and

$$(m-1)v\langle \nabla f, \nabla(\frac{v_t}{v}) \rangle = (m-1)\langle \nabla f, \nabla v_t \rangle - (m-1)\frac{v_t}{v}\langle \nabla f, \nabla v \rangle. \quad (2.7)$$

By (2.2), we have

$$\partial_t(\frac{v_t}{v}) = (m-1)\partial_t(\Delta_f v) + \partial_t(\frac{|\nabla v|^2}{v}) = (m-1)\Delta_f v_t + \frac{2}{v}\nabla v \nabla v_t - \frac{v_t}{v}\frac{|\nabla v|^2}{v}. \quad (2.8)$$

According to (2.4), (2.5), (2.6), (2.7), and (2.8), we conclude that

$$L(\frac{v_t}{v}) = \partial_t(\frac{v_t}{v}) - (m-1)v\Delta_f(\frac{v_t}{v}) = (m-1)\frac{v_t}{v}\Delta_f v + \frac{v_t}{v}\frac{|\nabla v|^2}{v} + 2m\nabla v \nabla(\frac{v_t}{v}).$$

On the other hand, by (2.1) we get

$$\begin{aligned} \partial_t(\frac{|\nabla v|^2}{v}) &= \frac{2v\nabla v \nabla v_t - |\nabla v|^2 v_t}{v^2} \\ &= \frac{2\nabla v}{v}\nabla((m-1)v\Delta_f v + |\nabla v|^2) - \frac{|\nabla v|^2}{v^2}((m-1)v\Delta_f v + |\nabla v|^2) \\ &= (m-1)\frac{|\nabla v|^2}{v}\Delta_f v + 2(m-1)\nabla v \nabla \Delta_f v + \frac{2\nabla v}{v}\nabla|\nabla v|^2 - \frac{|\nabla v|^4}{v^2}. \end{aligned} \quad (2.9)$$

Direct calculation shows that

$$\begin{aligned} \Delta_f \frac{|\nabla v|^2}{v} &= \Delta \frac{|\nabla v|^2}{v} - \langle \nabla f, \nabla \frac{|\nabla v|^2}{v} \rangle \\ &= \frac{1}{v}\Delta|\nabla v|^2 - \frac{|\nabla v|^2}{v^2}\Delta v - \frac{2}{v^2}\nabla v \nabla|\nabla v|^2 + \frac{2}{v^3}|\nabla v|^4 - \frac{1}{v}\langle \nabla f, \nabla|\nabla v|^2 \rangle + \frac{|\nabla v|^2}{v^2}\langle \nabla f, \nabla v \rangle \\ &= \frac{1}{v}\Delta_f|\nabla v|^2 - \frac{|\nabla v|^2}{v^2}\Delta_f v - \frac{2}{v^2}\nabla v \nabla|\nabla v|^2 + \frac{2}{v^3}|\nabla v|^4. \\ &= \frac{1}{v}\Delta_f|\nabla v|^2 - \frac{|\nabla v|^2}{v^2}\Delta_f v - \frac{2\nabla v}{v}\nabla(\frac{|\nabla v|^2}{v}). \end{aligned} \quad (2.10)$$

According to (2.9) and (2.10), we obtain

$$\begin{aligned} L(\frac{|\nabla v|^2}{v}) &= \partial_t(\frac{|\nabla v|^2}{v}) - (m-1)v\Delta_f \frac{|\nabla v|^2}{v} \\ &= 2(m-1)(\frac{|\nabla v|^2}{v}\Delta_f v + \nabla v \nabla \Delta_f v - \frac{1}{2}\Delta_f|\nabla v|^2) + 2m\nabla v \nabla(\frac{|\nabla v|^2}{v}) + \frac{|\nabla v|^4}{v^2}. \end{aligned} \quad (2.11)$$

According to [9,13], we have

$$\frac{1}{2}\Delta_f|\nabla v|^2 \geq \frac{1}{N+n}|\Delta_f v|^2 + \nabla v \nabla \Delta_f v + Ric_f^N(\nabla v, \nabla v). \quad (2.12)$$

By (2.11) and (2.12), we conclude that (2.4) is true.  $\square$

**Lemma 2.2** ([7]) *The function  $F$  satisfies the following inequality:*

$$L(F) \leq -\frac{2(m-1)}{N+n}|\Delta_f v|^2 - 2(m-1)Ric_f^N(\nabla v, \nabla v) + 2m\nabla v \nabla F - ((m-1)\Delta_f v)^2 + (1-\alpha)\left(\frac{v_t}{v}\right)^2 - \alpha' \frac{v_t}{v} - \varphi'. \tag{2.13}$$

**Proof** For the reader’s convenience, we give the details of the proof of Lemma 2.2. By (2.3) and (2.4), we have

$$\begin{aligned} L(F) &= L\left(\frac{|\nabla v|^2}{v}\right) - \alpha L\left(\frac{v_t}{v}\right) - \alpha' \frac{v_t}{v} - \varphi' \\ &\leq 2(m-1)\frac{|\nabla v|^2}{v}\Delta_f v - \frac{2(m-1)}{N+n}|\Delta_f v|^2 - 2(m-1)Ric_f^N(\nabla v, \nabla v) \\ &\quad + 2m\nabla v \nabla\left(\frac{|\nabla v|^2}{v}\right) + \frac{|\nabla v|^4}{v^2} - \alpha(m-1)\frac{v_t}{v}\Delta_f v - \alpha\frac{v_t}{v}\frac{|\nabla v|^2}{v} \\ &\quad - 2\alpha m \nabla v \nabla\left(\frac{v_t}{v}\right) - \alpha' \frac{v_t}{v} - \varphi'. \end{aligned} \tag{2.14}$$

By the definition of  $F$ , we have

$$2m\nabla v \nabla\left(\frac{|\nabla v|^2}{v}\right) - 2\alpha m \nabla v \nabla\left(\frac{v_t}{v}\right) = 2m\nabla v \nabla\left(\frac{|\nabla v|^2}{v} - \alpha\frac{v_t}{v}\right) = 2m\nabla v \nabla F. \tag{2.15}$$

According to (2.2), we get

$$\frac{|\nabla v|^4}{v^2} - \alpha\frac{v_t}{v}\frac{|\nabla v|^2}{v} = (1-\alpha)\frac{|\nabla v|^2}{v}\frac{v_t}{v} - (m-1)\frac{|\nabla v|^2}{v}\Delta_f v. \tag{2.16}$$

Using (2.2) again, we arrive at

$$\begin{aligned} &(m-1)\frac{|\nabla v|^2}{v}\Delta_f v - \alpha(m-1)\frac{v_t}{v}\Delta_f v + (1-\alpha)\frac{|\nabla v|^2}{v}\frac{v_t}{v} \\ &= \frac{|\nabla v|^2}{v}\left(\frac{v_t}{v} - \frac{|\nabla v|^2}{v}\right) - \alpha\frac{v_t}{v}\left(\frac{v_t}{v} - \frac{|\nabla v|^2}{v}\right) + (1-\alpha)\frac{|\nabla v|^2}{v}\frac{v_t}{v} \\ &= 2\frac{|\nabla v|^2}{v}\frac{v_t}{v} - \frac{|\nabla v|^4}{v^2} - \alpha\left(\frac{v_t}{v}\right)^2 \\ &= -\left(\alpha\frac{v_t}{v} - \frac{|\nabla v|^2}{v}\right)^2 + (1-\alpha)\left(\frac{v_t}{v}\right)^2 \\ &= -((m-1)\Delta_f v)^2 + (1-\alpha)\left(\frac{v_t}{v}\right)^2. \end{aligned} \tag{2.17}$$

Putting (2.15), (2.16), and (2.17) into (2.14), we conclude that (2.13) is true.

**3. The proof of Theorem 1.1**

Let  $\bar{F} = \frac{|\nabla v|^2}{v} - \alpha \frac{v_t}{v}$ , where  $\alpha = e^{2HKt}$ . We consider  $\bar{F}$  in the geodesic ball  $B_p(2R)$ , which is centered at  $p$  with radius  $2R$ . Since  $Ric_f^N(B_p(2R)) \geq -K$ , by (2.13) and the definition of  $H$  and  $a$ , we have

$$\begin{aligned} L(\bar{F}) &\leq \frac{-2}{(m-1)(N+n)}((m-1)\Delta_f v)^2 + 2vK \frac{|\nabla v|^2}{v} + 2m\nabla v \nabla \bar{F} \\ &\quad - ((m-1)\Delta_f v)^2 + (1-\alpha)\left(\frac{v_t}{v}\right)^2 - \alpha' \frac{v_t}{v} \\ &\leq -\frac{1}{a}((m-1)\Delta_f v)^2 + 2HK \frac{|\nabla v|^2}{v} + 2m\nabla v \nabla \bar{F} + (1-\alpha)\left(\frac{v_t}{v}\right)^2 - \alpha' \frac{v_t}{v}. \end{aligned}$$

Since  $L(\alpha^{-1}\bar{F}) = (\alpha^{-1})'\bar{F} + \alpha^{-1}L(\bar{F})$  and  $\alpha' = 2HK\alpha$ , then

$$L(\alpha^{-1}\bar{F}) \leq -\frac{1}{a}\alpha^{-1}((m-1)\Delta_f v)^2 + 2m\alpha^{-1}\nabla v \nabla \bar{F} + (1-\alpha)\alpha^{-1}\left(\frac{v_t}{v}\right)^2. \tag{3.1}$$

By (2.2) and the definition of  $\bar{F}$ , we get

$$(m-1)\Delta_f v = (\alpha^{-1}-1)\frac{|\nabla v|^2}{v} - \alpha^{-1}\bar{F}, \quad \alpha \frac{v_t}{v} = \frac{|\nabla v|^2}{v} - \bar{F}. \tag{3.2}$$

Putting (3.2) into (3.1), we obtain

$$\begin{aligned} L(\alpha^{-1}\bar{F}) &\leq -\frac{(1-\alpha)^2}{a\alpha^3} \frac{|\nabla v|^4}{v^2} - \frac{1}{a\alpha^3}\bar{F}^2 + \frac{2(1-\alpha)}{a\alpha^3} \frac{|\nabla v|^2}{v}\bar{F} + \frac{2m}{\alpha}\nabla v \nabla \bar{F} \\ &\quad + \frac{1-\alpha}{\alpha^3} \frac{|\nabla v|^4}{v^2} + \frac{1-\alpha}{\alpha^3}\bar{F}^2 - \frac{2(1-\alpha)}{\alpha^3} \frac{|\nabla v|^2}{v}\bar{F}. \end{aligned} \tag{3.3}$$

According to (2.4) and (2.5) in [8], we can construct a cut-off function  $\phi$  such that  $0 \leq \phi \leq 1$ ,  $\text{supp}(\phi) \subset B_p(2R)$ ,  $\phi|_{B_p(R)} = 1$  and

$$\frac{|\nabla \phi|^2}{\phi} \leq \frac{C}{R^2}, \quad -\Delta_f \phi \leq \frac{C}{R^2}(1 + R\sqrt{K}), \tag{3.4}$$

where  $C$  is a constant depending only on  $n$ . Set  $G = t\phi\alpha^{-1}\bar{F}$ . Assume that  $G$  achieves its maximum at the point  $(x_0, s) \in B_p(2R) \times [0, T]$  and assume  $G(x_0, s) > 0$ . By the maximum principle, we have

$$\nabla G = 0, \quad L(G) \geq 0, \quad \nabla(\alpha^{-1}\tilde{F}) = -\frac{\alpha^{-1}\tilde{F}}{\phi}\nabla\phi$$

at the point  $(x_0, s)$ , and

$$\begin{aligned} 0 \leq L(G) &= s\phi L(\alpha^{-1}\bar{F}) - (m-1)v \frac{\Delta_f \phi}{\phi} G + 2(m-1)v \frac{|\nabla \phi|^2}{\phi^2} G + \frac{G}{s} \\ &\leq s\phi \left\{ -\frac{1}{a\alpha^3}\bar{F}^2 + \frac{2(1-\alpha)}{a\alpha^3} \frac{|\nabla v|^2}{v}\bar{F} + \frac{2m}{\alpha}\nabla v \nabla \bar{F} + \frac{1-\alpha}{\alpha^3}\bar{F}^2 - \frac{2(1-\alpha)}{\alpha^3} \frac{|\nabla v|^2}{v}\bar{F} \right\} \end{aligned}$$

$$\begin{aligned}
 & -(m-1)v\frac{\Delta_f\phi}{\phi}G + 2(m-1)v\frac{|\nabla\phi|^2}{\phi^2}G + \frac{G}{s} \\
 = & -\frac{1}{a\alpha s\phi}G^2 + \frac{2(1-\alpha)}{a\alpha^2}\frac{|\nabla v|^2}{v}G - 2m\nabla v\frac{\nabla\phi}{\phi}G + \frac{1-\alpha}{\alpha s\phi}G^2 - \frac{2(1-\alpha)}{\alpha^2}\frac{|\nabla v|^2}{v}G \\
 & -(m-1)v\frac{\Delta_f\phi}{\phi}G + 2(m-1)v\frac{|\nabla\phi|^2}{\phi^2}G + \frac{G}{s} \\
 \leq & -\frac{1+a(\alpha-1)}{a\alpha s\phi}G^2 + \frac{2(1-\alpha)}{a\alpha^2}\frac{|\nabla v|^2}{v}G - \frac{2(1-\alpha)}{\alpha^2}\frac{|\nabla v|^2}{v}G \\
 & + 2\frac{mH^{\frac{1}{2}}}{(m-1)^{\frac{1}{2}}}\frac{|\nabla\phi|}{\phi}\frac{|\nabla v|}{v^{\frac{1}{2}}}G - H\frac{\Delta_f\phi}{\phi}G + 2H\frac{|\nabla\phi|^2}{\phi^2}G + \frac{G}{s}.
 \end{aligned} \tag{3.5}$$

Multiplying both sides of (3.5) by  $\frac{a\alpha s\phi}{(1+a(\alpha-1))G}$ , we get

$$\begin{aligned}
 G(x, T) \leq G(x_0, s) \leq & \frac{1}{1+a(\alpha-1)}\left\{\frac{2(1-\alpha)(1-a)}{a\alpha}s\phi\frac{|\nabla v|^2}{v}\right. \\
 & \left.+ 2\frac{mH^{\frac{1}{2}}}{(m-1)^{\frac{1}{2}}}as\phi|\nabla\phi|\frac{|\nabla v|}{v^{\frac{1}{2}}} - Has\alpha\Delta_f\phi + 2Has\alpha\frac{|\nabla\phi|^2}{\phi} + a\alpha\phi\right\}.
 \end{aligned} \tag{3.6}$$

By (3.4), (3.5), and the inequality  $2xy \leq x^2 + y^2$ , similar to the proof of Theorem 1.2 in [6] we have

$$\bar{F}(x, T) \leq \frac{1}{1+a(\alpha-1)}\left\{\left(\frac{bm^2Ha\alpha^4}{2(\alpha-1)(1-a)} + 3Ha\alpha^2\right)\frac{C}{R^2} + Ha\alpha^2\sqrt{K}\frac{C}{R} + \frac{a\alpha^2}{T}\right\}.$$

Since  $T$  is arbitrary, we obtain

$$\bar{F}(x, T) \leq \frac{1}{1+a(\alpha-1)}\left\{\frac{CHa\alpha^2}{R^2}\left(\frac{bm^2\alpha^2}{2(\alpha-1)(1-a)} + 3 + \sqrt{KR}\right) + \frac{a\alpha^2}{T}\right\}.$$

Thus, the proof of Theorem 1.1 is complete. Letting  $R \rightarrow \infty$  in (1.4), we get (1.5). Therefore, we conclude that Corollary 1.2 is true. □

#### 4. The proof of Theorem 1.3

We find that  $\varphi(t) = \frac{a}{t} + aSK + \frac{a}{3}(SK)^2t + \frac{\lambda}{t^2}$  and  $\alpha(t) = 1 + \frac{2}{3}SKt$  satisfy the following equations:

$$\begin{cases} -\frac{2}{t}\varphi + a\left(\frac{1}{t} + HK\right)^2 - \varphi' = 0, \\ 2\left(\frac{1}{t} + HK\right) - \alpha' = \frac{2\alpha}{t}. \end{cases} \tag{4.1}$$

On the other hand, by (2.2) and the definition of  $F$ ,  $\varphi(t)$ , and  $\alpha(t)$  we get

$$\begin{aligned}
 (m-1)\Delta_f v + a\left(\frac{1}{t} + HK\right) &= -\frac{1}{\alpha}\left\{F + \varphi + (\alpha-1)\frac{|\nabla v|^2}{v} - a\alpha\left(\frac{1}{t} + HK\right)\right\} \\
 &= -\frac{1}{\alpha}\left\{F + (\alpha-1)\frac{|\nabla v|^2}{v} - \left(\frac{2a}{3}HK + \frac{a}{3}(HK)^2t - \frac{\lambda}{t^2}\right)\right\}.
 \end{aligned}$$



Therefore, similar to (5.4) in [6] and some discussions in Section 2 and Section 3, we have

$$L(F) \leq -\frac{1}{a\alpha^2} \left\{ F + (\alpha - 1) \frac{|\nabla v|^2}{v} - \left( \frac{2a}{3} HK + \frac{a}{3} (HK)^2 t - \frac{\lambda}{t^2} \right) \right\}^2 - \frac{2}{t} F + 2m \nabla v \nabla F.$$

Construct a cut-off function  $\phi$  as that in Section 3. Define  $G = \beta(t)\phi F$ . Assume that  $G$  achieves its maximum at the point  $(x_0, s) \in B_p(2R) \times [0, T]$  and assume  $G(x_0, s) > 0$ . Similar to (5.6) in [6], we conclude that

$$\begin{aligned} G(x_0, s) &\leq \frac{\alpha\beta}{s} \phi \left\{ \frac{2}{3} (2HKs + (HK)^2 s^2 - \frac{\lambda}{s}) + s\alpha^2 \left( \frac{\beta'}{\beta} - \frac{2}{s} \right) \right\} \\ &\quad + \alpha\alpha^2 \beta H \left\{ -\Delta_f \phi + 2 \frac{|\nabla \phi|^2}{\phi} + \frac{bm^2 \alpha^2}{2(\alpha - 1)} \frac{|\nabla \phi|^2}{\phi} \right\}. \end{aligned} \quad (4.2)$$

Let  $\beta(t) = \tanh(HKt)$ , similar to [6] we have

$$\frac{2}{3} (2HKs + (HK)^2 s^2 - \frac{\lambda}{s}) + s\alpha^2 \left( \frac{\beta'}{\beta} - \frac{2}{s} \right) \leq 0. \quad (4.3)$$

By (3.4), (4.2), and (4.3), we get

$$G(x_0, s) \leq \{ M\alpha^2(T) \left( 3 \frac{C}{R^2} + \frac{C\sqrt{K}}{R} \right) + \frac{CMabm^2\alpha^4(T)}{R^2} \}.$$

Thus, similar to the discussions of (5.7) in [6], we affirm that Theorem 1.3 holds. Letting  $R \rightarrow \infty$  in (1.6), we get (1.7). Therefore, we conclude that Corollary 1.4 is true.

## References

- [1] Brighton K. A Liouville-type theorem for smooth metric measure spaces. *J Geom Anal* 2013; 23: 562–570.
- [2] Charalambous N, Lu ZQ. Heat kernel estimates and the essential spectrum on weighted manifolds. *J Geom Anal* 2013; DOI 10.1007/s12220-013-9438-1.
- [3] Chen L, Chen WY. Gradient estimates for positive smooth  $f$ -harmonic functions. *Acta Math Sci* 2010; 30B: 1614–1618.
- [4] Dung NT. A splitting theorem on smooth metric measure spaces. *Arch Math* 2012; 99: 179–187.
- [5] Futaki A, Li HZ, Li XD. On the first eigenvalue of the Witten-Laplacian and the diameter of compact shrinking solitons. *Ann Glob Anal Geom* 2013; 44: 105–114.
- [6] Huang GY, Huang ZJ, Li HZ. Gradient estimates for the porous medium equations on Riemannian manifolds. *J Geom Anal* 2013; 23: 1851–1875.
- [7] Huang GY, Li HZ. Gradient estimates and entropy formulae of porous medium and fast diffusion equations for the Witten Laplacian. *Pac J Math* 2014; 268: 47–78.
- [8] Huang GY, Ma BQ. Gradient estimates for a nonlinear parabolic equation on Riemannian manifolds. *Arch Math* 2010; 94: 265–275.
- [9] Li XD. Liouville theorems for symmetric diffusion operators on complete Riemannian manifolds. *J Math Pures Appl* 2005; 84: 1295–1361.
- [10] Lu P, Ni L, Vázquez J, Villani C. Local Aronson-Bénilan estimates and entropy formulae for porous medium and fast diffusion equations on manifolds. *J Math Pures Appl* 2009; 91: 1–19.

- [11] Ma L, Du SH. Extension of Reilly formula with applications to eigenvalue estimates for drifting Laplacians. *CR Math Acad Sci Paris* 2010; 348: 1203–1206.
- [12] Ma L, Liu BY. Convex eigenfunction of a drifting Laplacian operator and the fundamental gap. *Pac J Math* 2009; 240: 343–361.
- [13] Wei G, Wylie W. Comparison geometry for the Bakry-Emery Ricci tensor. *J Differ Geom* 2009; 83: 377–405.
- [14] Xia CY, Xu HW. Inequalities for eigenvalues of the drifting Laplacian on Riemannian manifolds. *Ann Glob Anal Geom* 2014; 45: 155–166.
- [15] Yang YY. Gradient estimates for a nonlinear parabolic equation on Riemannian manifolds. *Proc Am Math Soc* 2008; 136: 4095–4102.