

1-1-2011

## Computational comparison of $\beta$ -mannosidases of animals, humans, microbes, and plants

ZAHOOOR QADIR SAMRA

MUHAMMAD AMIN ATHAR

Follow this and additional works at: <https://journals.tubitak.gov.tr/veterinary>



Part of the [Animal Sciences Commons](#), and the [Veterinary Medicine Commons](#)

---

### Recommended Citation

SAMRA, ZAHOOOR QADIR and ATHAR, MUHAMMAD AMIN (2011) "Computational comparison of  $\beta$ -mannosidases of animals, humans, microbes, and plants," *Turkish Journal of Veterinary & Animal Sciences*: Vol. 35: No. 6, Article 4. <https://doi.org/10.3906/vet-0908-52>

Available at: <https://journals.tubitak.gov.tr/veterinary/vol35/iss6/4>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Veterinary & Animal Sciences by an authorized editor of TÜBİTAK Academic Journals. For more information, please contact [academic.publications@tubitak.gov.tr](mailto:academic.publications@tubitak.gov.tr).

## Computational comparison of $\beta$ -mannosidases of animals, humans, microbes, and plants

Zahoor Qadir SAMRA\*, Muhammad Amin ATHAR

Institute of Biochemistry and Biotechnology, Quaid-e-Azam Campus, University of the Punjab,  
Lahore, 54590 - PAKISTAN

Received: 21.08.2009

**Abstract:** The  $\beta$ -mannosidase (MANB) enzyme is involved in removing mannose residue from the nonreducing end, and its impaired activity leads to  $\beta$ -mannosidosis. MANB amino acid sequences of humans, other mammals, plants, fungi, and bacteria were compared to determine their similarities, differences, and predicted 3D structures. Our cloned MANB DNA sequence showed a 99% similarity to a previously reported human MANB DNA sequence but 16 nucleotide differences were observed, showing the polymorphic nature of the enzyme. The 9 changed codons coded the different amino acids *Ile, Lys, Ile, Thr, Arg, Leu, Leu, Gly, and Asp*, while 7 changed codons coded the same amino acids, *Ile, Arg, Gln, Val, Ile, Pro, and Val*. The amino acid sequence comparison of human MANB with bovine, goat, and mouse MANB showed a nearly 75% similarity, while 10%-13%, 17%-18%, and 9%-23% similarities were observed with plant, fungi, and bacteria MANB, respectively. The catalytic nucleophilic and proton donor sites were conserved in the  $\beta$ -mannosidase of mammals, plants, fungi, and bacteria, except *L. esculentum*, and the nucleophilic site of *P. furiosus* was also changed. The catalytic sites of MANB indicated that it follows a dyad catalytic mechanism. Additionally, 2 common putative glycosylation sites at N-residues 35 and 77 were conserved. The 3D structure prediction indicated differences in the  $\alpha$ -helix loop, while the  $\beta$ -pleated sheets were nearly the same. The comparison showed that the MANB enzyme is polymorphic in nature with conserved catalytic sites and has an evolutionary relationship among different species. The 3D structure comparison of MANB will be helpful to understand the disease process of  $\beta$ -mannosidosis.

**Key words:**  $\beta$ -Mannosidase, DNA sequence, amino acid comparison, computational analysis

### Introduction

$\beta$ -Mannosidase (MANB,  $\beta$ -D-mannosidase mannohydrolase, EC 3.2.1.25) is one of the exoglycosidases that cleave the single  $\beta$ -linked mannose residue from the nonreducing end of all N-linked glycoproteins. It is essential for the complete hydrolysis of polysaccharides such as galactoglucomannan and manno oligosaccharides, which are produced by  $\beta$ -mannanase activity. MANB is also important in the saccharification of hemicelluloses to monomeric sugar for further

conversion into biochemical products as well as in the synthesis of oligosaccharides. Oligosaccharides are long chains of sugar molecules used in the building of bones, cartilage, skin, tendons, and many other tissues in the body (1-4). This enzyme also provides useful information for structural studies of polysaccharides and glycoproteins that have  $\beta$ -mannosidic linkages.

MANB enzyme has been purified from fungi (1,5), hyperthermophilic microbes (6,7), plants (8,9), goats (10), bovines (3), and humans (11). The defective

\* E-mail: samra201@hotmail.com

or impaired activity of MANB leads to a lysosomal storage disease commonly known as  $\beta$ -mannosidosis. This disease has been detected in Nubian goats (12) and bovines (2,13), as well as in humans (14,15). Facial dysmorphism, abnormal skull and joints, weakness and shrinkage of muscles, marked intention tremor, and deafness have been reported as pathological symptoms (16). The defective activity of the MANB enzyme leads to the accumulation of mannose-conjugated oligosaccharides or glycoproteins in many tissues, along with the loss of the myelin sheath in the nervous system (17).

In humans, the clinical expressions of  $\beta$ -mannosidosis are reported as peripheral neuropathy (18,19), hearing loss, speech impairment (20), epileptic encephalopathy (21), and angiokeratoma (22). The MANB enzyme activity was completely lost in the fibroblast, plasma, and leukocytes of  $\beta$ -mannosidosis patients (16).

As a continuation of our previous report about the cloning and expression of the MANB gene, there is a need for computational analysis of MANB DNA and its amino acid sequence similarity to or differences from reported MANB DNA and amino acid sequences of other species. The computational analysis will be helpful in finding the evolutionary conservation as well as in understanding the degradation pathway of mannose-linked glycoproteins in the progression of  $\beta$ -mannosidosis in mammals.

## Materials and methods

### Comparison of DNA sequences of human $\beta$ -mannosidase

The cloned DNA sequence of human  $\beta$ -mannosidase (EU009130) was compared with the reported human  $\beta$ -mannosidase DNA sequence (U60337) (15). Both DNA sequences were aligned using the nucleotide-nucleotide BLAST (BLASTN) program of the National Center for Biotechnology Information (NCBI), and the nucleotide differences were noted.

### Comparison of deduced amino acid sequences of $\beta$ -mannosidase

MANB enzyme activity was characterized in the species mentioned below and their NCBI accession

numbers were used to learn the deduced amino acid sequences of MANB from the NCBI. The deduced amino acid sequences of MANB of different species were aligned with deduced amino acid sequences of our cloned human MANB in ClustalW 1.83 and similarities and differences of the amino acid residues were noted.

1) Mammals: human (*Homo sapiens*) (U60337), bovine (*Bos taurus*) (U17432), goat (*Capra hircus*) (U46067), and mouse (*Mus musculus*) (AF306557).

2) Fungi: *Aspergillus niger* (AJ251874) and *Aspergillus aculeatus* (AB015509).

3) Plants: Arabidopsis (*Arabidopsis thaliana*) (AB122060), cotton (*Gossypium hirsutum*) (AY187062), and tomato (*Lycopersicon esculentum* Mill.) (AF415204).

4) Bacteria: *Cellulomonas fimi* (AF126472), *Thermotoga neapolitana* (AY033395), *Thermobifida fusca* (CAD33708), *Pyrococcus furiosus* (U60214), *Streptomyces coelicolor* (NP\_630333), *Xylella fastidiosa* (NP\_298136), and *Thermotoga maritime* (NP\_229424).

### Prediction of 3-dimensional structures

The MANB amino acid sequences of human, bovine, goat, mouse, *Cellulomonas fimi*, *Thermotoga neapolitana*, *Pyrococcus furiosus*, *Streptomyces coelicolor*, and *Thermotoga maritime* were inserted into the ExpASY proteomics server and the 3-dimensional structures were predicted.

## Results

### Sequence comparison

#### Human

The cloned human MANB DNA sequence was aligned with the reported human MANB cDNA sequence. A 99% DNA sequence similarity was observed, but 16 nucleotide differences in the open reading frame of  $\beta$ -mannosidase DNA were identified. The positions of nucleotide differences are shown in the Table. Due to a change in triplet codons, it was noted that 7 triplet codons with changed nucleotide positions at 390, 479, 486, 489, 757, 1011, and 1137 maintained the same amino acid residues as *Ile*, *Arg*, *Gln*, *Val*, *Ile*, *Pro*, and *Val*, respectively. In

Table. Computational comparison of DNA sequences of the human  $\beta$ -mannosidase gene and deduced amino acid sequences.

No.	Nucleotide position	Alkhatay (U60337)	Amino acid residue triplet codon	Samra (EU009130)	Amino acid residue triplet codon
1.	208	<b>CAC</b>	His	TAC	Tyr
2.	390	ATT	Ile*	ATC	Ile*
3.	479	CSC	Arg*	CGC	Arg*
4.	486	<b>CAG</b>	Gln*	CAA	Gln*
5.	489	GTT	Val*	GTC	Val*
6.	757	NTA	Val/Ile*	ATA	Ile*
7.	862	<b>GAA</b>	Glu	AAA	Lys
8.	889	AAC	Asn	ATC	Ile
9.	1011	CCT	Pro*	CCC	Pro*
10.	1037	AAT	Asn	ACT	Thr
11.	1039	<b>GGA</b>	Gly	CGA	Arg
12.	1130	<b>CAG</b>	Gln	CTG	Leu
13.	1137	GTT	Val*	GTA	Val*
14.	1598	TTT	Phe	CTT	Leu
15.	1766	<b>GAA</b>	Glu	GGA	Gly
16.	2590	AAT	Asn	GAT	Asp

Bold letters indicate nucleotide differences. Asterisk indicates common amino acid residue.

the 9 remaining triplet codons, the first nucleotide position of the triplet codons was changed at 208, 862, 1039, 1130, 1598, and 2590, while the second nucleotide position was changed at 889, 1037, and 1766 (Table). The positions of the changed amino acids numbers are shown (Figure 1).

#### Other mammals

The comparison of deduced amino acid sequences of human  $\beta$ -mannosidase with MANB of mouse, bovine, and goat indicated that a 74.85% sequence similarity was observed with bovine and mouse MANB, whereas a 74.63% similarity was observed with goat MANB. The proposed catalytic nucleophilic site (Glu-554) and proton donor site (Glu-457) of human  $\beta$ -mannosidase was suggested after alignment with an amino acid sequence of *A. thaliana*  $\beta$ -mannosidase (9). The common potential

glycosylation sites of MANB were observed at N-residues 35 and 57 of goat, human, and bovine, but not in mouse, while glycosylation at residues 28, 280, 284, and 763 was only observed in human MANB. The glycosylation at residue 297 is conserved among bovine, mouse, human, and goat.

#### Plant

The alignment of deduced amino acids of human  $\beta$ -mannosidase with reported amino acids sequences of MANB of *G. hirsutum* and *A. thaliana* showed a 13.42% and a 13.69% similarity, respectively, while it showed a 10.70% similarity with tomato  $\beta$ -mannosidase. The catalytic nucleophilic and proton donor sites of  $\beta$ -mannosidase are conserved in human, cotton, and Arabidopsis, and was not found conserved in *L. esculentum*  $\beta$ -mannosidase (Figure 2).

Samra 1 253 288 297 346 347 377 533 589 864 879  
**Met-----I-----K-----I-----T-----R-----L-----L-----G-----D-----stop**  
 -----  
 Alkhyat **Met-----V/I-----E-----N-----N-----G-----Q-----F-----E-----N-----stop**

Figure 1. Alignment and comparison of deduced amino acid sequences of human  $\beta$ -mannosidase (EU009130) and reported human  $\beta$ -mannosidase (Alkhyat, U60337). The amino acids differences are represented by numbers with 1-letter amino acid abbreviations. Dashes indicate the same amino acid sequence. The 9 changed amino acid residues are shown.

Amino acid	1	48	55	136	184	185	236	282	295	297	299	328	380	390
residue no	1	48	55	134	181	182	233	279	292	294	296	325	377	387
of species	1	45	5	119	178	179	224	266	280	282	284	312	369	379
	1	6	13	54	62	63	79	100	114	116	118	130	173	183

<i>G.hirsutum</i>	M--P--L---T---W---D---E---F---P--G---G---R---D---G
<i>A.thaliana</i>	M--P--L---T---W---D---E---F---P--G---G---R---D---G
SMANB	M--P--L---T---W---D---E---F---P--G---G---R---D---G
<i>L.esculentum</i>	M--P--L---T---W---D---E---F---P--G---G---R---D---G
	* * * * * * * * * * * * * *

Amino acid	399	414	453	455	465	558	597	623	909	933	976
residue no	396	411	450	452	464	549	588	612	876	897	944
of species	388	403	434	435	457	554	578	592	803	827	879
	192	207	240	242			334	349	452	476	514

<i>G.hirsutum</i>	F---F---H---S--- <b>E</b> --- <b>E</b> ---N---P---F---D-----L
<i>A.thaliana</i>	F---F---H---S--- <b>E</b> --- <b>E</b> ---N---P---F---D-----G
SMANB	F---F---H---S--- <b>E</b> --- <b>E</b> ---N---P---F---D-----Y
<i>L.esculentum</i>	F---F---H---S---N---P---F---D-----H
	* * * * * <b>▲</b> <b>▲</b> * * * *

Figure 2. Computational comparison of deduced amino acid sequences of human and plant  $\beta$ -mannosidases. SMANB represents cloned human  $\beta$ -mannosidase (EU009130). Amino acid sequence alignment of human with reported plant  $\beta$ -mannosidase of *A. thaliana* (AB122060), *G. hirsutum* (AY187062), and *L. esculentum* (AF415204) was done using ClustalW 1.83. The common conserved aligned amino acid sequence is shown by an asterisk. The proposed catalytic nucleophilic (*Glu-549*) and proton donor (*Glu-464*) sites of plant endo- $\beta$ -mannosidase of *A. thaliana* (9) are marked with a bolded “E” and shown by an arrowhead. These sites were used to compare the catalytic sites of other  $\beta$ -mannosidases. Dashes indicate the nonconserved amino acid sequences.

### Fungi

The comparison of the deduced amino acid sequence of human MANB with the reported amino acid sequences of  $\beta$ -mannosidase of *A. niger* and *A. aculeatus* showed 18.99% and 17.86% similarity, respectively (Figure 3). The first amino acid after the start codon “M” is “R”, which resembles the amino acid sequence of human  $\beta$ -mannosidase and suggests the

pattern of the eukaryotic translation signal sequence (23). One glycosylation site was also found conserved in the  $\beta$ -mannosidase of *A. niger*, *A. aculeatus*, and human. The proposed catalytic nucleophilic and proton donor sites are also conserved.

### Bacteria

The deduced amino acid sequences of bacterial  $\beta$ -mannosidases showed conserved proton donor and

*A.niger* MRHSIGLAAALLAPTLPLVALG--QHIRDLSSEKWTLSRRLNRTVPAQFPSQVHLDLLRA 58  
*A.aculaetus* MRALPTTATLLGLVFFPSASRSQYVRDLGTEQWTLSSATLNRTVPAQFPSQVHMDLLRE 60  
SMANB MRLHLLLLLLALCGAGTTAAELS-----YSLRGNWSICNGNGSLELPGAVPGCVHSALFQQ 55  
\*\* : \* . : : \* : . . : \* . \* . \* \* \* :

*A.niger* GVIGEYHG-LNDFNLRWIAAANWTTYT-SQPIKGLLDNYGSTWLVFDGLDTFATISILWTA 58  
*A.aculaetus* GIIDEPYNDLNDNFNLRWIADANWTTYT-SGKIEGLGEDYESTWLVFDGLDTFASISFCGQF 59  
SMANB GLIQDSYYRFNDLNYRWVSLDNWTTYSKEFKIPFEISKWQKVNLIILEGVDTVSKILFNEVT 60  
\* : \* : : : \* : \* \* \* : : \* \* \* : . \* . . : . . \* : : \* : \* : \* : \* : \* : :

*A.niger* NRIHQSVSPVSGSMYLPALAEAC-QRRILIRKVSFRGGVTAEVNTCYLHIEWPDDVQLTLY 59  
*A.aculaetus* VGATDNQFRQYMFVSSILKACP-EEPTLGIQFGSAPNIVDAIAQDPSSPTWPEGVQITY 59  
SMANB IGETDNMFNRYSFEDITNVVRDVNSIELRFQSAVLYAAQQSKAHTRYQVPPDCPPLVQKGE 60  
. : . . : : . : . . : . : . : \* \* \*

*A.niger* EYPNRWFMRKEQSDFGWDWGPAPAGPWKPAYIVQLDKKESVYVLTNDLDIRKNQINY 60  
*A.aculaetus* EYPNRWFMRKEQSDFGWDWGPAPAGPWKPGYVVLKQAAPVYVNTDLDIRLQINY 60  
SMANB CHVN--FVRKEQCSFSWDWGPSFPTQGIWKDVRIEAY-----NICHLYFTFSPIY 49  
: \* \* : \* \* \* \* . \* . \* \* \* \* : \* \* \* \* : \* \* \* \* . \*

*A.niger* LPPDQSQPVVNASIDILGPLPAKPTMSIEVRDTHSGTILTSRTLNNVSVAGNAITGVTV 60  
*A.aculaetus* LPPDQTPVNVNASLDYLGSLPENPSMAIEVKDLQSGEILASRPLTNITVTEGSVTGVTV 60  
SMANB DKSAQEWNLIEIESTFDVVSSKPVGGQVIAIPKLQTTQTYSELQPGKRIVELFVN--I 57  
. \* : : : \* : . . \* : \* : . . : : . . . : . . : . . :

*A.niger* LDGLNPKLWWPQSSVIRTSTMFLSLKVEGTRPWPVWTNGRASAPFFLNQRNITEVQRAQ 60  
*A.aculaetus* LEGVDPKLLWWPQGLGDQNLNVYTI SVTDGGNQSVAEVTKRTGFRTIFLNQRNITDAQLAQ 60  
SMANB SKNITVKTWWPHGHGIQTGYNMTVLFELDG-----GLNIEKSAKVYFRTVELIEEPI 52  
. . : \* \* \* : . . : . \* . : . : \* : : . . :

*A.niger* GIAPGANWHFEVNGHEFYAKGSNLI PPDSFWTRVTEERISRLFDVAVVGNQNMLRVWSSG 60  
*A.aculaetus* GIAPGANWHFEVNGHEFYAKGSNLI PPDCFWTRVTEEDTMTRLFDVAVVAGNQNMLRVWSSG 60  
SMANB KGSPGLSFYFKITRFPIFLKGSNWI PADS FQDRVTSELLRLLLLSVVDANMNTLRVWGGG 60  
: \* \* : : \* : . . : : \* \* \* \* \* \* \* \* \* \* : : \* : : \* \* \* \* \* \* \* \*

*A.niger* AYLHDYIYDLADEKGI LLWSEFEFS DALYPSDDAFLENVA AEIVYVRRVNHHP SLALWA 60  
*A.aculaetus* AYLHDYIYDLADEKGI LLCEFEFS DALYPTDDAFLENVA AEVYVRRVNHHP SLALWA 60  
SMANB IYEQDEFYELCDELGIMVWQDFMFACALYPTDQGF LDSVTAEVAYQIKRLKSHPSII IWS 60  
\* : \* : \* : \* \* \* \* : . : \* \* : \* \* \* \* \* : \* \* \* \* : \* \* \* \* : \* \* \* \* : \* \* \* \* :

▼  
*A.niger* GGNEIESLMLPRVKDAAPSSYSYVGEY EKMYISLFLPLVYENTRSISYSPSSTTEGYLY 60  
*A.aculaetus* GGNEIESLMLLLVEAADPESYPFYVGEY EKMYISLFLPLVYENTRSISYSPSSTTEGYLD 60  
SMANB GNNENEALMMNWHYHISFTDRPIYIKDYVTLYVKNIRELVLAGDKSRPFITSSPTNG--- 57  
\* . \* \* \* . : : . . \* : \* : \* : . : \* \* . : \* : . : \* \* \* \* \* \*

▼  
*A.niger* IDLSAPVPM AERYDNTTSGSYGDTDH YDYSVAFDYGSYPVGRFANEFGFHSMP SLQT 60  
*A.aculaetus* IDLSAPVPM AERYSNTEGEY YGDTDH YNYDASIAFDYGTYPVGRFANEFGFHSMP SLQT 60  
SMANB ----AETVAEAWVSQNPNSNYFGDVHLYDYIS-DCWNWKFVFKARFAS EYGYQSWPSFST 56  
\* . . : . . . . \* : \* \* . \* : \* : . : : : \* \* . \* \* \* : \* \* \* \* \* \*

*A.niger* WQQAVDT-EDLYFNSSVVMLRNHHPAGGLMTDNYANSATGMGEMTMGVISYYPISKSD 59  
*A.aculaetus* WQQALTDPADLTFNSSVVMLRNHHYPAGGLMTDNYHNTVARHGRNDPGRAGLLPDAQHSV 60  
SMANB LEKVSST-EDWSFN SKFSLHRQHGGGN-----KQMLYQAGLHF KLPQST 44  
: : . \* \* \* \* . : \* \* \* \* . . . . . : \* :

*A.niger* HISSN-FSAWCHATQLFQADMYKSQIQFYRRGSGMPERQ----LGSLYWQLEDIWQAPSWA 55  
*A.aculaetus* RPRGQLQRLVPRDPALPGGPLQVTNPVLPAGQRAARTP----ARVPVLAARGHLA GALVG 56  
SMANB DPLRTFKDTIYLTQVMQAQC VKTETEFYRRSRSEIVDQQGHTMGALYWQLNDIWQAPSWA 60  
: . : . : . . . . . : . . . . .

Figure 3.

```

A. niger      GIEYGGRWKVLHHVMRDIYQPVI VSPFWN-YTTGSLDVYVTS DLWSPAAGTVDLTWLDLS 59
A. aculaetus GDRVRRPLEGPHYVARDIYKPVIVSPFWN-YTTGALDIYVTS DLWTAAGSVTLTWRDLS 59
SMANB         SLEYGGKWKMLHYFAQNFFAPLLPVGFENENTFYIYGVSDLHSDYSMTLSVRVHTWSSLE 60
. .           : * : . : : : * : : * * * . : . : : . * * . * .

A. niger      GRPIAGNAGTP-KSVPFTVGGLNSTR IYGTNVSSLGLPDTKDAVLILSLSAHGRLPNSDR 59
A. aculaetus GKPIASNGGLPTKPLPFHVGALNSTR LYMNMKQQPLPRHEDAILALELTATGSLPNTDE 60
SMANB         PVCSR-----VTERFVMKGG EAVCLYE EEPVS--ELLRRCGNCTRESCVVSFYLSADHE 51
. * : . : : . : * : . * . . . * . .

A. niger      TTNLTHENYATLSWPKDLKIVDPGLK LGYSSKKTTVTVEATSGVSLYTWLDYPEGVVGYF 60
A. aculaetus EVTFTHEQWFTPAFPKDLDLVNLRV RVEYDAPLGKFAVEATAGVALYTWLEHPEGVVGYF 60
SMANB         LLSPTNYHFLSSPKEAVGLCKAQITAI ISQQGDI FVFDLETSAVAPFVWLDVGS-IPGRF 59
. * : : : . : . . * : * : : * : . : * *

A. niger      EENAFVLAPGEKKEIGFTVLDDTTNGAWVRNITVQSLWDQKVRG 43
A. aculaetus EENSFVVVPGQKKVVG FVVQADETDGEWVHDVTVRSLWDLNEGE 43
SMANB         SDNGFLMTEKTR-TILFYPWEPTSKDELEQSFHVTSLTDIY--- 39
.:* . : . : : : : : : . . * : * :

```

Figure 3. Amino acid sequence alignment of human  $\beta$ -mannosidase (SMANB, EU009130) with *A. niger* and *A. aculeatus* was done using ClustalW 1.83. The asterisk indicates a conserved amino acid sequence, (.) indicates conserved substitution, and (:.) indicates semiconserved substitution. The proposed catalytic nucleophile (*Glu-554*) and proton donor (*Glu-457*) sites based on the human  $\beta$ -mannosidase are marked with a bold “E” and shown by an arrowhead. One common glycosylation site is marked with a bold “N.”

catalytic nucleophilic sites, except in *P. furiosus*. DNA sequences of  $\beta$ -mannosidase in *L. esculentum* and *P. furiosus* were observed to be shorter as compared to mammalian  $\beta$ -mannosidase. It was also noted that in hyperthermophilic *P. furiosus*, the proton donor site at 320 is comparable with the mammalian proton donor site at amino acid residue 457, but the catalytic nucleophilic site of *P. furiosus* changed and shifted at amino acid residue 395 (Figure 4).

### 3D structure

The predicted 3-dimensional structures of MANB enzymes were divided into left (L), center (C), and right (R) parts. The comparison of the 3D structures of MANB enzymes of mammals indicated that in part L, ‘a’ has the same type of  $\beta$ -pleated sheets. In part C, an additional loop of  $\alpha$ -helix structures was observed in bovine and goat, while mouse resembles the 3D structure of human MANB enzyme (Figure 5). Part L of the mammals is different from part L of bacteria. Further comparison of the 3D structure of human MANB with the 3D structure of MANB of *C. fimi* and *P. furiosus* indicated an extra extended loop of  $\beta$ -pleated sheets in *C. fimi* and 2 extra  $\beta$ -pleated sheets in *P. furiosus*. The numbers of  $\alpha$ -helix turns

in part C are also different between mammalian and bacterial MANB enzymes.

### Discussion

$\beta$ -Mannosidase is an important enzyme in the degradation pathway of mannose polymers and mannose-linked glycoproteins. The deduced amino acid sequence of human  $\beta$ -mannosidase was aligned with the  $\beta$ -mannosidases of animals, plants, and microbes. A 75% similarity of deduced amino acid sequence was observed between human and animal  $\beta$ -mannosidases, while a 10%-23% similarity was observed when comparing human  $\beta$ -mannosidase with plant and microbe  $\beta$ -mannosidase. The catalytic nucleophilic and proton donor sites and 2 glycosylation sites are conserved among the  $\beta$ -mannosidases of all species. In the 3D structures, a difference in  $\alpha$ -helix structures was observed while the  $\beta$ -pleated sheets were nearly the same.

The  $\beta$ -mannosidases of all species are active at an acidic pH (5.0 to 5.5), except that of *P. furiosus*, which shows enzyme activity at pH levels of 7.2 to 7.3. The glycosylation sites of  $\beta$ -mannosidases are

Amino acid	1--29--48-78-120-184-289-423-435-455-457-501--554
residue no	1--10--29-59-110-164-255-384-397-417-419-460--505
	1--10--29-59-110-164-255-381-397-417-419-460--505
	1--21--48-78-128-183-280-409-421-441-443-484--531
	1--11--28-58-114-165-257-393-405-425-427-468--515
	1--17--41-71-120-175-265-396-408-428-430-471--518
	1--38--68-98-149-204-313-442-454-474-476-526--574
	1-----10-40--74-105-163-285-297-318-320-361-----
SMANB	M---W---V---W---D---W---P---P---E---N--- <b>E</b> ---P----- <b>E</b>
<i>T.neapolitana</i>	M---W---V---W---D---W---P---P---E---N--- <b>E</b> ---P----- <b>E</b>
<i>T.maritimia</i>	M---W---V---W---D---W---P---P---E---N--- <b>E</b> ---P----- <b>E</b>
<i>T.Fusca</i>	M---W---V---W---D---W---P---P---E---N--- <b>E</b> ---P----- <b>E</b>
<i>S.coelicolor</i>	M---W---V---W---D---W---P---P---E---N--- <b>E</b> ---P----- <b>E</b>
<i>C.Fimi</i>	M---W---V---W---D---W---P---P---E---N--- <b>E</b> ---P----- <b>E</b>
<i>X.fastidiosa</i>	M---W---V---W---D---W---P---P---E---N--- <b>E</b> ---P----- <b>E</b>
<i>P.furiosus</i>	M-----V---W---D---W---P---P---E---N--- <b>E</b> ---P-----
Amino acid	556--653--657--671 879
residue no	507--591--595--599 787
	507--591--585--599 785
	533--617--621--625 840
	517--601--605--609 820
	519--603--607--611 823
	576--662--666--670 891
	391--462--466--470 510
SMANB	G----G----W----D----Y
<i>T.neapolitana</i>	G----G----W----D----R
<i>T.maritimia</i>	G----G----W----D----R
<i>T.Fusca</i>	G----G----W----D----V
<i>S.coelicolor</i>	G----G----W----D----A
<i>C.Fimi</i>	G----G----W----D----H
<i>X.fastidiosa</i>	G----G----W----D----Q
<i>P.furiosus</i>	G- <b>E</b> --G----W----D----G



Figure 4. Computational comparison of a deduced amino acid sequence of human MANB with bacterial  $\beta$ -mannosidases. SMANB represents cloned human  $\beta$ -mannosidase (EU009130). Amino acid sequence alignment was done using ClustalW 1.83. The common conserved aligned amino acid sequences and the amino acid residue numbers are shown. The proposed catalytic nucleophilic (*Glu-554*) and proton donor (*Glu-457*) sites based on the human  $\beta$ -mannosidase are marked with a bold "E." Dashes indicate the nonidentical amino acid sequences. The nucleophilic catalytic site of *P.furiosus* is shifted at amino acid residue 395, shown by an arrowhead.

conserved in mammals but different among species. In mammals, a disorder of the  $\beta$ -mannosidase enzyme leads to  $\beta$ -mannosidosis, a lysosomal storage disease. The defective activity of MANB has not yet been studied in mammals other than goat, bovine, and human. In human  $\beta$ -mannosidosis, the splice mutation of A to G in intron 15 and the 4 base insertions of ATAA between exon 7 and 8 disturb the normal transcription process and null mutation (15,18). In humans, the amino acid sequence of MANB after methionine (the start codon) has the

characteristic eukaryotic signal peptide sequence (24), which is in agreement with bovine  $\beta$ -mannosidase. The translation, maturation, and glycosylation processes of the human  $\beta$ -mannosidase enzyme are not completely known, but they can be compared with bovine  $\beta$ -mannosidase, which indicates the cleavage of the signal peptide at residue 17 following the (-3, -1) rule (24). The hydrophobic region in the deduced amino acid sequence of human MANB can be compared with the bovine  $\beta$ -mannosidase amino acid sequence, which is predicted at amino




Figure 5. Prediction of 3-dimensional models of the  $\beta$ -mannosidase enzyme on the basis of amino acid sequence using ExPASy. Asterisk indicates an extra  $\alpha$ -helix in bovine and goat, (▶) indicates an extra loop of  $\beta$ -pleated sheet in *C. fimi* and *P. furiosus*, and a double arrow indicates fewer  $\alpha$ -helices in part C of the microbial MANB.

acid residues 96-114 and 406-422 and is based on the Kyte-Doolittle hydrophathy plot (25).

The biochemical functions of proteins can be best understood by predicting their 3-dimensional structures on the basis of homology modeling. The numbers of protein structures being determined today are fewer than the numbers of known protein sequences. It is estimated that during 2008, nearly 53,000 experimentally proven protein structures were deposited in the Protein Data Bank (26). This is relatively fewer than the 6 million protein sequences admitted into the Uniport knowledge database (27). Generally, the structural similarity between 2 homologous proteins is determined by

the similarities of the amino acid sequences, which leads to the determination of the 3D structure of the proteins. On the basis of this homology modeling, the 3D structure of MANB of different species (prokaryotes to eukaryotes) was determined. The 3D structure provides fruitful insight for understanding the catalytic activity and mode of enzymatic reaction under different normal and pathological conditions. In MANB homology modeling, the open reading frames (ORFs) of species were used. The genomic ORFs of other species, such as *Saccharomyces cerevisiae*, *E. coli*, *M. genitalium*, *C. elegans*, and *M. jannaschii*, were used to identify the 3D structures. The homology among different sequences varied from

16% to 21% (28). The main advantages of 3D modeling are that it describes the right way of accepting or rejecting the match, that it can provide the active and binding sites of the proteins or enzymes, and that it can directly predict the functional properties of the enzymes or proteins that are not possible to elucidate only on the basis of amino acid sequence homology.

The  $\beta$ -mannosidosis disease is not completely understood due to lack of information about the structure and function of the  $\beta$ -mannosidase enzyme.

## References

- Ademark, P., Lundqvist, J., Hagglund, P., Tenkanen, M., Torto, N., Tjerneld, F., Stalbrand, H.: Hydrolytic properties of a beta-mannosidase purified from *Aspergillus niger*. *J. Biotechnol.*, 1999; 75: 281-289.
- Jones, M.Z., Rathke, E.J., Gage, D.A., Costello, C.E., Murakami, K., Ohta, M., Matsuura, F.: Oligosaccharides accumulated in the bovine beta-mannosidosis kidney. *J. Inherit. Metab. Dis.*, 1992; 15: 57-67.
- Sopher, B.L., Traviss, C.E., Cavanagh, K.T., Jones, M.Z., Friderici, K.H.: Bovine kidney beta-mannosidase: purification and characterization. *Biochem. J.*, 1983; 289: 343-347.
- Percheron, F., Foglietti, M.J., Bernard, M., Ricard, B.: Mammalian beta-D-mannosidase and beta-mannosidosis. *Biochimie*, 1992; 74: 5-11.
- Kurakake, M., Komaki, T.: Production of beta-mannanase and beta-mannosidase from *Aspergillus awamori* K4 and their properties. *Curr. Microbiol.*, 2001; 42: 377-380.
- Duffaud, G.D., McCutchen, C.M., Leduc, P., Parker, K.N., Kelly, R.M.: Purification and characterization of extremely thermostable beta-mannanase, beta-mannosidase, and alpha-galactosidase from the hyperthermophilic eubacterium *Thermotoga neapolitana* 5068. *Appl. Environ. Microbiol.*, 1997; 63: 169-177.
- Bauer, M.W., Bylina, E.J., Swanson, R.V., Kelly, R.M.: Comparison of a beta-glucosidase and a beta-mannosidase from the hyperthermophilic archaeon *Pyrococcus furiosus*. Purification, characterization, gene cloning, and sequence analysis. *J. Biol. Chem.*, 1996; 271: 23749-23755.
- Mo, B., Bewley, J.D.: Beta-mannosidase (EC 3.2.1.25) activity during and following germination of tomato (*Lycopersicon esculentum* Mill.) seeds. Purification, cloning and characterization. *Planta*, 2002; 215: 141-152.
- Ishimizu, T., Sasaki, A., Okutani, S., Meada, M., Yamagishi, M., Hase, S.: Endo-beta-mannosidase, a plant enzyme acting on N-Glycan. Purification, molecular cloning and characterization. *J. Biol. Chem.*, 2004; 279: 38555-38562.
- Pearce, R.D., Callahan, J.W., Novak, A., Little, P.B., Clarke, J.T.: Properties of partially purified goat kidney beta-D-mannosidase. *Br. Vet. J.*, 1990; 146: 270-280.
- Guadalupi, R., Bernard, M., Orlacchio, A., Foglietti, M.J., Emiliani, C.: Purification and properties of human urinary beta-D-mannosidase. *Biochem. Biophys. Acta*, 1996: 1293; 9-16.
- Jones, M.Z., Cunningham, J.G., Dade, A.W., Alessi, D.M., Mostosky, U.V., Vorro, J.R., Benitez, J.T., Lovell, K.L.: Caprine beta-mannosidosis: clinical and pathological features. *J. Neuropathol. Exp. Neurol.*, 1983; 42: 268-285.
- Jolly, R.D., Thompson, K.G., Bayliss, S.L., Vidler, B.M., Healy, P.J.: beta-Mannosidosis in a Salers calf: a new storage disease of cattle. *N.Z. Vet. J.*, 1990; 38: 102-105.
- Levade, T., Graber, D., Flurin, V., Delisle, M.B., Pieraggi, M.T., Testut, M.F., Carriere, J.P., Salvayre, R.: Human beta-mannosidase deficiency associated with peripheral neuropathy. *Ann. Neurol.*, 1994; 35: 116-119.
- Alkhayat, A.H., Kraemer, S.A., Leipprandt, J.R., Macek, M., Kleijer, W.J., Friderici, K.H.: Human beta-mannosidase cDNA characterization and first identification of a mutation associated with human beta-mannosidosis. *Hum. Mol. Genet.*, 1998; 7: 75-83.
- Bryan, L., Schmutz, S., Hodges, S.D., Snyder, F.F.: Bovine beta-mannosidase deficiency. *Biochem. Biophys. Res. Commun.*, 1990; 173: 491-495.
- Lovell, K.L., Kranich, R.J., Cavanagh, K.T.: Biochemical and histochemical analysis of lysosomal enzyme activities in caprine beta-mannosidosis. *Mol. Chem. Neuropathol.*, 1994; 21: 61-74.
- Uchino, Y., Fukushige, T., Yotsumoto, S., Hashiguchi, T., Taguchi, H., Suzuki, N., Konohana, I., Kanzak, T.: Morphological and biochemical studies of human beta-mannosidosis: identification of a novel beta-mannosidase gene mutation. *Br. J. Dermatol.*, 2003; 149: 23-29.
- Sedel, F., Friderici, K., Nummy, K., Caillaud, C., Chabli, A., Durr, A., Lubetzki, C., Agid, Y.: Atypical Gilles de la Tourette Syndrome with beta-mannosidase deficiency. *Arch. Neuro.*, 2006; 63: 129-131.
- Poenaru, L., Akli, S., Rocchiccioli, F., Eydoux, P., Zamet, P.: Human beta-mannosidosis: a 3-year-old boy with speech impairment and emotional instability. *Clin. Genet.*, 1992; 41: 331-334.

21. Cooper, A., Wraith, J.E., Savage, W.J., Thornley, M., Noronha, M.J.: Beta-mannosidase deficiency in a female infant with epileptic encephalopathy. *J. Inherit. Metab. Dis.*, 1991; 14: 18-22.
22. Rodriguez-Serna, M., Botella-Estrada, R., Chabas, A., Coll, M.J., Oliver, V., Febrer, M.I., Aliaga, A.: Angiokeratoma corporis diffusum associated with beta-mannosidase deficiency. *Arch. Dermatol.* 1996; 132: 1219-1222.
23. Kozak, M.: Compilation and analysis of sequence upstream from the translation start site in eukaryotic mRNA. *Nucleic Acid Res.*, 1984; 12: 857-872.
24. von Heijne, G. A new method for predicting signal sequence cleavage site. *Nucleic Acid Res.*, 1986; 14: 4683-4690.
25. Kyte, J., Doolittle, R.: A simple method for displaying the hydropathic character of a protein. *J. Mol. Biol.*, 1982; 157: 105-132.
26. Berman, H., Henrick, K., Nakamura, H., Markley, J.L.: The worldwide Protein Data Bank (wwPDB): ensuring a single, uniform archive of PDB data. *Nucleic Acid Res.* 2007; 35: D301-D303.
27. Wu, C.H., Apweiler, R., Bairoch, A., Natale, D.A., Barker, W.C., Boeckmann, B., Ferro, S., Gasteiger, E., Huang, H., Lopez, R., Magrane, M., Martin, M.J., Mazumder, R., O'Donovan, C., Redaschi, N., Suzek, B. The Universal Protein Resource (Uniport): an expanding universe of protein information. *Nucleic Acid Res.*, 2006; 34: D187-D189.
28. Mizuguchi, K., Deane, C.M., Blundell, T.L., Overington, J.P.: HOMSTRAD: a database of protein structure alignments for homologous families. *Protein Sci.* 1998; 7: 2469-2471.