

1-1-2014

Seiberg–Witten-like equations on 5-dimensional contact metric manifolds

NEDİM DEĞİRMENÇİ

ŞENAY BULUT

Follow this and additional works at: <https://dctubitak.researchcommons.org/math>



Part of the [Mathematics Commons](#)

Recommended Citation

DEĞİRMENÇİ, NEDİM and BULUT, ŞENAY (2014) "Seiberg–Witten-like equations on 5-dimensional contact metric manifolds," *Turkish Journal of Mathematics*: Vol. 38: No. 5, Article 2. <https://doi.org/10.3906/mat-1303-34>

Available at: <https://dctubitak.researchcommons.org/math/vol38/iss5/2>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Mathematics by an authorized editor of TÜBİTAK Academic Journals.

Seiberg–Witten-like equations on 5-dimensional contact metric manifolds**

Nedim DEĞİRMENCİ, Şenay BULUT*

Department of Mathematics, Faculty of Science, Anadolu University, Eskişehir, Turkey

Received: 18.03.2013 • Accepted: 26.02.2014 • Published Online: 01.07.2014 • Printed: 31.07.2014

Abstract: In this paper, we write Seiberg–Witten-like equations on contact metric manifolds of dimension 5. Since any contact metric manifold has a Spin^c -structure, we use the generalized Tanaka–Webster connection on a Spin^c spinor bundle of a contact metric manifold to define the Dirac-type operators and write the Dirac equation. The self-duality of 2-forms needed for the curvature equation is defined by using the contact structure. These equations admit a nontrivial solution on 5-dimensional strictly pseudoconvex CR manifolds whose contact distribution has a negative constant scalar curvature.

Key words: Seiberg–Witten equations, spinor, Dirac operator, contact metric manifold, self-duality

1. Introduction

Seiberg–Witten equations were defined on 4-dimensional Riemannian manifolds by Witten in [14]. The solution space of these equations gives differential topological invariants for 4-manifolds [1, 11]. Some generalizations were given later on higher dimensional manifolds [4, 7, 10].

Seiberg–Witten equations consist of 2 equations. The first is the Dirac equation, which is meaningful for the manifolds having Spin^c -structure. The second is the curvature equation, which couples the self-dual part of a connection 2-form with a spinor field. In order to be able to write down the curvature equation, the notion of the self-duality of a 2-form is needed. This notion is meaningful for 4-dimensional Riemannian manifolds. On the other hand, there are similar self-duality notions for some higher dimensional manifolds [5, 13]. In the present paper, we propose Seiberg–Witten-like equations for 5-dimensional contact metric manifolds by using the Spin^c -structure and the notion of self-duality given in [12] and [3], respectively.

The paper is organized as follows. We begin with a section introducing some basic facts concerning contact metric manifolds. In the following section, we study self-dual 2-forms on 5-dimensional contact metric manifolds. In Section 4, we discuss the Spin^c -structures and Dirac-type operators associated to the generalized Tanaka–Webster connection. In the final section we propose the Dirac and curvature equations and hence write Seiberg–Witten-like equations on contact metric manifolds of dimension 5. Finally, we obtain a special solution for these equations on the 5-dimensional strictly pseudoconvex CR manifolds whose contact distribution has a negative constant scalar curvature.

*Correspondence: skarapazar@anadolu.edu.tr

2010 AMS Mathematics Subject Classification: 15A66, 58Jxx.

** This paper was supported by Project 1105F099 of the Scientific Research Foundation of Anadolu University.

2. Contact metric manifolds

A contact form on a smooth manifold M of dimension $(2n+1)$ is a 1-form η such that $\eta \wedge (d\eta)^n \neq 0$ everywhere on M . The contact form η induces a hyperplane subbundle H of the tangent bundle TM given by $H = Ker \eta$. The Reeb vector field associated to η is the vector field ξ uniquely determined by $\eta(\xi) = 1$ and $d\eta(\xi, \cdot) = 0$. Then (M, η) is called a contact manifold.

Note that given $H = Ker \eta$ and ξ such that $\eta(\xi) = 1$, we can split the tangent bundle into $TM = H \oplus \mathbb{R}\xi$. If X is any vector field on M , then X decomposes as $X = X_H + f\xi$ for any $f \in C^\infty(M, \mathbb{R})$. X_H is called the horizontal part of X .

If (M, η) is a contact manifold, then the pair $(H, d\eta|_H)$ is a symplectic vector bundle. We fix an almost complex structure J_H on H compatible with $d\eta|_H$, i.e. $d\eta|_H(J_H(X), J_H(Y)) = d\eta|_H(X, Y)$. We can extend J_H to an endomorphism J of the tangent bundle TM by setting $J\xi = 0$. The relation $J^2 = -Id + \eta \otimes \xi$ then holds. With this in mind, g_η , given by

$$g_\eta(X, Y) = d\eta(X, JY) + \eta(X)\eta(Y),$$

defines a Riemannian metric on TM . The metric g_η is called a Webster metric and is said to be associated to η . Moreover, the following relations hold:

$$g_\eta(\xi, X) = \eta(X), \quad g_\eta(JX, Y) = d\eta(X, Y), \quad g_\eta(JX, JY) = g_\eta(X, Y) - \eta(X)\eta(Y)$$

for any $X, Y \in \Gamma(TM)$. We call $(M, g_\eta, \eta, \xi, J)$ a contact metric manifold. For detailed information, see [2, 12].

The generalized Tanaka–Webster connection ∇ is a well-known connection on the contact metric manifold $(M, g_\eta, \eta, \xi, J)$. This connection satisfies the conditions $\nabla\eta = 0$ and $\nabla g_\eta = 0$. Moreover, if J is integrable, i.e. $\nabla J = 0$, then the contact metric manifold $(M, g_\eta, \eta, \xi, J)$ is called a strictly pseudoconvex CR manifold [12].

3. Self-dual 2-forms on 5-dimensional contact metric manifolds

Let $(M, g_\eta, \eta, \xi, J)$ be a 5-dimensional contact metric manifold. The p -form α is called a horizontal p -form if $i(\xi)\alpha = 0$ where i is contraction operator. For any 2-form $\alpha \in \Omega^2(M)$ we have the splitting $\alpha = \alpha_H + \alpha_\xi$ where $\alpha_H = \alpha \circ \Pi$, $\Pi : TM \rightarrow H$ is the canonical projection and $\alpha_\xi = \eta \wedge i(\xi)\alpha$. The decomposition of $\Omega^2(M)$ is then given by

$$\Omega^2(M) = \Omega_H^2(M) \oplus \eta \wedge \Omega_H^1(M), \tag{1}$$

where $\Omega_H^2(M)$ and $\Omega_H^1(M)$ are the bundles of horizontal forms. Moreover, any horizontal 2-form can be split into its self-dual and anti-self dual parts as follows.

Let \star be the Hodge-star operator acting on the cotangent bundle T^*M . We can define the operator

$$\star : \Omega^2(M) \rightarrow \Omega^2(M), \quad \star(\beta) := \star(\eta \wedge \beta).$$

We can restrict the operator \star to the space of horizontal 2-forms $\Omega_H^2(M)$:

$$\star_H : \Omega_H^2(M) \rightarrow \Omega_H^2(M), \quad \star_H(\beta) := \star(\eta \wedge \beta).$$

This operator satisfies $\star_H^2 = id$. Then we have the following orthogonal decomposition:

$$\Omega_H^2(M) = \Omega_H^2(M)^+ \oplus \Omega_H^2(M)^-, \tag{2}$$

where $\Omega_H^2(M)^\pm$ is the eigenspace associated to eigenvalue ± 1 of the operator \star_H . The eigenspace $\Omega_H^2(M)^+$ is called as the space of self-dual 2-forms. In a similar way, the eigenspace $\Omega_H^2(M)^-$ is called the space of anti-self-dual 2-forms (see [3, 8]). From equalities (1) and (2), we have

$$\Omega^2(M) = \Omega_H^2(M)^+ \oplus \Omega_H^2(M)^- \oplus \eta \wedge \Omega_H^1(M).$$

Hence, any 2-form α can be written as $\alpha = \alpha_H^+ + \alpha_H^- + \eta \wedge \beta$ where β is a 1-form on H . The self-dual part of α is defined as the self-dual part of α_H , i.e. $\alpha^+ := \alpha_H^+$.

Locally, we can specify the self-dual and anti-self-dual 2-forms. For this, choose a local orthonormal frame field $\{e_1, e_2 = J(e_1), e_3, e_4 = J(e_3), \xi\}$ and denote by $\{e^1, e^2, e^3, e^4, \eta\}$ the dual basis. From (2), the 2-form $d\eta$ has the form $d\eta = e^1 \wedge e^2 + e^3 \wedge e^4$. The forms $e^1 \wedge e^2 + e^3 \wedge e^4$, $e^1 \wedge e^3 - e^2 \wedge e^4$ and $e^1 \wedge e^4 + e^2 \wedge e^3$ are an orthonormal basis for $\Omega_H^2(M)^+$. An orthonormal basis for $\Omega_H^2(M)^-$ is given by the forms $e^1 \wedge e^2 - e^3 \wedge e^4$, $e^1 \wedge e^3 + e^2 \wedge e^4$, and $e^1 \wedge e^4 - e^2 \wedge e^3$.

4. Dirac operators on contact metric manifolds

In this section we will describe Dirac operators on contact metric manifolds. For this, we need a $Spin^c$ -structure. Any contact metric manifold admits a canonical $Spin^c$ -structure. Then we have a $Spin^c$ -bundle $P_{Spin^c(2n)}$, an S^1 -bundle P_{S^1} , and the canonical line bundle \mathcal{L} . The spinor bundle S can be identified with the bundle $\wedge_{\mathbb{H}}^{0,*}M$ of the $(0, *)$ forms. For the definitions and more details about these notions, we refer to [12]. For our purpose, we use the following representation of the complex Clifford algebra $\mathbb{C}l_5$:

$$\begin{aligned} \kappa(e_1) &= \begin{pmatrix} 0 & i & 0 & 0 \\ i & 0 & 0 & 0 \\ 0 & 0 & 0 & i \\ 0 & 0 & i & 0 \end{pmatrix}, \quad \kappa(e_2) = \begin{pmatrix} 0 & 1 & 0 & 0 \\ -1 & 0 & 0 & 0 \\ 0 & 0 & 0 & -1 \\ 0 & 0 & 1 & 0 \end{pmatrix}, \\ \kappa(e_3) &= \begin{pmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & -1 & 0 \\ 0 & 1 & 0 & 0 \\ -1 & 0 & 0 & 0 \end{pmatrix}, \quad \kappa(e_4) = \begin{pmatrix} 0 & 0 & 0 & i \\ 0 & 0 & -i & 0 \\ 0 & -i & 0 & 0 \\ i & 0 & 0 & 0 \end{pmatrix}, \\ \kappa(e_5) &= \begin{pmatrix} i & 0 & 0 & 0 \\ 0 & -i & 0 & 0 \\ 0 & 0 & i & 0 \\ 0 & 0 & 0 & -i \end{pmatrix}, \quad \kappa(d\eta) = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 2i & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -2i \end{pmatrix}. \end{aligned}$$

Let $(M, g_\eta, \eta, \xi, J)$ be a contact metric manifold equipped with a $Spin^c$ -structure. Each unitary connection A on \mathcal{L} induces a spinorial connection ∇^A on S with the generalized Tanaka–Webster connection ∇ . The Kohn–Dirac operator D_H^A is defined as follows:

$$D_H^A = \sum_{i=1}^{2n} \kappa(e_i)(\nabla_{e_i}^A),$$

where $\{e_i\}$ is a local orthonormal frame of H . The Dirac operator D_A is defined by

$$D_A = D_H^A + \xi \cdot \nabla_\xi^A$$

(see also [12]).

5. Seiberg–Witten-like equations on 5-dimensional contact metric manifolds

In [6], Seiberg–Witten-like equations on 5-dimensional Euclidean space \mathbb{R}^5 were written. In this section, we will write Seiberg–Witten-like equations on 5-dimensional contact metric manifolds and give a solution to these equations on strictly pseudoconvex CR manifolds.

For a spinor ψ we define a 2-form $\sigma(\psi)$ by the following formula:

$$\sigma(\psi)(X, Y) = \langle X \cdot Y \cdot \psi, \psi \rangle + g_\eta(X, Y)|\psi|^2,$$

where $X, Y \in \Gamma(TM)$ and \langle, \rangle is the Hermitian inner product on the spinor space S . Note that $\sigma(\psi)$ is an imaginary valued 2-form. The restriction of $\sigma(\psi)$ to H is denoted by $\sigma_H(\psi)$.

Definition 1 *Let $(M, g_\eta, \eta, \xi, J)$ be a contact metric 5-manifold. Fix a $Spin^c$ -structure and a connection A in the $U(1)$ -principal bundle associated with the $Spin^c$ -structure. For any $\psi \in \Gamma(S)$ Seiberg–Witten equations are defined by*

$$\begin{aligned} D_A(\psi) &= 0, \\ F_A^+ &= -\frac{1}{4}\sigma(\psi)^+, \end{aligned} \tag{3}$$

where F_A^+ is the self-dual part of the curvature F_A and $\sigma(\psi)^+$ is the self-dual part of the 2-form $\sigma(\psi)$.

Now we give a solution for Seiberg–Witten equations in dimension 5. To do this, we follow the method given in [9]. From now on we suppose that $(M, g_\eta, \eta, \xi, J)$ is a strictly pseudoconvex CR manifold.

Let (M, g_η) be a contact metric manifold endowed with $Spin^c$ -structure. The spinor bundle is then $S = \wedge_H^{0,*}(M)$. Namely,

$$S = \wedge_H^{0,2}(M) \oplus \wedge_H^{0,1}(M) \oplus \wedge_H^{0,0}(M),$$

where $\wedge_H^{0,2}(M)$ is the eigenspace corresponding to the eigenvalue $2i$ of the mapping $\kappa(d\eta) : S \rightarrow S$ and has dimension 1, $\wedge_H^{0,1}(M)$ is the eigenspace corresponding to the eigenvalue 0 of the mapping $\kappa(d\eta) : S \rightarrow S$ and has dimension 2, and $\wedge_H^{0,0}(M)$ is the eigenspace corresponding to the eigenvalue $-2i$ of the mapping $\kappa(d\eta) : S \rightarrow S$ and has dimension 1.

If $\psi_0 \in \wedge_H^{0,0}(M)$, then ψ_0 denotes the spinor corresponding to the constant function 1 in the chosen coordinates

$$\psi_0 = \begin{pmatrix} 0 \\ 0 \\ 0 \\ 1 \end{pmatrix}.$$

Moreover, we have $d\eta \cdot \psi_0 = -2i\psi_0$. By using the expression of $\sigma_H(\psi)$ in the local coordinates, we obtain the following identity:

$$\sigma_H(\psi_0) = -id\eta. \tag{4}$$

5.1. Some identities

In this part, we collect some identities needed for the special solution of Seiberg–Witten equations.

When M is a strictly pseudoconvex CR manifold, M also has a complex CR structure [2]. Let $\{Z_1, \dots, Z_n\}$ be a local unitary frame of $T^{1,0}$ over $U \subset M$ where $Z_\alpha = \frac{1}{\sqrt{2}}(e_\alpha - \sqrt{-1}Je_\alpha)$, $1 \leq \alpha \leq n$. Let us denote by $\omega := (\omega_{\alpha\beta})$ the matrix of the connection form of ∇ with respect to the frame. Then we can write the following:

$$\nabla Z_\alpha = \sum_{\beta} \omega_{\alpha\beta} Z_\beta.$$

$\{Z_1, \dots, Z_n, \bar{Z}_1, \dots, \bar{Z}_n, \xi\}$ is a local frame of the complexified tangent bundle $TM^{\mathbb{C}}$ over U .

Let $\{\theta^1, \dots, \theta^n, \bar{\theta}^1, \dots, \bar{\theta}^n, \eta\}$ be the corresponding dual basis. Thus,

$$\zeta = \bar{\theta}^1 \wedge \dots \wedge \bar{\theta}^n : U \rightarrow \Lambda_H^{0,n}(M)$$

is a local section in determinant line bundle $\Lambda_H^{0,n}(M)$. The Webster connection ∇ defines a covariant derivative in the canonical line bundle $\Lambda_H^{0,n}(M)$ such that

$$\nabla(\bar{\theta}^1 \wedge \dots \wedge \bar{\theta}^n) = -Tr(\omega)\bar{\theta}^1 \wedge \dots \wedge \bar{\theta}^n.$$

Since ∇ is a metric with respect to g_η , the trace $Tr(\omega)$ is purely imaginary. Therefore, this connection ∇ in $\Lambda_H^{0,n}(M)$ induces a connection on the associated S^1 -principal bundle P_{S^1} . Let us denote this connection by A . Then,

$$\zeta^* A = -Tr \bar{\omega} = Tr \omega$$

is a local connection form on S^1 -bundle P_{S^1} . Let F_A be the curvature form of the connection A . The curvature form F_A is a 2-form on M with values in $i\mathbb{R}$. Over $U \subset M$ we have

$$F_A = dA = Tr d\omega. \tag{5}$$

Moreover,

$$Ric(X, Y) = Tr(d\omega) - Tr(\omega \wedge \omega) = Tr d\omega. \tag{6}$$

From (5) and (6) it follows that

$$F_A = Ric. \tag{7}$$

Here we follow the similar procedures given in [2].

In the following, the Ricci form ρ_H is defined by

$$\rho_H(X, Y) = Ric(X, J_H Y) = g_\eta(X, J_H Ric Y)$$

for any $X, Y \in \Gamma(H)$. In the case of a strictly pseudoconvex CR manifold, the almost complex structure J_H is complex. Therefore, we have the equation

$$Ric(X, Y) = i\rho_H(X, Y) \tag{8}$$

for any $X, Y \in \Gamma(H)$.

Proposition 2 Let ρ_H be a Ricci form on H and s_H be a scalar curvature of the subbundle H . Then the following identity holds:

$$\rho_H^+ = -\frac{s_H}{4}d\eta, \tag{9}$$

where ρ_H^+ is a the self-dual part of the Ricci form ρ_H .

Proof In local coordinates the almost complex structure J is given as follows.

$$J = \begin{pmatrix} 0 & -1 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -1 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix}$$

Since $J \circ Ric = Ric \circ J$, we obtain the reduced form of the Ric as follows.

$$Ric = \begin{pmatrix} R_{11} & 0 & R_{13} & R_{14} & 0 \\ 0 & R_{11} & -R_{14} & R_{13} & 0 \\ R_{13} & -R_{14} & R_{33} & 0 & 0 \\ R_{14} & R_{13} & 0 & R_{33} & 0 \\ 0 & 0 & 0 & 0 & R_{55} \end{pmatrix}$$

The Ricci form ρ_H can be written in the following way:

$$\rho_H = -R_{11}e^1 \wedge e^2 - R_{33}e^3 \wedge e^4 - R_{13}(e^1 \wedge e^4 - e^2 \wedge e^3) + R_{14}(e^1 \wedge e^3 + e^2 \wedge e^4).$$

Since the 2-forms $e^1 \wedge e^4 - e^2 \wedge e^3$ and $e^1 \wedge e^3 + e^2 \wedge e^4$ are anti-self-dual 2-forms, the self-dual part of ρ_H is given by

$$\rho_H^+ = \frac{-R_{11} - R_{33}}{2}d\eta = -\frac{R_{11} + R_{22} + R_{33} + R_{44}}{4}d\eta = -\frac{s_H}{4}d\eta,$$

where s_H is the restricted scalar curvature to H . □

5.2. A special solution to 5-dimensional Seiberg–Witten equations

Let $(M, g_\eta, \eta, \xi, J)$ be a strictly pseudoconvex contact manifold of dimension 5. Suppose that the scalar curvature s_H of the subbundle H is negative and constant. Then let $\psi = \sqrt{-s_H}\psi_0$. In this case, $\psi \in \wedge^0_{H^0}(M)$. From (4) we have

$$\sigma_H(\psi) = is_Hd\eta. \tag{10}$$

By using (7),(8), (9), and (10) we obtain

$$F_A^+ = Ric^+ = i\rho_H^+ = -i\frac{s_H}{4}d\eta = -\frac{1}{4}\sigma_H(\psi). \tag{11}$$

Note that since $d\eta$ is a self-dual 2-form, $\sigma_H(\psi)$ is also i.e., $\sigma_H(\psi)^+ = \sigma_H(\psi)$. Because of $\sigma(\psi)^+ = \sigma_H(\psi)^+$ and with identity (11), we get

$$F_A^+ = -\frac{1}{4}\sigma(\psi)^+.$$

One can show that $\nabla_{e_i}^A \psi_0 = 0$. Therefore, we deduce that

$$D_H^A \psi = 0.$$

Moreover,

$$D_A \psi = 0.$$

The pair $(A, \psi = \sqrt{-s_H} \psi_0)$ is a solution of Seiberg–Witten-like equations in (3).

References

- [1] Akbulut S. Lectures on Seiberg–Witten Invariants. Turk J Math 1996; 20: 95–118.
- [2] Baum H. Lorentzian twistor spinors and CR-geometry. Differ Geom Appl 1996; 11: 69–96.
- [3] Bellettini C. Almost complex structures and calibrated integral cycles in contact 5-manifolds. Adv Calc Var 2013; 6: 339–374.
- [4] Bilge AH, Dereli T, Koçak Ş. Monopole equations on 8-manifolds with Spin(7) holonomy. Commun Math Phys 1999; 203: 21–30.
- [5] Corrigan E, Devchand C, Fairlie DB, Nuyts J. First-order equations for gauge fields in spaces of dimension greater than four. Nucl Phys B 1983; 214: 452–464.
- [6] Degirmenci N, Karapazar Ş. Seiberg–Witten like monopole equations on \mathbb{R}^5 . J Partial Dif 2011; 24: 150–157.
- [7] Degirmenci N, Özdemir N. Seiberg–Witten like equations on 7-manifolds with G2-structures. J Nonlinear Math Phys 2005; 12: 457–461.
- [8] Fan H. Half De Rham complexes and line fields on odd-dimensional manifolds. Trans Am Math Society 1996; 348: 2947–2982.
- [9] Friedrich T. Dirac Operators in Riemannian Geometry. Providence, RI, USA: AMS, 2000.
- [10] Gao YH, Tian G. Instantons and the monopole-like equations in eight dimensions. J High Energy Phys 2000; 5: 036.
- [11] Morgan J. Seiberg–Witten Equations and Applications to the Topology of Smooth Manifolds. Princeton, NJ, USA: Princeton University Press, 1996.
- [12] Petit R. Spin^c-structures and Dirac operators on contact manifolds. Differ Geom Appl 2005; 22: 229–252.
- [13] Tian G. Gauge theory and calibrated geometry I. Ann Math 2000; 151: 193–268.
- [14] Witten E. Monopoles and four manifolds. Math Res Lett 1994; 1: 769–796.