

1-1-2004

Seasonal Distribution of Phytoplankton in Orduzu Dam Lake (Malatya, Turkey)

AHMET KADRİ ÇETİN

BÜLENT ŞEN

Follow this and additional works at: <https://journals.tubitak.gov.tr/botany>



Part of the [Botany Commons](#)

Recommended Citation

ÇETİN, AHMET KADRİ and ŞEN, BÜLENT (2004) "Seasonal Distribution of Phytoplankton in Orduzu Dam Lake (Malatya, Turkey)," *Turkish Journal of Botany*. Vol. 28: No. 3, Article 2. Available at: <https://journals.tubitak.gov.tr/botany/vol28/iss3/2>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Botany by an authorized editor of TÜBİTAK Academic Journals. For more information, please contact academic.publications@tubitak.gov.tr.

Seasonal Distribution of Phytoplankton in Orduzu Dam Lake (Malatya, Turkey)

A. Kadri ÇETİN

Firat University, Science and Art Faculty, Department of Biology, 23119, Elazığ - TURKEY

Bülent ŞEN

Firat University, Aquaculture Faculty, 23119, Elazığ - TURKEY

Received: 15.11.2002

Accepted: 17.10.2003

Abstract: The species composition and seasonal distribution of phytoplankton in Orduzu Dam Lake was studied for a year. Diatoms (*Bacillariophyta*) were most diverse, followed by green algae (*Chlorophyta*), blue-green algae (*Cyanophyta*), euglenoids (*Euglenophyta*) and dinoflagellates (*Dinophyta*). A total of 117 taxa were recorded and the phytoplankton of the lake contained a large number of detached benthic algae. Phytoplankton assemblages were dominated by diatoms in all the periods investigated and centric diatoms were the most abundant. Overall phytoplankton density was high during the spring and summer months and the highest phytoplankton density was observed in August. The abundance of phytoplankton was positively correlated with water temperature.

Key Words: Orduzu Dam Lake, Phytoplankton, Seasonal variations, Malatya, Turkey.

Orduzu Baraj Gölü (Malatya, Türkiye) Fitoplanktonunun Mevsimsel Değişimi

Özet: Bu çalışmada Orduzu Baraj Gölü fitoplanktonunun tür kompozisyonu ve mevsimsel değişimi bir yıl süreyle incelenmiştir. Fitoplankton içerisinde diyatome tür çeşitliliği bakımından en zengin grubu oluştururken onları *Chlorophyta*, *Cyanophyta*, *Euglenophyta* ve *Dinophyta* üyeleri izlemiştir. Fitoplanktonda toplam 117 taxa kaydedilmiştir. Diyatome araştırma süresince, fitoplanktonda baskın alg grubunu oluşturmuşlardır. Fitoplankton yoğunluğu ilkbahar ve yaz aylarında yüksek olurken maksimum fitoplankton yoğunluğu Ağustos ayında gözlenmiştir. Fitoplankton yoğunluğu su sıcaklığı ile pozitif bir korelasyon göstermiştir.

Anahtar Sözcükler: Orduzu Baraj Gölü, Fitoplankton, Mevsimsel değişim, Malatya, Türkiye

Introduction

Algae are the major primary producers in many aquatic systems and are an important food source for other organisms. They include planktonic and benthic forms. Species composition and the seasonal variations of planktonic and benthic forms in freshwaters are dependent on interactions between physical and chemical factors. A considerable amount of information has been gathered over the last 10 years on the ecology and distribution of algae in lakes and running waters in eastern Anatolia. The majority of these studies are concerned with seasonal variations of phytoplankton in natural lakes (Şen, 1988; Çetin, 2000) and reservoirs (Şen & Çetin, 1988; Çetin & Şen, 1998; Çetin & Yıldırım, 2000). Algae in running water have been studied less (Şen et al., 1999; Yavuz & Çetin, 2000).

There are no algological studies on Orduzu Dam Lake in the literature. The purpose of this study was to determine the species composition and seasonal variations of phytoplankton in Orduzu Dam Lake.

Study area

Orduzu Dam Lake was constructed for irrigation purposes in 1979 and is situated in the east of Turkey, 5 km from the city of Malatya (38°20' E, 38°25' N). The surface area, water capacity, average depth and altitude of the reservoir are 15 km², 1,600,000 m³, 6 m and 950 m respectively. The lake has a narrow, elongated shape (Figure 1). It is shallow, with a maximum depth of 9 m, and it does not stratify. The climate of region is arid; the winters are cold and the summers hot and dry.



Figure 1. Map of Orduzu Dam Lake showing the position of sampling stations (o).

The location of Orduzu Dam Lake and the sampling stations are shown in Figure 1. Three sampling stations were chosen in the lake. The first station was on the north side of the lake (average depth of 3 m). The second sampling station was in the middle of the lake (average depth of 4 m), and the third was on the south side (average depth of 3 m).

Materials and Methods

Phytoplankton and water samples were taken from a 0-0.5 m water column monthly from the stations between April 1997 and May 1998 using a 1.5 l Nansen water sampler. Dissolved oxygen concentration and temperature were measured with a combined electrode (HACH oxygenmeter). Conductivity was measured through a conductivimeter (HACH 17250 model). pH was measured with a CyberScan (PD300 model) pH meter and transparency with a Secchi disc in situ. Total hardness, calcium, magnesium (by titration methods), nitrate-nitrogen (by brucinsulfate method), silica (by molybdosilicate method) and sulphate (by turbidimetric

method) concentrations of the lake water were determined in the laboratory (APHA, 1985).

Phytoplankton individuals were counted using an inverted microscope (Lund et al., 1958). The phytoplankton density was estimated by counting all individuals. Single cells, colonies and filaments were all considered as individuals and the results are expressed as individuals ml⁻¹ (ind. ml⁻¹). The phytoplankton was identified mainly using the works of Geitler (1925), Germain (1981), Patrick & Reimer (1966, 1975), Huber-Pestalozzi (1968), Prescott (1982), Ettl (1983) and Krammer & Lange-Bertalot (1986).

Results

Physical and chemical variables

The highest and lowest water temperatures in Orduzu Dam Lake were 23.5 °C and 4.8 °C in August and February, respectively. The annual cycle of water temperature at all stations showed a clear maximum in summer and a minimum in winter. Dissolved oxygen was 8.9-10.0 mg l⁻¹. Dissolved oxygen concentrations

decreased in summer and increased in winter. The maximum conductivity (375 μs) was measured in July and minimum conductivity (200 μs) occurred in November (Table 1).

The lowest and highest light visibilities were 0.36 m and 1.46 m in April and August respectively. The lake was slightly basic and pH values varied between 7.5 and 8.3. The total hardness showed an irregular pattern, fluctuating between 143 and 180 mg l^{-1} CaCO_3 . Seasonal fluctuations of nitrate-nitrogen were negligible in the lake. The maximum concentration of nitrate-nitrogen (0.023 mg l^{-1}) was determined in February and the minimum concentration (0.008 mg l^{-1}) was observed in July. Concentrations of sulphate were almost stable throughout the sampling period. The maximum (5.06 mg l^{-1}) and minimum (2.16 mg l^{-1}) values of sulphate were measured in August and September, respectively. Silica showed seasonal variations in Orduzu Dam Lake. The maximum concentration of silica (16.50 mg l^{-1}) was observed in February whilst the minimum value (5.65 mg l^{-1}) was recorded in October (Table 1).

Seasonal variations of phytoplankton

The total of 117 algal taxa belonging to *Bacillariophyta*, *Chlorophyta*, *Cyanophyta*, *Euglenophyta* and *Dinophyta* were identified in the phytoplankton. They are listed in Table 2. Diatoms were the most significant algae with respect to number of species and abundance in

the phytoplankton, and centric diatoms were more conspicuous compared to pennate forms. *Cyclotella bodanica* Grun., *C. comta* (Ehrenb.) Kütz., *C. krammeri* Håk., *C. stelligera* Cleve & Grun., *C. ocellata* Pant., *Cymbella amphicephala* Naegeli ex Kütz., *Navicula veneta* Kütz. and *Fragilaria ulna* (Nitzsch) Lange-Bert. were the most dominant algae at all stations in the phytoplankton.

In general, the dynamics of algal numbers of all algal groups in the phytoplankton were quite similar (Figure 2). All algal groups started to increase in numbers in April and continued to increase slightly until the end of the summer. During this growth period the highest numbers of *Chlorophyta* were observed in May, whilst that of diatoms occurred in August. The numbers of individual of diatoms started to decrease after reaching their maximum and decreased regularly and continuously until the end of winter. By contrast, the number of individuals of other algal groups was either decreasing or low during autumn and winter. However, an exception was recorded in October when individual numbers of all algae excluding diatoms increased suddenly. *Dinophyta* and *Euglenophyta* were present from April to November but never exceeded 10 ind. ml^{-1} during the study. These algae were absent in winter.

In April, the phytoplankton was 358 ind. ml^{-1} and was dominated by diatoms (*Cyclotella bodanica*, *C. comta*, *C. krammeri*, *C. stelligera*, *C. ocellata*, *Achnanthes flexella* (Kütz.) Brunth., *Cymbella affinis* Kütz., *Navicula veneta* and *Fragilaria ulna*).

Table 1. Variations in concentrations of some physical and chemical parameters in Orduzu Dam Lake water.

Sampling dates	Water temp., °C	pH	Dissolved O ₂ , (mg l ⁻¹)	Transparency (cm)	Total hardness CaCO ₃ (mg l ⁻¹)	Ca ²⁺ (mg l ⁻¹)	Mg ²⁺ (mg l ⁻¹)	Conductivity (μs)	Silica (mg l ⁻¹)	SO ₄ ²⁻ (mg l ⁻¹)	NO ₃ -N (mg l ⁻¹)
Apr. 1997	10.90 ± 0.36	8.23 ± 0.15	9.71 ± 0.02	39.00 ± 2.64	161.00 ± 1.00	44.53 ± 0.61	12.06 ± 0.56	281.66 ± 2.88	11.93 ± 0.11	3.62 ± 0.60	0.050 ± 0.003
May 1997	14.93 ± 0.11	8.10 ± 0.10	9.41 ± 0.18	60.00 ± 2.64	154.00 ± 2.30	40.80 ± 0.80	12.79 ± 1.01	293.33 ± 2.88	15.50 ± 0.80	3.98 ± 0.15	0.017 ± 0.002
Jun. 1997	18.50 ± 0.50	7.96 ± 0.05	9.11 ± 0.12	101.00 ± 1.00	144.00 ± 1.00	39.46 ± 0.46	11.01 ± 0.50	276.66 ± 5.77	9.33 ± 0.35	3.40 ± 0.62	0.015 ± 0.005
Jul. 1997	20.06 ± 0.11	8.16 ± 0.05	9.06 ± 0.05	116.33 ± 4.04	179.00 ± 1.00	48.00 ± 0.40	14.33 ± 0.48	205.00 ± 5.00	7.10 ± 0.26	4.63 ± 0.57	0.017 ± 0.002
Aug. 1997	23.26 ± 0.25	7.96 ± 0.05	9.20 ± 0.00	144.33 ± 2.08	170.66 ± 1.15	46.66 ± 0.61	13.12 ± 0.24	321.66 ± 7.63	7.31 ± 0.40	5.06 ± 0.10	0.015 ± 0.005
Sep. 1997	23.00 ± 0.00	7.93 ± 0.11	9.01 ± 0.12	98.33 ± 0.57	147.33 ± 1.15	38.26 ± 1.00	12.63 ± 0.87	296.66 ± 2.88	6.01 ± 0.75	2.16 ± 0.07	0.015 ± 0.005
Oct. 1997	19.86 ± 0.23	8.03 ± 0.23	9.13 ± 0.05	75.33 ± 1.52	157.33 ± 1.15	41.33 ± 0.46	13.12 ± 0.00	360.00 ± 0.00	5.65 ± 0.21	3.03 ± 0.23	0.017 ± 0.002
Nov. 1997	11.90 ± 0.36	8.03 ± 0.15	9.30 ± 0.05	78.00 ± 1.00	161.33 ± 5.50	41.33 ± 1.40	13.87 ± 1.40	366.66 ± 7.63	8.18 ± 0.02	2.26 ± 0.12	0.015 ± 0.005
Dec. 1997	8.10 ± 0.17	7.83 ± 0.25	9.45 ± 0.13	80.33 ± 0.57	159.33 ± 1.15	40.66 ± 0.46	14.02 ± 0.30	366.66 ± 7.63	8.28 ± 0.02	2.23 ± 0.07	0.015 ± 0.000
Jan. 1998	5.06 ± 0.11	7.66 ± 0.20	9.90 ± 0.10	100.00 ± 1.00	154.33 ± 1.15	39.60 ± 0.69	13.44 ± 0.60	318.33 ± 2.88	14.33 ± 1.13	3.10 ± 0.25	0.017 ± 0.002
Feb. 1998	4.93 ± 0.11	8.13 ± 0.20	10.03 ± 0.05	75.66 ± 0.57	148.66 ± 1.52	38.13 ± 1.00	12.96 ± 0.27	303.33 ± 5.77	16.50 ± 0.50	2.15 ± 0.10	0.067 ± 0.007
Mar. 1998	6.96 ± 0.05	8.06 ± 0.11	9.96 ± 0.05	58.66 ± 1.52	163.66 ± 1.15	43.60 ± 1.44	13.28 ± 0.85	321.66 ± 2.88	14.25 ± 0.25	2.53 ± 0.02	0.013 ± 0.005

Table 2. Algal taxa recorded in the phytoplankton of Orduzu Dam Lake.

<i>Bacillariophyta</i>	<i>Gomphonema affine</i> Kütz.
<i>Cyclotella bodanica</i> Grun.	<i>G. dichotomum</i> S.Wunsam
<i>C. comta</i> (Ehrenb.) Kütz.	<i>G. subtile</i> Ehrenb.
<i>C. krammeri</i> Håk.	<i>Hannaea arcus</i> (Ehrenb.) in Patr. & Reimer
<i>C. stelligera</i> Cleve & Grun.	<i>Hantzschia amphioxys</i> (Ehrenb.) Grun.
<i>C. ocellata</i> Pant.	<i>H. amphioxys</i> var. <i>maior</i> Grun.
<i>C. planctonica</i> Brunnth.	<i>Navicula bacillum</i> Ehrenb.
<i>Achnanthes delicatula</i> Kütz.	<i>N. cari</i> Ehrenb.
<i>A. flexella</i> (Kütz.) Brunnth.	<i>N. cincta</i> (Ehrenb.) Ralfs in A.Pritch.
<i>A. gibberula</i> Grun in Cleve & Grun.	<i>N. cocconeiformis</i> Greg. ex Greville
<i>A. minutissima</i> Kütz.	<i>N. veneta</i> Kütz.
<i>Amphora ovalis</i> (Kütz.) Kütz.	<i>N. cuspidata</i> (Kütz.) Kütz.
<i>A. veneta</i> Kütz.	<i>N. cuspidata</i> var. <i>heribaudii</i> M.Peragallo in Héríb.
<i>Asterionella formosa</i> Hassall	<i>N. tripunctata</i> (O.F.Müll.) Bory
<i>Caloneis alpestris</i> (Grun.) Cleve	<i>N. neoventricosa</i> Hust.
<i>C. ventricosa</i> (Ehrenb.) F.Meister	<i>N. pupula</i> Kütz.
<i>Cocconeis placentula</i> Ehrenb.	<i>N. radiosa</i> Kütz.
<i>Cymatopleura elliptica</i> (Breb. ex Kütz.) W.Sm.	<i>N. reinhardtii</i> Grun. in Van Heurck
<i>C. librile</i> (Ehrenb.) Pant.	<i>N. rhyncocephala</i> Kütz.
<i>Cymbella affinis</i> Kütz.	<i>N. salinarum</i> Grun. in Cleve & Grun.
<i>C. amphicephala</i> Naegeli ex Kütz.	<i>N. veneta</i> Kütz.
<i>C. aspera</i> (Ehrenb.) H.Perag	<i>Neidium affine</i> (Ehrenb.) Pfitzer
<i>C. cistula</i> (Ehrenb.) Kirchner	<i>N. binodis</i> (Ehrenb.) Hust.
<i>C. cistula</i> var. <i>maculata</i> (Kütz.) Van Heurck	<i>N. dubium</i> (Ehrenb.) Cleve
<i>C. cuspidata</i> Kütz.	<i>N. iridis</i> (Ehrenb.) Cleve
<i>C. helvetica</i> Kütz.	<i>Nitzschia acicularis</i> (Kütz.)W.Sm.
<i>C. obtusiuscula</i> Kütz.	<i>N. amphibia</i> Grun.
<i>C. parva</i> (W.Sm.) Kirchner	<i>N. apiculata</i> (Gregory) Grun.
<i>C. tumida</i> (Breb. ex Kütz.) Grun in Van Heurck	<i>N. gracilis</i> Hantzsch
<i>Denticula elegans</i> Kütz.	<i>N. linearis</i> W.Sm.
<i>D. tenue</i> C. Agardh	<i>N. palea</i> (Kütz.) W.Sm.
<i>Diatoma hyemale</i> (Roth) Heib.	<i>N. sigmoidea</i> (Nitzsch) W.Sm.
<i>D. vulgare</i> Bory	<i>N. sinuata</i> (Thwaites ex W.Sm.) Grun. in Cleve & Grun.
<i>Diploneis ovalis</i> (Hilse) Cleve	<i>N. tryblionella</i> Hantzsch in Rabenh.
<i>D. oblongella</i> (Naegeli ex Kütz.) R. Ross	<i>Pinnularia acoricola</i> Hust.
<i>Epithemia adnata</i> (Kütz.) Rabenh.	<i>P. brebissonii</i> (Kütz.) Rabenh.
<i>E. turgida</i> (Ehrenb.) Kütz.	<i>P. mesolepta</i> (Ehrenb.)W.Sm.
<i>Eunotia arcus</i> Ehrenb.	<i>P. viridis</i> (Nitzsch) Ehrenb.
<i>Fallacia pygmaea</i> (Kütz.) Stickle & D.G.Mann	<i>Rhoichosphenia abbreviata</i> (C.Agardh) Lange-Bertalot
<i>Fragilaria construens</i> (Ehrenb.) Grun.	<i>Stauroneis anceps</i> Ehrenb.
<i>F. construens</i> var. <i>binodis</i> (Ehrenb.) Grun.	<i>S. phoenicenteron</i> (Nitzsch) Ehrenb.
<i>F. ulna</i> (Nitzsch) Lange-Bert.	<i>Surirella linearis</i> W.Sm.
<i>Frustulia vulgaris</i> (Thwaites) De Toni	<i>S. robusta</i> var. <i>splendida</i> (Ehrenb.)Van Heurck

Table 2. continued

<i>Tabularia tabulata</i> (C.Agardh) D.M.Williams & Round	<i>Tetraëdron minimum</i> (A.Braun) Hansg.
<i>Chlorophyta</i>	<i>Cyanophyta</i>
<i>Eudorina elegans</i> Ehrenb.	<i>Chroococcus turgidus</i> (Kütz.) Naegeli
<i>Pandorina morum</i> (Mull.) Bory	<i>Merismopedia elegans</i> A.Braun in Kütz.
<i>Pediastrum boryanum</i> (Turp.) Menegh.	<i>M. punctata</i> Meyen
<i>P. duplex</i> Meyen	<i>M. tenuissima</i> Lemmerm.
<i>P. duplex</i> var. <i>clathratum</i> (A.Braun) Lagerh.	<i>O. limosa</i> (Roth) C.A.Agardh ex Gomont
<i>P. simplex</i> Meyen	<i>O. princeps</i> Vaucher ex Gomont
<i>Scenedesmus acuminatus</i> (Lagerh.) Chodat	<i>O. rubescens</i> (de Candolle) ex Gomont
<i>S. acutus</i> Meyen ex Ralfs	<i>O. tenuis</i> C.A. Agardh ex Gomont
<i>S. armatus</i> Chodat	<i>O. tenuis</i> var. <i>natans</i> Gomont
<i>S. intermedius</i> Chodat	<i>Phormidium formosum</i> (Gomont) Anagn. et Komárek
<i>S. longus</i> Meyen	<i>Dinophyta</i>
<i>S. obliquus</i> (Turp.) Kütz.	<i>Ceratium hirundinella</i> (O.F.Müll.) Dujard.
<i>S. protuberans</i> F.E.Fritsch in M.F.Rich	<i>Peridinium cinctum</i> (O.F.Müll.) Ehrenb.
<i>S. quadricauda</i> (Turp.) Chodat	<i>Euglenophyta</i>
<i>S. quadricauda</i> var. <i>longispina</i> (Chod.) G.M.Smith	<i>Euglena viridis</i> Ehrenb.
<i>S. quadricauda</i> var. <i>maximus</i> West & West	<i>Phacus acuminatus</i> A.Stokes




Figure 2. Seasonal variations in individual numbers of phytoplankton groups in Orduzu Dam Lake

Numbers of individuals in all algal groups started to increase in May and reached 519 ind. ml⁻¹. The dominant species were *Cyclotella comta*, *C. krammeri*, *C. stelligera*, *C. ocellata*, *Asterionella formosa* Hassal, *Cymbella affinis*, *Navicula veneta*, *N. tripunctata* (O.F.Müll.) Bory, *N. rhyncocephala* Kütz., *Pediastrum duplex* Meyen, *Tetraëdron minimum* (A.Braun) Hansg. and *Scenedesmus quadricauda* (Turp.) Chodat.

The highest summer number of individuals occurred in August (691 ind. ml⁻¹) and the proportions of the algal groups in the phytoplankton were as follows: *Bacillariophyta* (93.77%), *Chlorophyta* (3.03%), *Cyanophyta* (1.45%), *Dinophyta* (1.15%) and *Euglenophyta* (0.6%) (Figure 2). The phytoplankton was dominated by *Cyclotella comta*, *C. krammeri*, *C. ocellata*, *Achnanthes flexella*, *A. minutissima* Kütz., *Cymbella affinis*, *C. amphicephala*, *C. obtusiuscula* Kütz., *Navicula veneta*, *N. tripunctata*, *N. rhyncocephala*, *Nitzschia gracilis* Hantzsch, *Tetraëdron minimum* and *Ceratium hirundinella* (O.F.Müll.) Dujard. The numbers of individuals of centric diatoms were far higher than those of other algae. Diatoms were at their maximum in this month.

In September, the phytoplankton was dominated by *Bacillariophyta*, and centric diatoms were the most abundant (56%) once more. The phytoplankton composition in October was made up of *Bacillariophyta* (83.6%), *Chlorophyta* (7.49%), *Cyanophyta* (5.16%), *Dinophyta* (0.77%) and *Euglenophyta* (0.28%). The phytoplankton numbers in February were lowest (52 ind. ml⁻¹) at all sampling stations except for *Chlorophyta*. The phytoplankton increased (163 ind. ml⁻¹) in March again and *Bacillariophyta* (89.5%), *Chlorophyta* (7.36%),

Cyanophyta (1.84%) and *Euglenophyta* (1.3%) were all present.

Discussion

The phytoplankton of Lake Orduzu was dominated by diatoms, whilst green algae, euglenoids, blue-green algae and dinoflagellates were less significant. The species composition of the phytoplankton of Lake Orduzu showed similarities to those of many lakes and reservoirs in Turkey (e.g., Aykulu et al., 1983; Şen, 1988; Çetin & Şen, 1998; Çetin, 2000; Çetin & Yıldırım, 2000; Yıldız, 1985).

Although pennate diatoms were represented by a higher number of taxa, they were much lower in numbers of individuals than were the centric forms in this lake. Although centric diatoms were poor in number of species they were much more abundant as individuals than were pennate forms and other algae. The dominance of centric diatoms, represented mainly by *Cyclotella krammeri*, *C. stelligera*, *C. ocellata* and *C. comta* in this study, has been previously documented by several authors in Turkish lakes (Çetin, 2000; Çetin & Yıldırım, 2000; Gönüloğlu, 1985) and elsewhere (Round 1984; M'harzi et al., 1998; Piirsoo, 2001),

Centric diatoms are one of the best adapted algal groups to turbulent and turbid systems (Izaguirre et al., 2001), whereas pennate diatoms are regarded as benthic forms. However, plankton samples from shallow and turbulent water often contain benthic diatoms (mostly pennate forms), which are whirled up into the water as a result of water movement. It has been reported that pennate diatoms were richer in number of taxa than centric forms in the phytoplankton of many shallow Turkish lakes (Aykulu et al., 1983; Altuner, 1984; Gönüloğlu, 1985).

The seasonal succession and species composition of the phytoplankton were similar at all the sampling stations during the study. The reasons for the similarities might be due to similar environmental conditions at the stations since physical and chemical properties at the different stations were quite similar.

The seasonal variations of phytoplankton are related to a variety of environmental factors in aquatic environments (Wu & Chou, 1998). Water temperature and transparency are among the most important physical

factors affecting the distribution and seasonal variations of phytoplankton in lakes (Simon & Hildrev 1998; Mosisch et al., 1999). The effects of water temperature on phytoplankton have been examined in many freshwater ecosystems, and it was found that water temperature strongly regulates the seasonal variations of phytoplankton (Lund 1965; Richardson et al., 2000; Izaguirre et al., 2001). The increase in phytoplankton during the spring and summer months in Dam Lake Orduzu could also be a result of the increasing water temperature.

Light is a major resource for phytoplankton and has a complex pattern of spatial and temporal variability (Litchman, 2000). Suspended matter in lake water increases in autumn and spring, resulting in minimum transparency. During the summer the transparency was at its maximum level. There was also a significant correlation between the growth of phytoplankton and transparency in Orduzu Dam Lake since the largest populations of all algae occurred during the summer whilst individual numbers were low in winter.

No relation was observed between diatom growth and pH level since high and low individual numbers were observed at similar pH levels.

A notable relation was observed between diatom growth and silica concentrations in the present study. Silica concentrations decreased gradually during the vernal growth of diatoms and were recorded at their lowest when the vernal maximum ended. In fact, silica concentrations decreased during such growth periods and increased whilst individual numbers of diatoms were decreasing. This finding is in harmony with that of Pearsall (1930), who suggested that the fall in silica concentrations coincides with the diatom maxima.

The lake appeared to have moderately hard water. Calcium had no strong effects on the growth of any algal group since concentrations of calcium varied only between 38 and 48 mg l⁻¹ during the study. A similar situation was also observed for magnesium since variations in its concentrations were negligible.

There appeared to be no correlation between the growth of algae and nitrate and sulphate since concentrations of these ions changed only slightly throughout the study.

References

- Altuner Z (1984). Tortum Gölünde Bir İstasyondan Alınan Fitoplanktonun Kalitatif ve Kantitatif Olarak İncelenmesi. *Doğa Bilim Dergisi* A₂ 8(2): 161-182.
- APHA (1985). *Standard Methods for Examination of Water and Wastewater*. Washington: American Public Health Association.
- Aykulu G, Obalı O & Gönülol A (1983). Ankara Çevresindeki Bazı Göllerde Fitoplanktonun Yayılışı. *Doğa Bilim Dergisi: Temel Bilimler* 7: 227-288.
- Çetin AK & Şen B (1998). Diatoms (*Bacillariophyta*) in the Phytoplankton of Keban Reservoir and Their Seasonal Variations. *Turk J Bot* 22: 25-33.
- Çetin AK (2000). Phytoplankton of Gölbaşı Lake (Adıyaman, Turkey) and Their Seasonal Variations. *International Journal on Algae* 2(4): 87-96.
- Çetin AK & Yıldırım V (2000). Species Composition and Seasonal Variations of the Phytoplankton in Sürgü Reservoir (Malatya, Turkey). *Acta Hydrobiologica* 42: 21-28.
- Ettl H (1983). *Chlorophyta I, Protomonadina. Süßwasserflora von Mitteleuropa*. Stuttgart: Gustav Fisher-Verlag.
- Geitler L (1925). *Cyanophyceae, Die Süßwasser Flora Mitteleuropas*. Stuttgart: Gustav Fisher Pub.
- Germain H (1981). *Flore Des Diatomées: Diatomophycées*. Paris: Societe Nouvella Des Éditions Boubée.
- Gönülol A (1985). Studies on the Phytoplankton of the Bayındır Dam Lake. *Communications Serie C*, 3: 21-38.
- Huber-Pestalozzi G (1968). *Das Phytoplankton des Süßwassers. Systematic and Biologie*. Teil. III. Stuttgart: Schweizer-Verlag.
- Izaguirre I, O'Farrell I & Tell G (2001). Variation in phytoplankton composition and limnological features in a water-water ecotone of Lower Parana Basin (Argentina). *Freshwater Biology* 46: 63-74.
- Krammer K & Lange-Bertalot H (1991a). *Süßwasser-flora von Mitteleuropas, Bacillariophyceae, Band2/3, 3. Teil: Centrales, Fragilariaceae, Eunotaceae*. Stuttgart: Gustav Fischer Verlag.
- Litchman E (2000). Growth Rates of Phytoplankton under Fluctuating Light. *Freshwater Biology* 44: 223-235.
- Lund JWG, Kipling, C., & Le Cren ED (1958). The Inverted Microscope Method of Estimating Algal Numbers and Statistical Basis of Estimations by Counting. *Hydrobiologia* 11: 143-170.
- Lund JWG (1965). The Ecology of the Freshwater Phytoplankton. *Biological Reviews* 40: 231-293.
- M'harzi A, Tackx M, Daro HM, Kesaulia I, Caturao R & Podoor R (1998). Winter Distribution of Phytoplankton and Zooplankton around Some Sandbanks of the Belgian Coastal Zone. *Journal of Plankton Research* 20: 2031-2052.
- Mosisch TD, Bunn SE, Davies PM & Marshall CJ (1999). Effects of Shade and Nutrient Manipulation on Periphyton Growth in a Subtropical Stream. *Aquatic Botany* 64: 167-177.
- Patrick R & Reimer CW (1966). *The Diatoms of the United States. Vol. I: Fragilariaceae, Eunotiaceae, Achnantheaceae, Naviculaceae*. Philadelphia: The Academy of National Science of Philadelphia.
- Patrick R & Reimer CW (1975). *The Diatoms of the United States. Vol. II: Entomoneidaceae, Cymbellaceae, Gomphonemaceae, Epithemiaceae*. Philadelphia: The Academy of National Science of Philadelphia.
- Pearsall WH (1930). Phytoplankton in the English Lakes. 1. The proportions in the water of some dissolved substances of biological importance. *Journal of Ecology* 18: 306-320.
- Piirsoo K (2001). Phytoplankton of Estonian rivers in midsummer. *Hydrobiologia* 444: 135-146.
- Prescott GW (1982). *Algae of the Western Great Lakes Area*. Germany, Koenigstein: Otto Koeltz Science Pub.
- Richardson TL, Gibson CE & Heaney SI (2000). Temperature, Growth and Seasonal Succession of Phytoplankton in Lake Baikal, Siberia. *Freshwater Biology* 44: 43-440.
- Round FE (1984). *The Ecology of Algae*. Cambridge: Cambridge University Press.
- Şen B & Çetin AK (1988). Seasonal Dynamics of Benthic Diatoms in a Reservoir in South-East Turkey. *Proceedings of the 10th International Diatom Symposium Joensuu Finland*, 505-511.
- Şen B (1988). Hazar Gölü (Elazığ) Alg Florası ve Mevsimsel Değişimleri Üzerine Gözlemler, Kısım I. Litoral Bölge. *IX. Ulusal Biyoloji Kongresi Sivas* 3: 289-298.
- Şen B, Alp MT, Özrenk F, Ercan T & Yıldırım V (1999). A Study on the Amounts of Plant Nutrients and Organic Matter Carried into Lake Hazar (Elazığ-Turkey). *Fresenius Environmental Bulletin* 8: 272-279.
- Simon SCH & Hildrev AG (1998). Patterns in the Epilithic Community of a Lake Littoral. *Freshwater Biology* 39: 477-492.
- Wu JT & Chou JW (1998). Dinoflagellate Associations in Feitsui Reservoir, Taiwan. *Botanical Bulletin of Academia Sinica* 39: 137-145.
- Yavuz OG & Çetin AK (2000). Cıp Çayı (Elazığ, Türkiye) Pelajik Bölge Algleri ve Mevsimsel Değişimleri. *Fırat Üniv. Fen ve Müh.Bil. Dergisi* 12(2): 9-14.
- Yıldız K (1985). Altınapa Baraj Gölü Alg Toplulukları üzerinde Araştırmalar. *Doğa Bilim Dergisi* A₂ 9(2): 419-427.