

1-1-1998

## Calculation of the Reaction Cross-Section for Proton-Nucleus Using Recently Optical Potential

M. M. OSMAN

Follow this and additional works at: <https://journals.tubitak.gov.tr/physics>



Part of the [Physics Commons](#)

---

### Recommended Citation

OSMAN, M. M. (1998) "Calculation of the Reaction Cross-Section for Proton-Nucleus Using Recently Optical Potential," *Turkish Journal of Physics*: Vol. 22: No. 6, Article 3. Available at: <https://journals.tubitak.gov.tr/physics/vol22/iss6/3>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Physics by an authorized editor of TÜBİTAK Academic Journals. For more information, please contact [academic.publications@tubitak.gov.tr](mailto:academic.publications@tubitak.gov.tr).

## Calculation of the Reaction Cross-Section for Proton-Nucleus Using Recently Optical Potential

M. M. OSMAN

*Department of Physics, Faculty of Science,  
Cairo University, Giza - EGYPT*

Received 11.12.1996

### Abstract

The eikonal approximation is used to calculate the reaction cross-section for proton scattering from different nuclei. The imaginary proton - nucleus optical potential is derived from relativistic mean field theory approach. We found a fair agreement with experimental data.

### 1. Introduction

Recently, successful attempts [1,2] have been made to apply the Dirac-Bruckner -Hartree-Fock (DBHF) approach [3] to nuclear matter problem<sup>5</sup>. Common to all DBHF result is that a repulsive relativistic many-body effect is obtained, which is strongly density dependent such that empirical nuclear matter properties can be explained starting from a realistic nucleon-nucleon interaction.

In spite of the success of DBHF approach to calculate nuclear matter properties, it is much more elaborate when applied to finite nuclei. For this reason, an alternative approach in which the DBHF results for nuclear matter are parameterized in terms of an effective Lagrangian which reproduce the binding energy, scalar and vector self-energy term as the original DBHF calculations. This approach has been used by Machleit et al to derive the proton-nucleus optical model potential. In the present paper, we apply this relativistic approach to calculate several proton-nucleus reaction cross-sections and compare our results with the experimental data. This enable us to show to what extent the method of Ref.[4] is successful in reproducing the proton-nucleus optical potential.

In the next section we briefly describe the method and in section III we compare our results for the reaction cross-section with experiment.

## 2. Theory

The nucleon self energy in a finite nucleus is obtained by means of the local density approximation, in which the spatial dependence of the microscopic optical potential is directly related to the density of the target nucleus. For a self-consistent description of the nucleon-nucleus scattering, the target density should also be determined from the effective Lagrangian [4],

$$L = \bar{\Psi}[i\gamma_\mu\partial^\mu - m - g_\sigma(\rho)\Phi_\sigma - g_w(\rho)\gamma_\mu\Psi_w^\mu]\Psi + \frac{1}{2}(\partial^\mu\Phi_\sigma)^2 - \frac{1}{2}m_\sigma^2\Phi_\sigma^2 - \frac{1}{4}(\partial_\mu\Phi_w^v - \partial_v\Phi_w^\mu)^2 + \frac{1}{2}m_w^2\Phi_w^{\mu^2}, \quad (1)$$


The Dirac equation for the single-particle motion of the projectile nucleon in the mean field of the target nucleus can be written as

$$[\alpha.K + \beta(m + U_S) + U_V + V_C]\Psi = E\Psi \quad (2)$$

with

$$U_S = \frac{\sum_S - m \sum_V}{1 + \sum_V}, U_V = \frac{\sum_O + E \sum_V}{1 + \sum_V}, \quad (3)$$

where E is the energy of the projectile in the center of mass (c.m) system of the projectile and nucleus, which is related to the incident energy  $T_{lab}$  by



**Figure 1.** Calculated reaction cross - section  $\sigma_R$ , together with experimental data [6], for the reaction of proton with  $^{13}\text{C}$  and  $^{27}\text{Al}$ .

$$E = \frac{m^2 + m_T(m + T_{lab})}{[(m + M_T)^2 + 2m_T T_{lab}]^{1/2}} \quad (4)$$

with  $m$  and  $m_T$  being the mass of the projectile and target, respectively.  $V_C$  is the coulomb field and is treated as in Ref.[5].

In terms of the scalar potential  $U_S$  and the vector potential  $U_V$ , the momentum of a nucleon propagating through a uniform nuclear medium can be determined from

$$E = [(m + U_S)^2 + K^2]^{1/2} + U_V \quad (5)$$

and can be written as

$$\frac{K^2}{2m} + V + iW = E - m + \frac{(E - m)^2}{2m}. \quad (6)$$

Here,  $V$  is the real part and  $W$  is imaginary part of the optical potential. They are given by[4]

$$V = U_{SR} + U_{VR} + \frac{(E - m)}{m} U_{VR} + \frac{1}{2m} (U_{SR}^2 + U_{VI}^2 - U_{SI}^2 - U_{VR}^2) \quad (7)$$

$$W = U_{SI} + U_{VI} + \frac{(E - m)}{m} U_{VI} + \frac{1}{m} (U_{SR} U_{SI} - U_{VR} U_{VI}), \quad (8)$$


where the real and imaginary part of the scalar and vector potential are given by  $U_S = U_{SR} + iU_{SI}$  and  $U_V = U_{VR} + iU_{VI}$ ,

To calculate the reaction cross-section for nucleon-nucleus system, we used the eikonal approximation

$$\sigma_R = 2\pi \int b db (1 - \exp[-4\delta_1(b)]) \quad (9)$$

where  $\delta_1(b)$  is the imaginary part of the nuclear elastic phase shift within the eikonal approximation,

$$\delta_1(b) = -\frac{1}{4} \frac{2\mu}{\hbar^2 K} \int dZ' W(\sqrt{b^2 + Z'^2}). \quad (10)$$



**Figure 2.** Same as Fig.(1) but for  $^{16}\text{O}$ ,  $^{40}\text{Ca}$ ,  $^{90}\text{Zr}$  and  $^{208}\text{Pb}$ , with experimental data are taken from Ref.[7].

### 3. Results and Discussion

The scalar potentials ( $U_{SR}$  and  $U_{SI}$ ) and the vector potentials ( $U_{VR}$  and  $U_{VI}$ ) can be calculated for protons scattered from nuclei from ref.(4). Using equations (9) and (10), we calculated the reaction cross - section  $\sigma_R$  for 20-800 MeV proton scattering from  $^{12}\text{C}$ ,  $^{16}\text{O}$ ,  $^{27}\text{Al}$ ,  $^{40}\text{Ca}$ ,  $^{90}\text{Zr}$ , and  $^{208}\text{pb}$ . Figure(1) shows the calculated  $\sigma_R$  together with the experimental data [6], for the reaction of proton with  $^{12}\text{C}$ , and  $^{27}\text{Al}$ . In figure (2), the calculated  $\sigma_R$  and the experimental data [7], for the reaction of proton with  $^{16}\text{O}$ ,  $^{27}\text{Al}$ ,  $^{40}\text{Ca}$ ,  $^{90}\text{Zr}$  and  $^{208}\text{pb}$ , are shown. From the figures, we can say that the calculated reaction cross-section  $\sigma_R$  for proton - nucleus systems using relativistic mean field theory approach is in fair agreement with the experimental results. This shows that this parameter free model can be used to describe the proton - nucleus scattering processes over a wide range of incident proton energy.

The author would to thank Prof. Dr. M. Ismail for helpful discussions.

### References

- [1] R. Brockmann and R. Machleidt, Phys. Lett., 149B (1984) 283; R. Machleidt and R. Brockmann, *ibid*, 160 (1985) 364.
- [2] R. Machleidt, Adv. Nucl. Phys., 19 (1989) 189; R-Brockmann and R Machleidt, Phys. Rev., c42 (1990) 1965.
- [3] M.R. Ansatasio, L.S. Celenza, W.S. Pong and C.M. Shakin, Phys. Rep, 100 (1983) 327; L.S. Celenza and C.M. Shakin, Relativistic Nuclear physics, theories of structure and scattering, Lecture Notes in Physics Vol.2 (World Scientific, Singapore, 1986).

OSMAN

- [4] G.Q.Li, R. Machleidt, R. Fritz, H. Muther and Y. Z. Zhuo, Phys. Rev. C48, (1993). 2443;  
G. Q. Li, R. Machleidt and Y. Z. Zhuo, Phys. Rev, C48, (1993) 1062.
- [5] D. P. Murdock and C. J. Horowitz, Phys. Rev., C 83 (1987) 1442.
- [6] L.Sihver, et. al., Phys. Rev, C47 (1993) 1225.
- [7] W. Bauhoff, At. Data Nucl. Data Tables, 35 (1986) 429.