

1-1-2014

A class of uniquely (strongly) clean rings

ORHAN GÜRGÜN

AYŞE ÇİĞDEM ÖZCAN

Follow this and additional works at: <https://journals.tubitak.gov.tr/math>



Part of the [Mathematics Commons](#)

Recommended Citation

GÜRGÜN, ORHAN and ÖZCAN, AYŞE ÇİĞDEM (2014) "A class of uniquely (strongly) clean rings," *Turkish Journal of Mathematics*: Vol. 38: No. 1, Article 5. <https://doi.org/10.3906/mat-1209-9>
Available at: <https://journals.tubitak.gov.tr/math/vol38/iss1/5>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Mathematics by an authorized editor of TÜBİTAK Academic Journals. For more information, please contact academic.publications@tubitak.gov.tr.

A class of uniquely (strongly) clean rings

Orhan GÜRGÜN^{1,*}, Ayşe Çiğdem ÖZCAN²

¹Department of Mathematics, Ankara University, Ankara, Turkey

²Department of Mathematics, Hacettepe University, Beytepe Ankara, Turkey

Received: 10.09.2012 • Accepted: 03.01.2013 • Published Online: 09.12.2013 • Printed: 20.01.2014

Abstract: In this paper we call a ring R δ_r -clean if every element is the sum of an idempotent and an element in $\delta(R_R)$ where $\delta(R_R)$ is the intersection of all essential maximal right ideals of R . If this representation is unique (and the elements commute) for every element we call the ring *uniquely (strongly) δ_r -clean*. Various basic characterizations and properties of these rings are proved, and many extensions are investigated and many examples are given. In particular, we see that the class of δ_r -clean rings lies between the class of uniquely clean rings and the class of exchange rings, and the class of uniquely strongly δ_r -clean rings is a subclass of the class of uniquely strongly clean rings. We prove that R is δ_r -clean if and only if $R/\delta_r(R_R)$ is Boolean and $R/Soc(R_R)$ is clean where $Soc(R_R)$ is the right socle of R .

Key words: Clean ring, strongly clean ring, uniquely clean ring, strongly J-clean ring

1. Introduction

Clean rings have been studied by many ring and module theorists since 1977, and it is still a very popular subject. They were defined by Nicholson as a subclass of exchange rings. An associative ring with unity is called *clean* if every element is the sum of an idempotent and a unit [14]. If this representation is unique for every element, Nicholson and Zhou [17] call the ring *uniquely clean*. They proved that a ring R is uniquely clean if and only if for all $a \in R$ there exists a unique idempotent $e \in R$ such that $a - e \in J(R)$ where $J(R)$ is the Jacobson radical of R (we call the ring with this property *uniquely J-clean*). Chen et al. [7] call a ring *uniquely strongly clean* if every element can be written uniquely as the sum of an idempotent and a unit that commute. They proved that R is uniquely strongly clean if and only if for every $a \in R$, there exists a unique idempotent $e \in R$ such that $a - e \in J(R)$ and $ae = ea$ (we call the ring with this property *uniquely strongly J-clean*). Recently, Chen [6] defined strongly J -clean rings. A ring R is called *strongly J-clean* if for all $a \in R$ there exists an idempotent $e \in R$ such that $a - e \in J(R)$ and $ea = ae$ [6]. Note that strongly J -clean rings are strongly clean but the converse need not be true [6, Proposition 2.1 and Example 2.2].

These results motivate us to define the class of uniquely $\delta(R_R)$ -clean and uniquely strongly $\delta(R_R)$ -clean rings where $\delta(R_R)$ is the ideal defined by Zhou [21]. These classes of rings give some new classes of uniquely clean and uniquely strongly clean rings and also give some ideas on the cleanness of $R/Soc(R_R)$ where $Soc(R_R)$ is the right socle of R . Firstly basic properties of $\delta(R_R)$ -clean rings are given in Section 2. Interestingly we see that the class of $\delta(R_R)$ -clean rings lies between the class of uniquely clean rings and exchange rings. We also

*Correspondence: orhangurgun@gmail.com

2010 AMS Mathematics Subject Classification: 16S50, 16S70, 16U99.

prove that if R is $\delta(R_R)$ -clean, then $R/Soc(R_R)$ is clean and partially unit regular, i.e. every regular element is unit regular. In Section 3, uniquely $\delta(R_R)$ -clean rings are studied. We see that any uniquely $\delta(R_R)$ -clean ring is uniquely clean. Contrary to the result in [17] saying that R is uniquely clean if and only if $R[[x]]$ is uniquely clean, just the necessity is true for uniquely $\delta(R_R)$ -clean rings. Section 4 is devoted to uniquely strongly $\delta(R_R)$ -clean rings (USDC for short). Any uniquely $\delta(R_R)$ -clean ring is USDC, and any USDC ring is uniquely strongly clean. We prove that if R is a commutative ring, then R is USDC if and only if the ring of 2×2 upper triangular matrices, $T_2(R)$, is USDC. In the last section $\delta(R_R)$ -cleanness of the formal triangular matrix ring is investigated.

Recall some definitions. Following [21], a submodule N of a module M is called δ -small in M (denoted by $N \ll_{\delta} M$) if $N + K \neq M$ for any submodule K of M with M/K singular. Denote $\delta(M)$ to be the sum of all δ -small submodules of M (see [21, Lemma 1.5]). We use δ_r (or $\delta_r(R)$) for $\delta(R_R)$ for a ring R . Clearly $J(R) \subseteq \delta_r(R) \ll_{\delta} R_R$. If S is simple and M is essential, then $S \cap M$ must equal S (as it cannot be zero). Since every simple right ideal is contained in every essential right ideal, then $S_r := Soc(R_R) \subseteq \delta_r(R)$ (see also [21, Lemma 1.9]). By view of [21, Corollary 1.7], $J(R/S_r) = \delta_r/S_r$; in particular, R is semisimple if and only if $\delta(R_R) = R$.

A ring R is an *exchange ring* if, for every $a \in R$, there exists an idempotent $e \in aR$ such that $1 - e \in (1 - a)R$ (see [14]). For example, (von Neumann) regular rings and clean rings are exchange. If I is a left ideal of a ring R , *idempotents lift modulo I* if, given $a \in R$ with $a^2 - a \in I$, there exists $e^2 = e \in R$ such that $a - e \in I$ [14]. Note that R is an exchange ring if and only if idempotents lift modulo every left ideal of R [14, Corollary 1.3]. A ring R is called δ -semiregular if R/δ_r is a regular ring and idempotents lift modulo δ_r [21, Theorem 3.5]. A ring R is called *abelian* if every idempotent of R is central.

Throughout this article, all rings are associative with unity and all modules are unitary. We denote $S_r = Soc(R_R)$ and $Z_r = Z(R_R)$ for the right socle and the right singular ideal of a ring R . We write J (or $J(R)$) for the Jacobson radical of R . $U(R)$ is the set of all units in R . The ring of integers modulo n is denoted by \mathbb{Z}_n , and we write $M_n(R)$ (resp. $T_n(R)$) for the rings of all (resp., all upper triangular) $n \times n$ matrices over the ring R .

2. δ_r -clean rings

Chen [6] calls a ring R *strongly J -clean* if for every element $a \in R$ there exists an idempotent $e \in R$ such that $a - e \in J$ and $ea = ae$. Call a ring R *J -clean* if for any element $a \in R$, there exists an idempotent $e \in R$ such that $a - e \in J$.

Any J -clean ring is clean. Let $a \in R$ and $a = e + w$ where $e^2 = e \in R$, $w \in J$. Then $a = (1 - e) + (2e - 1 + w)$. Since $(2e - 1)^2 = 1$ we see that $a - (1 - e) \in U(R)$ (see [6, Proposition 2.1]). It is easy to give an example of a ring that is clean but not J -clean (e.g., \mathbb{Z}_3). Now we introduce the notion of δ_r -clean rings.

Definition 2.1 A ring R is called δ_r -clean if for every element $a \in R$ there exists an idempotent $e \in R$ such that $a - e \in \delta_r$.

The class of δ_r -clean rings contains Boolean rings, semisimple rings, and J -clean rings. Clearly, R is δ_r -clean if and only if R/δ_r is Boolean and idempotents lift modulo δ_r . Note that there exists a ring R with R/δ_r is Boolean but such that idempotents do not lift modulo δ_r . There is a ring R with $R/J(R)$ Boolean

but such that idempotents do not lift modulo $J(R)$ (see [13, Example 15]). In this ring, idempotents do not lift modulo δ_r , for, if they did, then R would be δ_r -clean and therefore exchange, by Theorem 2.2 below. Then idempotents would lift modulo $J(R)$, a contradiction.

On the other hand, if R is δ_r -clean, then R/J need not be a Boolean ring. For example, \mathbb{Z}_3 is semisimple but not Boolean.

Theorem 2.2 *If R is a δ_r -clean ring, then*

- 1) R/S_r is a semiregular ring, i.e. R is δ_r -semiregular;
- 2) R is an exchange ring;
- 3) R/S_r is a clean ring;
- 4) $Z_r \subseteq J$.

Proof 1) Since R/δ_r is a Boolean ring and idempotents lift modulo δ_r , R is δ -semiregular. By [19, Theorem 1.4], R is δ_r -semiregular if and only if R/S_r is semiregular.

2) If R/S_r is semiregular, then R is exchange by [19, Corollary 1.5].

3) If R is δ_r -clean, then R/S_r is $J(R/S_r)$ -clean since $J(R/S_r) = \delta_r/S_r$. Any J -clean ring is clean. We thus conclude that R/S_r is a clean ring.

4) Since R is δ_r -semiregular, $Z_r \subseteq \delta_r$ by [16, Theorem 1.2]. Then Z_r is δ -small in R . This gives that Z_r is small in R . Hence, $Z_r \subseteq J$. □

Example 2.3 *If R is a semisimple ring that is not a Boolean ring (e.g., \mathbb{Z}_3), then R is δ_r -clean but not J -clean since $J = 0$ and $\delta_r = R$.*

Example 2.4 *There exist clean rings that are not δ_r -clean.*

Proof 1) Let V_D be a nonzero vector space over a division ring D and let $R = \text{End}_D(V)$. Then R is regular (see [1, Exercise 15.13]) and clean [15, Lemma 1] (see also [3, Lemma 3.1]) and $S_r = S_l = \{f \in R \mid \text{rank } f < \infty\}$ (see [1, Exercise 18.4]). Since $J(R/S_r) = \delta_r/S_r$ and R is regular, we have that $\delta_r = S_r$.

Now assume that V_D is a countably infinite dimensional vector space and let $\{v_1, v_2, \dots\}$ be a basis of V . Define the shift operator f on V by $f(v_n) = v_{n+1}$ for $n = 1, 2, 3, \dots$. Then $f^2 - f \notin S_r$. This shows that $R/S_r = R/\delta_r$ is not Boolean. Hence, R is not δ_r -clean.

2) Let p be a prime integer and consider the local ring $\mathbb{Z}_{(p)} = \{\frac{m}{n} \mid m, n \in \mathbb{Z}, (m, n) = 1, p \nmid n\}$. Since $\mathbb{Z}_{(p)}$ is not semisimple, $J = \delta_r = p\mathbb{Z}_{(p)}$. Then $\mathbb{Z}_{(p)}$ is clean but not δ_r -clean, because $\mathbb{Z}_{(p)}/\delta_r$ is not Boolean. □

Note that any clean ring is exchange [14, Proposition 1.8]. Bergman's example is an example of an exchange ring that is not clean. We prove below that this ring is not δ_r -clean, and so we pose the following question.

Question: Is any δ_r -clean ring clean?

Example 2.5 (Bergman) Let F be a field with $\text{char}(F) \neq 2$, and $A = F[[x]]$. Let Q be the field of fractions of A . Define

$$R = \{r \in \text{End}_F(A) \mid \exists q \in Q \text{ and } \exists n > 0 \text{ with } r(a) = qa \text{ for all } a \in x^n A\}.$$

Then R is a regular (so exchange) ring [10], but not clean [4]. There is also an epimorphism $\theta : R \rightarrow Q$ given by $r \mapsto q$, where r agrees with q on $x^n A$ for some $n > 0$ with $\text{Ker } \theta = S_r = \delta_r$ (see [12, Example 1]). Now assume that R is δ_r -clean. Then, for any $r \in R$, there exists an idempotent $e \in R$ such that $r - e \in \delta_r$. This gives that $\theta(r - e) = \theta(r) - \theta(e) = 0$ and $\theta(r) = \theta(e)$ is an idempotent in Q . Since Q is a field, $\theta(r) = 0$ or 1 , which contradicts the fact that θ is an epimorphism. Therefore, R is not δ_r -clean.

Thus we conclude that

$$\{ \text{Boolean} \} \subsetneq \{ J\text{-clean} \} \subsetneq \{ \delta_r\text{-clean} \} \subsetneq \{ \text{exchange} \}.$$

Now we give a few conditions for a δ_r -clean ring to be clean or J -clean. First note that Baccella [2] proved the important fact that idempotents lift modulo S_r for any ring R .

Proposition 2.6 *Any δ_r -clean ring R is J -clean if*

- 1) R/J is Boolean, or 2) $S_r \subseteq J$.

Proof 1) Assume that R is δ_r -clean and R/J is Boolean. Let $a \in R$. Then $a^2 - a \in J$. By Theorem 2.2, idempotents lift modulo J . Hence, there exists an idempotent $e \in R$ such that $a - e \in J$.

2) Assume that R is δ_r -clean. If $S_r \subseteq J$, then $J/S_r = J(R/S_r) = \delta_r/S_r$, and we have that $J = \delta_r$. Hence, R is J -clean. \square

Proposition 2.7 *If R is δ_r -clean and R/J is abelian, then R is clean.*

Proof Assume that R is δ_r -clean. According to Theorem 2.2, R is exchange and so R/J is exchange and idempotents lift modulo J by [14, Corollary 1.3]. Thus, R/J is abelian exchange and it is clean by [14, Proposition 1.8]. By [9, Proposition 6], R is clean. \square

Recall that a ring R is called *right quasi-duo* if every maximal right ideal is a 2-sided ideal. If R is an exchange ring, then R/J is right quasi-duo iff R/J is reduced iff R/J is abelian [20, Proposition 4.1]. Hence, the following corollary is immediate.

Corollary 2.8 *If R is δ_r -clean and right (or left) quasi-duo, then R is clean.*

Proposition 2.9 *Let R be a ring with only trivial idempotents (e.g., a local ring). Then R is δ_r -clean if and only if R is either a division ring or $R/J(R) \cong \mathbb{Z}_2$.*

Proof Assume that R is δ_r -clean. Then R is exchange by Theorem 2.2. Since R is exchange and has only trivial idempotents, R is local. Then either $J(R) = 0$ or $J(R) = \delta_r$. If $J(R) = 0$, then R is a division ring. If $J(R) = \delta_r$, then R is J -clean and so R is strongly J -clean by hypothesis. Hence, $R/J(R) \cong \mathbb{Z}_2$ by [6, Lemma 4.2]. Conversely, if R is a division ring, then R is semisimple and so R is δ_r -clean. If $R/J(R) \cong \mathbb{Z}_2$, then R is J -clean by [17, Theorem 15] and so R is δ_r -clean. \square

A characterization of δ_r -clean rings can be given as follows.

Theorem 2.10 *Let R be a ring. The following statements are equivalent.*

- 1) R is δ_r -clean.
- 2) R/S_r is J -clean.

3) R/δ_r is Boolean and R/S_r is clean.

Proof Since $J(R/S_r) = \delta_r/S_r$, (1) \Leftrightarrow (2). By Theorem 2.2, (1) \Rightarrow (3).

(3) \Rightarrow (1) Let $a \in R$. Then $a^2 - a \in \delta_r$. Since $\bar{R} = R/S_r$ is clean, idempotents of $\bar{R}/J(\bar{R})$ lift to idempotents of \bar{R} . By [19, Lemma 1.3], idempotents of R/δ_r lift to idempotents of R . Hence, there exists $e^2 = e \in R$ such that $a - e \in \delta_r$. Thus, R is δ_r -clean. \square

Bergman's example (see Example 2.5) also shows that if R/S_r is a clean ring, then R need not be clean [12, Example 1].

Recall that a ring R is said to have *stable range 1*, written $sr(R) = 1$, if given $a, b \in R$ for which $aR + bR = R$, there exists a $y \in R$ such that $a + by \in U(R)$. It is obvious that $sr(R) = 1$ if and only if $sr(R/J) = 1$.

Lemma 2.11 *Let R be a ring. Then $sr(R/\delta_r) = 1$ if and only if $sr(R/S_r) = 1$.*

Proof It can be easily seen by the fact that $J(R/S_r) = \delta_r/S_r$. \square

Recall that an element a of a ring R is called *regular* (resp., *unit regular*) if there exists $u \in R$ (resp., $u \in U(R)$) such that $a = aua$. A ring R is called *partially unit regular* if every regular element of R is unit regular. These rings are also called *IC-ring* in [11].

Theorem 2.12 *If R is a δ_r -clean ring, then R/S_r is partially unit regular.*

Proof Since R/δ_r is a Boolean ring, $sr(R/\delta_r) = 1$. By Theorem 2.2, R is an exchange ring. Hence, by Lemma 2.11 and [5, Theorem 3], R/S_r is partially unit regular. \square

The following example shows that if R is δ_r -clean, then R/S_r need not be a regular ring in general.

Example 2.13 *Let $R = \mathbb{Z}_8$. Then $Soc(R) = 4R$ and $J = 2R$. It is clear that R is J -clean, but since $J \not\subseteq Soc(R)$, $R/Soc(R)$ is not regular.*

3. Uniquely δ_r -clean rings

Definition 3.1 A ring R is called *uniquely δ_r -clean* if for every element $a \in R$ there exists a unique idempotent $e \in R$ such that $a - e \in \delta_r$.

Let I be an ideal of R . Then *idempotents lift uniquely modulo I* if whenever $a^2 - a \in I$, there exists a unique idempotent $e \in R$ such that $e - a \in I$ [17]. This condition implies that if $e - f \in I$, $e^2 = e$, $f^2 = f$, then $e = f$; in particular, 0 is the only idempotent in I .

Clearly, R is uniquely δ_r -clean if and only if R/δ_r is Boolean and idempotents lift uniquely modulo δ_r .

Theorem 3.2 *If R is uniquely δ_r -clean, then the following hold.*

1) $\delta_r = J$.

2) R is uniquely clean.

Proof 1) Since idempotents lift uniquely modulo δ_r , by the remark above, the only idempotent in δ_r is 0. Now let $a \in \delta_r$. Then there exists a semisimple right ideal Y of R such that $R = (1-a)R \oplus Y$ by [21, Theorem 1.6]. Since $Y \subseteq S_r \subseteq \delta_r$, we have that $Y = 0$. Hence $1-a$ is right invertible in R , and so $a \in J$.

2) It is clear by (1) and [17, Theorem 20]. □

Note that any uniquely clean ring is abelian by [17, Lemma 4].

Examples 3.3 1) *No semisimple ring is uniquely δ_r -clean, for, if R is a semisimple ring, then $\delta_r = R$ and for any $a \in R$, $a-0 \in R$ and $a-1 \in R$.*

2) *If $R \not\cong \mathbb{Z}_2$, then $R/J \cong \mathbb{Z}_2$ if and only if R is local uniquely δ_r -clean, for, if $R/J \cong \mathbb{Z}_2$, then $J = \delta_r$ and R is uniquely clean by [17, Theorem 15] and so R is uniquely δ_r -clean. The converse is also true by Proposition 2.9.*

Therefore, for example, the rings $R = \left\{ \begin{bmatrix} a & b \\ 0 & a \end{bmatrix} \mid a, b \in \mathbb{Z}_2 \right\}$, $R = \left\{ \begin{bmatrix} x & y \\ 0 & x \end{bmatrix} \mid x \in \mathbb{Z}_4, y \in \mathbb{Z}_4 \oplus \mathbb{Z}_4 \right\}$, or $R = \mathbb{Z}_{2^n}$ where $1 \neq n \in \mathbb{N}$ are uniquely δ_r -clean.

Uniquely clean rings need not be uniquely δ_r -clean.

Example 3.4 1) \mathbb{Z}_2 is uniquely clean but not uniquely δ_r -clean.

2) Let $R = \prod_{i=1}^{\infty} R_i$ where $R_i \cong \mathbb{Z}_2$ for all $i = 1, 2, \dots$. Then R is a Boolean ring with $S_r = \bigoplus_{i=1}^{\infty} R_i$. Since R/S_r is Boolean, $J(R/S_r) = 0$ and so $S_r = \delta_r$. Clearly R is uniquely J -clean, that is, uniquely clean but not uniquely δ_r -clean.

It is easy to see that every uniquely clean ring is δ_r -clean by the fact that R is uniquely clean if and only if R is uniquely J -clean [17, Theorem 20]. But if R is a semisimple ring that is not Boolean, then R is δ_r -clean but not uniquely clean (see Example 2.3).

Thus, we conclude that

$$\{ \text{uniquely } \delta_r\text{-clean} \} \subsetneq \{ \text{uniquely clean} \} \subsetneq \{ \delta_r\text{-clean} \} \subsetneq \{ \text{exchange} \}.$$

If $S_r \subseteq J$ for a ring R , then $J/S_r = J(R/S_r) = \delta_r/S_r$ and so $J = \delta_r$. Hence, Proposition 3.5 below is obvious by Proposition 2.6.

Proposition 3.5 *If R is a uniquely clean ring with $S_r \subseteq J$, then R is uniquely δ_r -clean.*

By [17, Theorem 20] we know that R is uniquely clean if and only if R/J is Boolean, R is abelian, and idempotents lift modulo J . However, this result cannot be restated for δ_r in general. The following theorem and examples prove our claim.

Theorem 3.6 *Let R be a ring and consider the following conditions.*

- 1) R is uniquely δ_r -clean.
- 2) R/δ_r is Boolean, R is abelian, and idempotents lift modulo δ_r .
- 3) R/δ_r is Boolean, R/S_r is abelian, and idempotents lift modulo δ_r .

4) R/S_r is uniquely clean.

Then (1) \Rightarrow (2) \Rightarrow (3) \Leftrightarrow (4).

Proof (1) \Rightarrow (2) Since R is uniquely clean, it is abelian by [17, Lemma 4].

(2) \Rightarrow (3) Since idempotents always lift modulo S_r , it is clear.

(3) \Leftrightarrow (4) It is by [17, Theorem 20]. Note that idempotents lift modulo $J(R/S_r)$ if and only if idempotents lift modulo δ_r [19, Lemma 1.3]. □

In Theorem 3.6, (2) $\not\Rightarrow$ (1) in general.

Example 3.7 We consider again the ring $R = \prod_{i=1}^{\infty} R_i$ where $R_i \cong \mathbb{Z}_2$, $i = 1, 2, \dots$ (see Example 3.4). Since R is uniquely clean, R is abelian and δ_r -clean. But R is not uniquely δ_r -clean.

In Theorem 3.6, (4) $\not\Rightarrow$ (2) in general.

Example 3.8 Let $R = \begin{bmatrix} \mathbb{Z}_2 & \mathbb{Z}_2 \\ 0 & \mathbb{Z}_2 \end{bmatrix}$. Then $S_r = \delta_r = \begin{bmatrix} 0 & \mathbb{Z}_2 \\ 0 & \mathbb{Z}_2 \end{bmatrix}$ and $R/S_r \cong \mathbb{Z}_2$ is Boolean. Obviously R is δ_r -clean but not abelian.

Theorem 3.9 If R is uniquely δ_r -clean and $e^2 = e \in R$, then eRe is uniquely δ_r -clean.

Proof Since R is abelian, $\delta_r(eRe) = e\delta_r e$ by [18, Theorem 3.11]. By Theorem 3.2, $\delta_r = J$, so we have that $J(eRe) = eJe = \delta_r(eRe)$. If R is uniquely δ_r -clean, then R is uniquely clean by Theorem 3.2. By [17, Corollary 6], eRe is uniquely clean. By [17, Theorem 20], eRe is uniquely δ_r -clean. □

Although every factor ring of a uniquely clean ring is uniquely clean [17, Theorem 22], the same property does not hold for uniquely δ_r -clean.

Remark 3.10 1) If R is a uniquely δ_r -clean ring, then factor rings of R need not be uniquely δ_r -clean in general. For example, if $R \not\cong \mathbb{Z}_2$ and $R/J \cong \mathbb{Z}_2$, then R is uniquely δ_r -clean by Example 3.3, but R/J is not uniquely δ_r -clean.

(2) Since matrix ring $M_n(R)$ and upper triangular matrix ring $T_n(R)$ are not abelian for $n \geq 2$, they are not uniquely δ_r -clean by Theorem 3.2.

Let R be a ring and V an (R, R) -bimodule that is a general ring (possibly with no unity) in which $(vw)r = v(wr)$, $(vr)w = v(rw)$, and $(rv)w = r(vw)$ hold for all $v, w \in V$ and $r \in R$. Then the *ideal-extension* (also called the Dorroh extension) $I(R; V)$ of R by V is defined to be the additive abelian group $I(R; V) = R \oplus V$ with multiplication $(r, v)(s, w) = (rs, rw + vs + vw)$.

Uniquely clean ideal-extensions are considered in [17, Proposition 7]. Now we deal with uniquely δ_r -clean ideal-extensions.

Proposition 3.11 An ideal-extension $S = I(R; V)$ is uniquely δ_r -clean if the following conditions are satisfied:

- 1) R is uniquely δ_r -clean;
- 2) if $e^2 = e \in R$ then $ev = ve$ for all $v \in V$;
- 3) if $v \in V$ then $v + w + vw = 0$ for some $w \in V$.

Proof Assume that (1), (2), and (3) are satisfied. Since R is uniquely δ_r -clean, R is uniquely clean by Theorem 3.2 and so S is uniquely clean by [17, Proposition 7]. Then S is δ_r -clean. Note by the proof of [17, Proposition 7] that any idempotent in S is of the form $(e, 0)$ where $e^2 = e \in R$. Now suppose that $(e, 0) + (u, v) = (e_1, 0) + (u_1, v_1)$ in S where $(e, 0)$ and $(e_1, 0)$ are idempotents and $(u, v), (u_1, v_1) \in \delta_r(S)$. Then $e + u = e_1 + u_1$ in R where e and e_1 are idempotents in R and $u, u_1 \in \delta_r(R)$ by the following result, and so $(e, 0) = (e_1, 0)$ by (1).

Claim. If $(u, v) \in \delta_r(S)$ then $u \in \delta_r(R)$.

Proof. Let $(u, v) \in \delta_r(S)$. Then $(u, 0) \in \delta_r(S)$ because $(0, V) \subseteq J(S) \subseteq \delta_r(S)$ by (3). Let L be a right ideal of R such that $uR + L = R$. It is enough to show that L is a direct summand of R by [21, Theorem 1.6]. Since $(u, 0)S + (L \oplus V) = S$ and $(u, 0) \in \delta_r(S)$, we have that $L \oplus V$ is a direct summand of S and so is generated by an idempotent $(e, 0) \in S$ where $e^2 = e \in R$. Then we see that $L = eR$, and hence L is a direct summand of R , as desired. \square

Example 3.12 Let R be a uniquely δ_r -clean ring and let $S = \{[a_{ij}] \in T_n(R) \mid a_{11} = \dots = a_{nn}\}$. Then S is uniquely δ_r -clean and is noncommutative if $n \geq 3$.

Proof If $V = \{[a_{ij}] \in T_n(R) \mid a_{11} = \dots = a_{nn} = 0\}$, then $S \cong I(R; V)$. The conditions in Proposition 3.11 hold as in [17, Example 8]. \square

If R is a ring and $\sigma : R \rightarrow R$ is a ring endomorphism, let $R[[x, \sigma]]$ denote the ring of skew formal power series over R , that is, all formal power series in x with coefficients from R with multiplication defined by $xr = \sigma(r)x$ for all $r \in R$. In particular, $R[[x]] = R[[x, 1_R]]$ is the ring of formal power series over R . Since $R[[x, \sigma]] \cong I(R; \langle x \rangle)$ where $\langle x \rangle$ is the ideal generated by x , the proof of [17, Example 9] and Proposition 3.11 give the next results.

Corollary 3.13 Let R be a ring and $\sigma : R \rightarrow R$ a ring endomorphism and $e = \sigma(e)$ for all $e^2 = e \in R$. If R is uniquely δ_r -clean, then $R[[x, \sigma]]$ is uniquely δ_r -clean

Corollary 3.14 If R is a uniquely δ_r -clean ring, then $R[[x]]$ is uniquely δ_r -clean.

Corollary 3.14 can be proven by using Proposition 3.15 below, for, if R is uniquely δ_r -clean, then $R[[x]]$ is a uniquely clean ring by Theorem 3.2 and [17, Corollary 10]. By Proposition 3.15, $J(R[[x]]) = J(R) + \langle x \rangle \subseteq \delta_r(R[[x]]) \subseteq \delta_r(R) + \langle x \rangle$. Then since $J(R) = \delta_r(R)$ by Theorem 3.2(1), $J(R[[x]]) = \delta_r(R[[x]])$. Hence, $R[[x]]$ is a uniquely δ_r -clean ring.

Proposition 3.15 Let R be a ring. Then $\delta_r(R[[x]]) \subseteq \delta_r(R) + \langle x \rangle$.

Proof Let $f(x) = a_0 + a_1x + a_2x^2 + \dots \in \delta_r(R[[x]])$. Since $\langle x \rangle \subseteq J(R[[x]])$, $a_0 \in \delta_r(R[[x]])$. Let L be a right ideal of R such that $a_0R + L = R$. It is enough to show that L is a direct summand of R by [21, Theorem 1.6]. Since $a_0R[[x]] + L[[x]] = R[[x]]$ and $a_0 \in \delta_r(R[[x]])$, we have that $L[[x]]$ is a direct summand of $R[[x]]$ and so is generated by an idempotent $e(x) = e_0 + e_1x + e_2x^2 + \dots \in R[[x]]$. Then e_0 is an idempotent in R and it can be seen that $L = e_0R$. Thus, $a_0 \in \delta_r(R)$, as desired. \square

Note that $J(\mathbb{Z}_2[[x]]) = \delta_r(\mathbb{Z}_2[[x]]) \subsetneq \delta_r(\mathbb{Z}_2) + \langle x \rangle = \mathbb{Z}_2[[x]]$.

Corollary 3.16 *If $R[[x]]$ is δ_r -clean, then R is δ_r -clean.*

Proof Let $a \in R$. Then there exist $e(x)^2 = e(x) \in R[[x]]$ and $w(x) \in \delta_r(R[[x]])$ such that $a = e(x) + w(x)$ and so $w(0) \in \delta_r(R)$ by Proposition 3.15. Thus, $a = e(0) + w(0)$ where $e(0)^2 = e(0) \in R$, as asserted. \square

If $R[[x]]$ is uniquely δ_r -clean, then R need not be uniquely δ_r -clean. For example, \mathbb{Z}_2 is not uniquely δ_r -clean but since $\mathbb{Z}_2[[x]]/J(\mathbb{Z}_2[[x]]) \cong \mathbb{Z}_2$, $\mathbb{Z}_2[[x]]$ is uniquely δ_r -clean by Example 3.3(2).

4. Uniquely strongly δ_r -clean rings

Uniquely strongly clean rings were studied in [7]. A ring R is called *uniquely strongly clean* if for every element $a \in R$ there exists a unique idempotent $e \in R$ such that $a - e \in U(R)$ and $ea = ae$. In Theorem 17 of [7] it is proven that a uniquely strongly clean ring is exactly the same as a uniquely strongly J -clean, i.e. for any $a \in R$ there exists a unique idempotent $e \in R$ such that $a - e \in J$ and $ea = ae$.

Definition 4.1 A ring R is called *uniquely strongly δ_r -clean* if for every element $a \in R$ there exists a unique idempotent $e \in R$ such that $a - e \in \delta_r$ and $ea = ae$.

Proposition 4.2 *A ring R is uniquely δ_r -clean if and only if R is an abelian USDC ring.*

Proof Since uniquely δ_r -clean rings are abelian by Theorem 3.6, the proof is obvious. \square

Proposition 4.3 *Let R be a USDC ring. Then the following hold:*

- 1) *If $e^2 = e \in \delta_r$, then $e = 0$.*
- 2) *R/J is Boolean.*
- 3) *$\delta_r = J$.*
- 4) *R is uniquely strongly clean.*

Proof 1) Let $e^2 = e \in \delta_r$. Then $e + 0 = 0 + e$ and $0.e = e.0$ yield $e = 0$.

2) R is exchange by Theorem 2.2. If we show that every nonzero idempotent of R is not the sum of 2 units, then by [13, Theorem 13], R/J will be Boolean. Let e be a nonzero idempotent in R . Write $e = u + v$, where $u, v \in U(R)$. Since R is USDC, R/δ_r is Boolean and so $2 \in \delta_r$. Therefore, u and v are congruent to 1, modulo δ_r , which means that their sum is in δ_r . This contradicts with (1).

3) Let $a \in \delta_r$. Since R/J is Boolean, $a^2 - a \in J$. By Theorem 2.2, R is exchange and so idempotents lift modulo J . Thus, there exist $e^2 = e \in R$ such that $a - e \in J$. Since $J \subseteq \delta_r$, $e = 0$ by (1). Hence, $a \in J$, as asserted.

4) It is clear by (3) and [7, Theorem 17]. \square

However, a uniquely strongly clean ring need not be USDC. The ring $R = \begin{bmatrix} \mathbb{Z}_2 & \mathbb{Z}_2 \\ 0 & \mathbb{Z}_2 \end{bmatrix}$ is uniquely strongly clean by [7, Theorem 10] but not USDC by Example 3.8.

Thus, we conclude that

$$\{ \text{uniquely } \delta_r\text{-clean} \} \subsetneq \{ \text{USDC} \} \subsetneq \{ \text{uniquely strongly clean} \} \subsetneq \{ \delta_r\text{-clean} \}.$$

The first and the last containments above are proper because, for example, the ring \mathbb{Z}_p where $2 \neq p$ is a prime is δ_r -clean but not uniquely strongly clean because $J(\mathbb{Z}_p) = 0$ and \mathbb{Z}_p is not Boolean. If R is a commutative uniquely δ_r -clean ring, then $T_n(R)$ is USDC by Theorem 4.5 for any $n \in \mathbb{N}$, but $T_n(R)$ is never uniquely δ_r -clean by Remark 3.10(2).

Any factor ring of any USDC ring need not be USDC. For example, since \mathbb{Z}_4 is uniquely δ_r -clean by Example 3.3, it is USDC by Proposition 4.2. However, $\mathbb{Z}_4/J(\mathbb{Z}_4) \cong \mathbb{Z}_2$ is not USDC by Proposition 4.2 and Example 3.3.

Proposition 4.4 *Let e be an idempotent of a ring R such that $eR = eRe$ (i.e. right semicentral) or $ReR = R$ (i.e. full idempotent). If R is USDC, then eRe is USDC.*

Proof Assume that R is USDC. For any idempotent e of R , eRe is uniquely strongly clean by Proposition 4.3(4) and [7, Example 5]. Since uniquely strongly clean rings are uniquely strongly J -clean, for any $a \in eRe$, there exists an idempotent $f \in eRe$ and $v \in \delta_r(eRe)$ such that $a = f + v$ and $fv = vf$. It remains to show the uniqueness. Let $a = f + v = g + w$ where f and g are idempotents in eRe and $v, w \in \delta_r(eRe)$ such that $fv = vf$ and $gw = wg$. If e is an idempotent as in the hypothesis, then $\delta_r(eRe) \subseteq e\delta_re \subseteq \delta_r(R)$ by [18, Theorems 3.9 and 3.11]. Hence, by assumption, $f = g$. \square

Since $M_n(R)$ is never uniquely strongly clean by [7, Lemma 6], $M_n(R)$ is never USDC.

Theorem 4.5 *Let R be a commutative ring. Then the following are equivalent.*

- (1) R is USDC.
- (2) R is uniquely δ_r -clean.
- (3) $T_n(R)$ is USDC for all $n \geq 1$.
- (4) $T_2(R)$ is USDC.

Proof (1) \Leftrightarrow (2) This follows by Proposition 4.2.

(3) \Rightarrow (4) It is clear.

(4) \Rightarrow (1) Suppose that $T_2(R)$ is USDC and let $e = \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} \in T_2(R)$. Since e is right semicentral and $eT_2(R)e \cong R$, R is USDC by Proposition 4.4.

(1) \Rightarrow (3) If R is USDC, then $T_n(R)$ is uniquely strongly clean by Proposition 4.3(4) and [7, Theorem 10]. According to Proposition 4.3(3) and Lemma 5.1, $\delta_r(T_n(R)) = J(T_n(R))$ and so $T_n(R)$ is USDC by [7, Theorem 17]. Therefore, the proof is completed. \square

5. On the formal triangular matrix rings

Let S and T be any ring, M an (S, T) -bimodule, and R the formal triangular matrix ring $\begin{bmatrix} S & M \\ 0 & T \end{bmatrix}$. It

is well known that $J(R) = \begin{bmatrix} J(S) & M \\ 0 & J(T) \end{bmatrix}$ (e.g., [8, Corollary 2.2]), but for $\delta_r(R)$ the similar property does

not hold in general. For example, if $S = M = T = F$ is a field, then $\delta_r(R) = Soc_r(R) = \begin{bmatrix} 0 & F \\ 0 & F \end{bmatrix}$ since $R/Soc_r(R)$ has zero Jacobson radical, but $\begin{bmatrix} \delta_r(S) & M \\ 0 & \delta_r(T) \end{bmatrix} = \begin{bmatrix} F & F \\ 0 & F \end{bmatrix} = R$. Now we prove the following.

Lemma 5.1 *Let $R = \begin{bmatrix} S & M \\ 0 & T \end{bmatrix}$ where S, T are any ring and M is an (S, T) -bimodule. Then $\delta_r(R) \subseteq \begin{bmatrix} \delta_r(S) & M \\ 0 & \delta_r(T) \end{bmatrix}$.*

Proof Let $r = \begin{bmatrix} s & m \\ 0 & t \end{bmatrix} \in \delta_r(R)$ where $s \in S, t \in T$ and $m \in M$. We claim that $s \in \delta_r(S)$. Let I be a right ideal of S such that $sS + I = S$. It is enough to show that I is a direct summand of S by [21, Theorem 1.6]. Since $rR + \begin{bmatrix} I & M \\ 0 & T \end{bmatrix} = R$ and $r \in \delta_r(R)$, we have that $\begin{bmatrix} I & M \\ 0 & T \end{bmatrix}$ is a direct summand of R and so is generated by an idempotent $e \in R$. Let $e = \begin{bmatrix} g & n \\ 0 & f \end{bmatrix}$ where $g \in S, f \in T$ and $n \in M$. Then g is an idempotent in S and we see that $I = gS$, and hence I is a direct summand of S , as desired. By a similar argument we see that $t \in \delta_r(T)$. Hence, the proof is completed. \square

According to [8, Proposition 6.3], $R = \begin{bmatrix} S & M \\ 0 & T \end{bmatrix}$ is clean if and only if S and T are clean. This result also holds for J -clean ring.

Proposition 5.2 *Let $R = \begin{bmatrix} S & M \\ 0 & T \end{bmatrix}$. Then R is J -clean if and only if S and T are J -clean.*

Proof Since S and T are factor rings of R , the necessity is obvious. Now assume that S and T are J -clean. Let $r = \begin{bmatrix} s & m \\ 0 & t \end{bmatrix} \in R$ where $s \in S, t \in T$ and $m \in M$. Then $s = e + w$ where $e^2 = e \in S$ and $w \in J(S)$, and $t = f + v$ where $f^2 = f \in T$ and $v \in J(T)$. This gives that $\begin{bmatrix} s & m \\ 0 & t \end{bmatrix} = \begin{bmatrix} e & 0 \\ 0 & f \end{bmatrix} + \begin{bmatrix} w & m \\ 0 & v \end{bmatrix}$ where $\begin{bmatrix} e & 0 \\ 0 & f \end{bmatrix}$ is an idempotent in R and $\begin{bmatrix} w & m \\ 0 & v \end{bmatrix} \in J(R)$. Hence, R is J -clean. \square

If S and T are local rings with nonzero maximal left ideal, then $J(S) = \delta_r(S)$ and $J(T) = \delta_r(T)$. By Lemma 5.1, one can thus deduce that $J(R) = \delta_r(R)$. Hence, the following corollary is immediate from Proposition 5.2.

Corollary 5.3 *Let $R = \begin{bmatrix} S & M \\ 0 & T \end{bmatrix}$ where S and T are local rings with nonzero maximal left ideals. Then R is δ_r -clean if and only if S and T are δ_r -clean.*

If $R = \begin{bmatrix} \mathbb{Z}_3 & \mathbb{Z}_3 \\ 0 & \mathbb{Z}_3 \end{bmatrix}$, then \mathbb{Z}_3 is a δ_r -clean ring, but R is not δ_r -clean since no quotient of it is Boolean.

Acknowledgement

The first author thanks the Scientific and Technological Research Council of Turkey (TÜBİTAK) for grant support.

References

- [1] Anderson, F.W., Fuller, K.R: Rings and Categories of Modules. New York. Springer-Verlag 1974.
- [2] Baccella, G.: Exchange property and the natural preorder between simple modules over semi-Artinian rings. *J. Algebra* 253, 133–166 (2002).
- [3] Camillo, V.P., Khurana, D., Lam, T.Y., Nicholson, W.K., Zhou, Y.: Continuous modules are clean. *J. Algebra* 304, 94–111 (2006).
- [4] Camillo, V.P., Yu, H.P.: Exchange rings, units and idempotents. *Comm. Algebra* 22, 4737–4749 (1994).
- [5] Camillo, V.P., Yu, H.P.: Stable range one for rings with many idempotents. *Trans. Amer. Math. Soc.* 347, 3141–3147 (1995).
- [6] Chen, H.: On strongly J-clean rings. *Comm. Algebra* 38, 3790–3804 (2010).
- [7] Chen, J., Wang, Z., Zhou, Y.: Rings in which elements are uniquely the sum of an idempotent and a unit that commute. *J. Pure Appl. Algebra* 213, 215–223 (2009).
- [8] Haghany, A., Varadarajan, K.: Study of formal triangular matrix rings. *Comm. Algebra* 27, 5507–5525 (1999).
- [9] Han, J., Nicholson, W.K.: Extensions of clean rings. *Comm. Algebra* 29, 2589–2595 (2001).
- [10] Handelman, D.: Perspectivity and cancellation in regular rings. *J. Algebra* 48, 1–16 (1977).
- [11] Khurana, D., Lam, T.Y.: Rings with internal cancellation. *J. Algebra* 284, 203–235 (2005).
- [12] Lee, T.K., Yi, Z., Zhou, Y.: An example of Bergman’s and the extension problem for clean rings. *Comm. Algebra* 36, 1413–1418 (2008).
- [13] Lee, T.K., Zhou, Y.: A class of exchange rings. *Glasgow Math. J.* 50, 509–522 (2008).
- [14] Nicholson, W.K.: Lifting idempotents and exchange rings. *Trans. Amer. Math. Soc.* 229, 269–278 (1977).
- [15] Nicholson, W. K., Varadarajan, K., Zhou, Y.: Clean endomorphism rings. *Arch. Math.* 83, 340–343 (2004).
- [16] Nicholson, W.K., Yousif, M.F.: Weakly continuous and C2 conditions. *Comm. Algebra* 29, 2429–2446 (2001).
- [17] Nicholson, W.K., Zhou, Y.: Rings in which elements are uniquely the sum of an idempotent and a unit. *Glasgow Math. J.* 46, 227–236 (2004).
- [18] Özcan, A.Ç., Aydoğdu, P.: A generalization of semiregular and almost principally injective rings. *Algebra Coll.* 17, 905–916 (2010).
- [19] Yousif, M.F., Zhou, Y.: Semiregular, semiperfect and perfect rings relative to an ideal. *Rocky Mountain J. Math.* 32, 1651–1671 (2002).
- [20] Yu, H.P.: On quasi-duo rings. *Glasgow Math. J.* 37, 21–31 (1995).
- [21] Zhou, Y.: Generalizations of perfect, semiperfect and semiregular rings. *Algebra Colloq.* 7, 305–318 (2000).