

1-1-2009

Antioxidant and anticholinesterase constituents of *Salvia pocalata*

UFUK KOLAK

IŞIL HACİBEKİROĞLU

MEHMET ÖZTÜRK

FEVZİ ÖZGÖKÇE

GÜLAÇTI TOPÇU

See next page for additional authors

Follow this and additional works at: <https://journals.tubitak.gov.tr/chem>

 Part of the [Chemistry Commons](#)

Recommended Citation

KOLAK, UFUK; HACİBEKİROĞLU, IŞIL; ÖZTÜRK, MEHMET; ÖZGÖKÇE, FEVZİ; TOPÇU, GÜLAÇTI; and ULUBELEN, AYHAN (2009) "Antioxidant and anticholinesterase constituents of *Salvia pocalata*," *Turkish Journal of Chemistry*. Vol. 33: No. 6, Article 10. <https://doi.org/10.3906/kim-0902-15>
Available at: <https://journals.tubitak.gov.tr/chem/vol33/iss6/10>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Chemistry by an authorized editor of TÜBİTAK Academic Journals. For more information, please contact academic.publications@tubitak.gov.tr.

Antioxidant and anticholinesterase constituents of *Salvia pocalata*

Authors

UFUK KOLAK, IŞIL HACİBEKİROĞLU, MEHMET ÖZTÜRK, FEVZİ ÖZGÖKÇE, GÜLAÇTI TOPÇU, and AYHAN ULUBELEN

Antioxidant and anticholinesterase constituents of *Salvia poculata*

Ufuk KOLAK^{1,*}, Işıl HACİBEKİROĞLU¹, Mehmet ÖZTÜRK²
Fevzi ÖZGÖKÇE³, Gülaçtı TOPÇU⁴, Ayhan ULUBELEN¹

¹Department of General and Analytical Chemistry, Faculty of Pharmacy, İstanbul University
34116, İstanbul-TURKEY
e-mail: ufukkolak@yahoo.com

²Department of Chemistry, Faculty of Arts and Sciences, Muğla University 48121, Muğla- TURKEY

³Department of Biology, Faculty of Science and Letters, Yüzüncü Yıl University
65080, Van-TURKEY

⁴Department of Chemistry, Faculty of Science and Letters, İstanbul Technical University
34469 İstanbul-TURKEY

Received 13.02.2009

Two triterpenoids, namely $2\alpha,3\alpha$ -dihydroxy-24-nor-4(23),12-oleanadien-28-oic acid (**1**) and ursolic acid (**2**); 4 flavonoids, namely 5-hydroxy-7,4'-dimethoxyflavone (**3**), cirsimaritin (**4**), eupatilin (**5**), and salvigenin (**6**); a diterpenoid, namely sclareol (**7**); and a steroid, namely β -sitosterol (**8**), were isolated from the aerial parts of *Salvia poculata* Nab., a Turkish endemic *Salvia* species. Total phenolic and flavonoid contents of the crude extract were determined as pyrocatechol and quercetin equivalents, respectively. The antioxidant activity of the crude extract and the isolated compounds (**2-8**) was established using β -carotene bleaching, superoxide anion radical, and ABTS cation radical scavenging activity. In addition, the anticholinesterase activity of the crude extract and the isolated compounds (**2-8**) against the enzymes acetylcholinesterase (AChE) and butyrylcholinesterase (BChE) was determined. The phytochemistry and antioxidant and anticholinesterase activities of *S. poculata* were investigated for the first time in this study. The crude extract of *S. poculata* exhibited a significant antioxidant effect as well as butyrylcholinesterase inhibitory activity. Ursolic acid (**2**) and cirsimaritin (**4**) possessed high butyrylcholinesterase inhibitory activity.

Key Words: *Salvia poculata*; Lamiaceae; antioxidant activity; anticholinesterase activity.

*Corresponding author

Introduction

Free radicals such as superoxide anion, and hydroxyl and peroxy radicals, which are produced in biological systems and foods, are responsible for oxidation of cell lipids and DNA damage, and they may cause serious diseases (e.g. cancer, coronary arteriosclerosis, diabetes mellitus).¹ Dietary antioxidants may be effective in prevention of oxidative damage. Many scientists have focussed on medicinal and edible plants to discover natural antioxidants since some synthetic antioxidants have toxic effects. In addition, natural antioxidants may have an important role in protecting human health. Studies performed to find natural antioxidants indicated that many *Salvia* species and some of their constituents have shown significant antioxidant activity. Some *Salvia* species have been used commercially in the food industry to prevent or delay spoilage of foods.²

Alzheimer's disease (AD), which is the most common form of dementia, is frequently seen among elderly people all around the world. Some synthetic acetylcholinesterase inhibitors such as tacrine and donepezil have been used for the treatment of AD but they have several adverse effects.³ As a result of the search to find natural acetylcholinesterase inhibitors from plants, more than 35 alkaloids (e.g. physostigmine and galantamine) were found to be active. In addition, some terpenoids, glycosides, and coumarins exhibited similar effects.³ *Salvia* is one of the most investigated genera, some of its species such as *S. multiorrhiza*, *S. officinalis* L. and *S. lavandulaefolia* Vahl. have been used traditionally for dementia therapy.^{4,5} Since investigations on AD suggest that antioxidants may retard its progression,⁴ both antioxidant and anticholinesterase activity tests were carried out on the crude extract and constituents of *S. pocolata* in this study.

Salvia L., with 900 species, is the largest genus of the family Lamiaceae and is widespread throughout the world.⁶ Its species have been widely used in traditional medicine all around the world since ancient times. They have various activities such as anti-inflammatory, analgesic, antipyretic, cardioactive, antifungal, antituberculosis, antitumor, and antioxidant.² In addition, they have been used for the treatment of some cases of dementia.⁷ *Salvia* species have been consumed for different purposes such as tea, spices, and flavouring agents in perfumery and cosmetics; they are among the economically important species of the family Lamiaceae.⁸ There are about 90 *Salvia* species growing naturally in Turkey, half of which are endemic.⁹ In Turkish folk medicine, they have been used as antiseptic, antibacterial, diuretic, haemostatic, spasmolytic, and carminative agents.¹⁰ Since 1968, the phytochemistry and biological activity of about 60 *Salvia* species have been investigated by our research group. These studies indicated that the roots of Turkish *Salvia* species are rich in diterpenoids, especially abietane-type, whereas the aerial parts contain essential oils, flavonoids, and triterpenoids. The main constituents of *Salvia* species possess various biological activities.¹¹

In a continuation of our studies on Turkish *Salvia* species we report herein the constituents and the biological activity of the aerial parts of *S. pocolata* Nab. for the first time. The fatty acid composition of *S. pocolata* seed oil was determined.¹² In this study, 8 known compounds, namely 2 α ,3 α -dihydroxy-24-nor-4(23),12-oleanadien-28-oic acid (**1**),¹³ ursolic acid (**2**),¹⁴ 5-hydroxy-7,4'-dimethoxyflavone (**3**),¹⁵ cirsimaritin (**4**),¹⁶ eupatilin (**5**),¹⁷ salvigenin (**6**),¹⁸ sclareol (**7**),¹⁹ and β -sitosterol (**8**),²⁰ were isolated from the methanol extract of *S. pocolata* (Figure 1). The antioxidant activity of the crude extract and the isolated compounds (**2-8**) was determined using 3 different methods: β -carotene bleaching, and superoxide anion radical and ABTS cation radical scavenging activity assays. In addition, the acetyl- and butyryl-cholinesterase inhibitory activity of the crude extract and the isolated compounds (**2-8**) was established using the Ellman method.




Figure 1. Chemical formulae of compounds 1-8.

Experimental

General Experimental Procedures. The UV spectra (λ_{\max}) were recorded on a Shimadzu UV-1601 in MeOH, IR spectra (ν_{\max}) on a Perkin-Elmer One B in CHCl_3 , NMR spectra on a Varian UNITY INOVA spectrometer operating at 500 MHz for ^1H -NMR and 125 MHz for ^{13}C -NMR (TMS as an internal standard) including APT, HMQC, and HRESI-MS spectra on Bruker Microsoft Q spectrometer. Melting points were recorded on a Kofler apparatus (Reichert) and are uncorrected.

Plant Material. The aerial parts of *S. poculata* Nab. were collected from Eastern Turkey (Alacabük Dagı-Bitlis) at 2250 m altitude in July 2005, and identified by Dr. Fevzi Özgökçe. A voucher specimen was deposited in the Herbarium of Yüzüncü Yıl University (VANF 12938).

Extraction, Isolation, and Identification: The dried and powdered aerial parts of *S. poculata* (350 g) were macerated with 1.5 L of methanol at room temperature (25 °C) 3 times (24 h \times 3). After filtration, the solvent was evaporated to dryness in vacuo. The crude extract (10.6 g) was fractionated on a silica gel column (2.5 \times 100 cm). The column was eluted with petroleum ether (40-60°) (5 \times 150 mL), and a gradient

of dichloromethane was added in 10 mL increments into 100 mL petroleum ether until reaching 100%, thus 10 × 100 mL were used, followed by methanol in 10 mL increments up to 100% (10 × 100 mL). The similar fractions were combined with TLC control and then further subjected to preparative thin layer chromatography to yield compounds **1-8** using the following solvent systems: from Frac. 13-15 β -sitosterol (22 mg) (toluene/diethylether, 7:1) and 5-hydroxy-7,4'-dimethoxyflavone (8 mg) (toluene/ diethylether, 4:1), from Frac. 18-19 salvigenin (6.5 mg) (toluene/diethylether, 7:1), from Frac. 20-25 ursolic acid (13 mg) (toluene/diethylether, 2:1), from Frac. 26-28 sclareol (12 mg) (toluene/diethylether, 2:1), and eupatilin (9 mg) (toluene/chloroform/acetone, 1:9:1), from Frac. 40-45 cirsimaritin (3 mg) (chloroform/ acetone, 9:0.5), and $2\alpha,3\alpha$ -dihydroxy-24-nor-4(23),12-oleanadien-28-oic acid (2 mg) (chloroform/acetone, 8:1).

By correlating our spectral data (UV, IR, 1D- and 2D- NMR, mass) with those of standard samples, compounds **1-8** were identified as $2\alpha,3\alpha$ -dihydroxy-24-nor-4(23),12-oleanadien-28-oic acid (**1**), ursolic acid (**2**), 5-hydroxy-7,4'-dimethoxyflavone (**3**), cirsimaritin (**4**), eupatilin (**5**), salvigenin (**6**), sclareol (**7**), and β -sitosterol (**8**).

$2\alpha,3\alpha$ -Dihydroxy-24-nor-4(23),12-oleanadien-28-oic acid (1): UV (MeOH) λ_{\max} : 207 nm; IR (CHCl₃) ν_{\max} : 3437, 3080, 2950, 1695, 1650, 1462, 1382, 1053, 895, 755 cm⁻¹; ¹H-NMR (500 MHz, CDCl₃) δ : 5.26 (1H, *t*, *J* = 3.6 Hz, H-12), 5.00 (1H, *s*, H-23_b), 4.68 (1H, *d*, *J* = 1.45 Hz, H-23_a), 4.17 (1H, *d*, *J* = 3.66 Hz, H-3 β), 3.72 (1H, *ddd*, *J* = 11.5; 4.76; 3.7 Hz, H-2 β), 2.78 (1H, *dd*, *J* = 4.1; 13.2 Hz, H-18 β), 2.06 (1H, *br dd*, *J* = 2.0; 11.9 Hz, H-5 α), 1.10 (3H, *s*, Me-27), 0.87 (3H, *s*, Me-30), 0.84 (3H, *s*, Me-29), 0.78 (3H, *s*, Me-26), 0.68 (3H, *s*, Me-25); ¹³C-NMR (125 MHz, CDCl₃) δ : 178.52 (*s*, C-28), 148.96 (*s*, C-4), 142.89 (*s*, C-13), 121.67 (*d*, C-12), 110.69 (*t*, C-23), 74.71 (*d*, C-3), 68.15 (*d*, C-2), 47.16 (*s*, C-17), 45.00 (*t*, C-19), 45.10 (*d*, C-9), 44.84 (*d*, C-5), 43.61 (*t*, C-1), 41.87 (*d*, C-18), 40.36 (*s*, C-14), 40.36 (*s*, C-8), 38.00 (*s*, C-10), 32.82 (*t*, C-21), 32.04 (*q*, C-29), 31.41 (*t*, C-22), 28.68 (*t*, C-7), 30.07 (*s*, C-20), 26.57 (*t*, C-15), 24.97 (*q*, C-27), 23.20 (*t*, C-11), 22.56 (*q*, C-30), 22.05 (*t*, C-16), 19.25 (*t*, C-6), 16.14 (*q*, C-26), 13.08 (*q*, C-25). HRESI-MS: *m/z* 456.3229 (calcd for C₂₉H₄₄O₄, 456.3239).

Ursolic acid (2): UV (MeOH) λ_{\max} : 207 nm; IR (CHCl₃) ν_{\max} : 3360, 2940, 2860, 1695, 1500, 1370, 1270, 1185, 1030, 995, 660 cm⁻¹; ¹H-NMR (500 MHz, CDCl₃) δ : 5.24 (1H, *t*, *J* = 4.0 Hz, H-12), 3.23 (1H, *dd*, *J* = 5.0; 10.0 Hz, H-3 α), 1.24 (3H, *s*, Me-27), 1.05 (3H, *s*, Me-26), 0.97 (6H, *d*, *J* = 7.0 Hz, Me-29 and Me-30), 0.90 (3H, *s*, Me-25), 0.78 (3H, *s*, Me-24), 0.73 (3H, *s*, Me-23); ¹³C-NMR (125 MHz, CDCl₃) δ : 180.35 (*s*, C-28), 140.36 (*s*, C-13), 123.63 (*d*, C-12), 78.24 (*d*, C-3), 55.10 (*d*, C-5), 52.62 (*d*, C-9), 47.50 (*s*, C-17), 46.50 (*d*, C-18), 42.28 (*s*, C-14), 40.11 (*s*, C-8), 40.01 (*d*, C-19), 39.01 (*s*, C-10), 38.82 (*d*, C-20), 38.42 (*t*, C-1), 38.13 (*s*, C-4), 32.89 (*t*, C-21), 31.73 (*t*, C-22), 30.35 (*t*, C-7), 28.87 (*t*, C-15), 28.12 (*q*, C-23), 25.92 (*q*, C-27), 24.83 (*t*, C-2), 23.94 (*t*, C-16), 23.40 (*t*, C-11), 21.22 (*q*, C-30), 18.15 (*t*, C-6), 17.65 (*q*, C-24), 17.23 (*q*, C-29), 15.98 (*q*, C-26), 15.20 (*q*, C-25). HRESI-MS: *m/z* 456.3597 (calcd for C₃₀H₄₈O₃, 456.3603).

5-Hydroxy-7,4'-dimethoxyflavone (3): UV (MeOH) λ_{\max} : 268, 328; MeOH+ NaOMe: 288, 340; MeOH + AlCl₃: 277, 380; MeOH + AlCl₃ + HCl: 277, 380; MeOH + NaOAc: 268, 330; MeOH + NaOAc + H₃BO₃: 268, 332; ¹H-NMR (500 MHz, CDCl₃) δ : 12.80 (1H, *br s*, 5-OH), 7.85 (2H, *dd*, *J* = 1.95; 6.83 Hz, H-2' and H-6'), 7.02 (2H, *dd*, *J* = 1.95; 6.83 Hz, H-3' and H-5'), 6.58 (1H, *s*, H-3), 6.48 (1H, *d*, *J* = 2.44 Hz, H-8), 6.37 (1H, *d*, *J* = 2.44 Hz, H-6), 3.89 (3H, *s*, OMe), 3.88 (3H, *s*, OMe); ¹³C-NMR (125 MHz, CDCl₃) δ : 182.41 (*s*, C-4), 164.30 (*s*, C-2), 156.41 (*s*, C-9), 154.61 (*s*, C-4'), 154.02 (*s*, C-5), 148.14 (*s*, C-7), 124.03 (*s*, C-1'), 120.32 (*d*, C-2' and C-6'), 111.54 (*d*, C-3' and C-5'), 103.42 (*s*, C-10), 103.34 (*d*, C-3), 98.06 (*d*, C-6),

93.51 (*d*, C-8), 61.51 (*q*, OMe), 56.15 (*q*, OMe). HRESI-MS: m/z 298.0837 (calcd for C₁₇H₁₄O₅, 298.0841).

Cirsimaritin (5,4'-dihydroxy-6,7-dimethoxyflavone) (4): UV (MeOH) λ_{\max} : 274, 333; MeOH+ NaOMe: 272, 290 (*sh*), 383; MeOH + AlCl₃: 258 (*sh*), 296, 362; MeOH + AlCl₃ + HCl: 262 (*sh*), 299, 355; MeOH + NaOAc: 274, 339, 390; MeOH + NaOAc + H₃BO₃: 274, 335; ¹H-NMR (500 MHz, CDCl₃) δ : 12.70 (1H, *br s*, 5-OH), 7.74 (2H, *d*, $J = 8.7$ Hz, H-2' and H-6'), 6.90 (2H, *d*, $J = 8.7$ Hz, H-3' and H-5'), 6.52 (1H, *s*, H-3), 6.48 (1H, *s*, H-8), 3.89 (3H, *s*, OMe), 3.85 (3H, *s*, OMe); ¹³C-NMR (125 MHz, CDCl₃) δ : 181.52 (*s*, C-4), 164.15 (*s*, C-2), 157.16 (*s*, C-5), 152.88 (*s*, C-4'), 150.22 (*s*, C-7), 149.97 (*s*, C-9), 132.15 (*s*, C-6), 124.38 (*s*, C-1'), 121.44 (*d*, C-2' and C-6'), 115.43 (*d*, C-3' and C-5'), 104.13 (*s*, C-10), 103.32 (*d*, C-3), 91.42 (*d*, C-8), 62.15 (*q*, OMe), 61.22 (*q*, OMe). HRESI-MS: m/z 314.0787 (calcd for C₁₇H₁₄O₆, 314.0790).

Eupatilin (5,7-dihydroxy-6, 3',4'-trimethoxyflavone) (5): mp = 230-232 °C; UV (MeOH) λ_{\max} : 243, 275, 340; MeOH+ NaOMe: 242, 270, 377; MeOH + AlCl₃: 262, 291 (*sh*), 364; MeOH + AlCl₃ + HCl: 261, 295, 360; MeOH + NaOAc: 235, 275, 340; MeOH + NaOAc + H₃BO₃: 236, 275, 341; ¹H-NMR (500 MHz, CDCl₃) δ : 12.82 (1H, *br s*, 5-OH), 7.42 (1H, *d*, $J = 2.0$ Hz, H-2') 7.37 (1H, *dd*, $J = 2.0$; 8.79 Hz, H-6'), 6.90 (1H, *d*, $J = 8.79$ Hz, H-5'), 6.52 (1H, *s*, H-3), 6.48 (1H, *s*, H-8), 3.92 (3H, *s*, OMe), 3.92 (3H, *s*, OMe), 3.86 (3H, *s*, OMe); ¹³C-NMR (125 MHz, CDCl₃) δ : 182.01 (*s*, C-4), 162.18 (*s*, C-2), 158.46 (*s*, C-5), 156.00 (*s*, C-4'), 152.87 (*s*, C-3'), 151.26 (*s*, C-9), 149.07 (*s*, C-7), 133.11 (*s*, C-6), 123.41 (*s*, C-1'), 120.13 (*d*, C-6'), 111.62 (*d*, C-5'), 108.22 (*d*, C-2'), 105.37 (*s*, C-10), 103.46 (*d*, C-3), 92.00 (*d*, C-8), 61.42 (*q*, OMe), 56.18 (*q*, OMe). 56.01 (*q*, OMe). HRESI-MS: m/z 344.0891 (calcd for C₁₈H₁₆O₇, 344.0895).

Salvigenin (6): mp = 188 °C, UV (MeOH) λ_{\max} : 277, 330; MeOH+ NaOMe: 295, 332 (*sh*), 375; MeOH + AlCl₃: 235, 302 (*sh*), 360; MeOH + AlCl₃ + HCl: 235, 262, 301 (*sh*), 350; MeOH + NaOAc: 277, 329, 376; MeOH + NaOAc + H₃BO₃: 276, 329; ¹H-NMR (500 MHz, CDCl₃) δ : 12.76 (1H, *br s*, 5-OH), 7.86 (2H, *d*, $J = 10.0$ Hz, H-2' and H-6'), 7.02 (2H, *d*, $J = 10.0$ Hz, H-3' and H-5'), 6.59 (1H, *s*, H-3), 6.55 (1H, *s*, H-8), 3.90 (3H, *s*, OMe), 3.93 (3H, *s*, OMe), 3.97 (3H, *s*, OMe); ¹³C-NMR (125 MHz, CDCl₃) δ : 181.52 (*s*, C-4), 163.12 (*s*, C-2), 158.13 (*s*, C-5), 154.20 (*s*, C-4'), 152.00 (*s*, C-7), 151.74 (*s*, C-9), 132.06 (*s*, C-6), 124.03 (*s*, C-1'), 120.32 (*d*, C-2' and C-6'), 111.54 (*d*, C-3' and C-5'), 104.92 (*s*, C-10), 102.93 (*d*, C-3), 91.05 (*d*, C-8), 62.03 (*q*, OMe), 61.54 (*q*, OMe), 56.13 (*q*, OMe). HRESI-MS: m/z 328.0940 (calcd for C₁₈H₁₆O₆, 328.0946).

Sclareol (7): UV (MeOH) λ_{\max} : 217 nm; IR (CHCl₃) ν_{\max} : 3400, 2940, 1465, 1395, 900 cm⁻¹; ¹H-NMR (500 MHz, CDCl₃) δ : 5.94 (1H, *dd*, $J = 10.74$; 17.08 Hz, H-14), 5.21 (1H, *dd*, $J = 1.47$; 17.08 Hz, H-15_a), 5.02 (1H, *dd*, $J = 1.47$; 10.74 Hz, H-15_b), 1.27 (3H, *s*, Me-16), 1.16 (3H, *s*, Me-17), 0.86 (3H, *s*, Me-20), 0.78 (6H, *s*, Me-18 and Me-19); ¹³C-NMR (125 MHz, CDCl₃) δ : 146.23 (*d*, C-14), 111.43 (*t*, C-15), 74.98 (*s*, C-8), 73.84 (*s*, C-13), 61.89 (*d*, C-9), 55.51 (*d*, C-5), 45.24 (*t*, C-12), 44.64 (*t*, C-7), 42.24 (*t*, C-3), 39.96 (*t*, C-1), 39.50 (*s*, C-10), 33.61 (*q*, C-18), 33.46 (*s*, C-4), 27.52 (*q*, C-16), 24.48 (*q*, C-17), 21.71 (*q*, C-19), 20.75 (*t*, C-11), 19.33 (*t*, C-6), 18.66 (*t*, C-2), 15.57 (*q*, C-20). HRESI-MS: m/z 308.2710 (calcd for C₂₀H₃₆O₂, 308.2715).

β -Sitosterol (8): mp = 138-139 °C, UV (MeOH) λ_{\max} : 205 nm; IR (CHCl₃) ν_{\max} : 3445, 3305, 2925, 2860, 1643, 1375, 1063, 955, 835 cm⁻¹; ¹H-NMR (500 MHz, CDCl₃) δ : 5.35 (1H, *m*, H-6), 3.52 (1H, *m*, H-3 α), 1.01 (3H, *s*, Me-19), 0.92 (3H, *d*, $J = 6.4$ Hz, Me-21), 0.85 (3H, *t*, $J = 7.8$ Hz, Me-29), 0.83 (3H, *d*, $J = 6.8$ Hz, Me-26), 0.81 (3H, *d*, $J = 6.9$ Hz, Me-27), 0.69 (3H, *s*, Me-18); ¹³C-NMR (125 MHz, CDCl₃) δ : 140.71 (*s*, C-5), 121.63 (*d*, C-6), 71.73 (*d*, C-3), 56.75 (*d*, C-14), 56.02 (*d*, C-17), 51.13 (*d*, C-9), 45.82 (*d*, C-24), 42.37

(*s*, C-13), 42.20 (*t*, C-4), 39.79 (*t*, C-12), 37.33 (*t*, C-1), 36.43 (*s*, C-10), 36.07 (*d*, C-20), 33.95 (*t*, C-22), 31.96 (*t*, C-7), 31.81 (*d*, C-8), 31.63 (*t*, C-2), 29.15 (*d*, C-25), 28.25 (*t*, C-16), 26.10 (*t*, C-23), 24.25 (*t*, C-15), 23.13 (*t*, C-28), 21.09 (*t*, C-11), 19.77 (*q*, C-26), 19.46 (*q*, C-19), 19.21 (*q*, C-27), 18.68 (*q*, C-21), 11.84 (*q*, C-18), 11.04 (*q*, C-29). HRESI-MS: m/z 414.3857 (calcd for C₂₉H₅₀O, 414.3861).

Antioxidant activity

Chemicals: Methanol, chloroform, pyrocatechol, quercetin, sodium carbonate, aluminium nitrate, potassium acetate, potassium persulfate, sodium phosphate, silica gel for column chromatography (1.07734) and silica gel 60 F₂₅₄ TLC plates (1.05554) were obtained from E. Merck (Darmstadt, Germany). Folin-Ciocalteu's reagent, β -carotene, linoleic acid, polyoxyethylene sorbitan monopalmitate, butylated hydroxytoluene, butylated hydroxyanisole, trolox, (+)-catechin, nicotinamideadeninedinucleotide, acetyl-cholinesterase, butyrylcholinesterase, 5,5'-dithiobis (2-nitrobenzoic) acid, acetylthiocholine iodide, butyrylthiocholine iodide, and galanthamine were obtained from Sigma Chemical Co. (Sigma-Aldrich GmbH, Sternheim, Germany). 2,2'-Azinobis (3-ethylbenzothiazoline-6-sulfonic acid) diammonium salt, Tris-HCl, nitrotetrazoliumbluechloride, and *N*-methyl-phenazoniummethylsulphate were obtained from Fluka Chemie (Fluka Chemie GmbH, Sternheim, Germany).

Determination of Total Phenolic Content: The concentrations of the phenolic content in all samples were expressed as micrograms of pyrocatechol equivalents (PEs). The phenolic content was determined as described in the literature.²¹

Determination of Total Flavonoid Content: The measurement of the flavonoid concentration was based on the method described by Moreno et al. with a slight modification and the results were expressed as quercetin equivalents.²²

Determination of the Antioxidant Activity with the β -Carotene Bleaching Method: The antioxidant activity was established using the β -carotene–linoleic acid test system.²³

Superoxide Anion Radical Scavenging Activity: The measurement of the superoxide anion radical scavenging activity was based on the method described by Liu et al. with slight modification.²⁴

ABTS Cation Radical Decolorization Assay: The ABTS⁺ scavenging activity was determined according to the method of Re et al. with slight modifications.²⁵

Anticholinesterase Activity: Acetyl- and butyryl-cholinesterase inhibitory activities were established by slightly modifying the spectrophotometric method developed by Ellman et al.²⁶

Statistical analysis: The results were mean \pm SD of 3 parallel measurements. Analysis of variance was performed using ANOVA procedures. Significant differences between means were determined by Student's *t*-test, and *p* values < 0.05 were regarded as significant.

Results and discussion

Two triterpenoids, namely 2 α ,3 α -dihydroxy-24-nor-4(23),12-oleanadien-28-oic acid (**1**) and ursolic acid (**2**); 4 flavonoids, namely 5-hydroxy-7,4'-dimethoxyflavone (**3**), cirsimaritin (**4**), eupatilin (**5**), and salvigenin (**6**); a diterpenoid, namely sclareol (**7**); and a steroid, namely β -sitosterol (**8**), were obtained from the aerial parts of

the endemic plant *S. pocolata*. A rather rare nortriterpenoid $2\alpha,3\alpha$ -dihydroxy-24-nor-4(23),12-oleanadien-28-oic acid (**1**) was obtained for the first time from a Turkish *Salvia* species, previously isolated from *S. carduacea*.¹³ The structure determination of the compounds (**1-8**) was established using spectral methods (UV, IR, 1D- and 2D-NMR, mass spectrometry) and their spectral data were compared to those of standard samples.

In this study, the antioxidant activity of eupatilin (**5**) and sclareol (**7**) was determined for the first time. Although some antioxidant activity tests were performed for ursolic acid (**2**),^{27,28} 5-hydroxy-7,4'-dimethoxyflavone (**3**),²⁹ cirsimaritin (**4**),³⁰⁻³² salvigenin (**6**),³⁰ and β -sitosterol (**8**),^{33,34} we tested their antioxidant activity using β -carotene bleaching, and superoxide anion radical and ABTS cation radical scavenging activity assays.

As shown in Table 1, the crude extract is rich in phenolic compounds. The crude extract and β -sitosterol (**8**), which is a common steroid in *Salvia* species, showed significant inhibition of lipid peroxidation by the β -carotene bleaching method (Figure 2). Sclareol (**7**), a labdane diterpenoid, exhibited the highest activity among the tested compounds as a superoxide anion radical scavenger, while the crude extract was found to be moderately active in the ABTS cation radical scavenging activity test (Table 2). Since cirsimaritin (**4**) and salvigenin (**6**) did not exhibit antioxidant activity their results are not shown in Figure 2 and Table 2.

Table 1. Total phenolic and flavonoid contents of the crude extract of *S. pocolata*.^a

Sample	Phenolic content (μg PEs/mg extract) ^b	Flavonoid content (μg QEs/mg extract) ^c
ESP	85.56 ± 1.60	35.28 ± 0.13

^aValues expressed are means \pm S.D. of 3 parallel measurements ($p < 0.05$)

^bPEs, pyrocatechol equivalents

^cQEs, quercetin equivalents

Table 2. O_2^- and ABTS^+ scavenging activities of compounds **2-3**, **5-8**.^a

Samples	O_2^- assay		ABTS^+ assay	
	$\text{IC}_{50}(\mu\text{g}/\text{mL})$	r^2	$\text{IC}_{50}(\mu\text{g}/\text{mL})$	r^2
ESP	57.99 ± 1.11	0.9804	17.99 ± 0.11	0.9944
2	94.43 ± 0.13	0.9930	> 100	0.9911
3	71.65 ± 1.95	0.9828	44.80 ± 1.56	0.9961
5	59.14 ± 0.98	0.9961	22.34 ± 0.89	0.9969
6	78.46 ± 1.55	0.9987	78.68 ± 1.32	0.9908
7	48.74 ± 0.25	0.9994	> 100	0.9970
8	> 100	0.9904	> 100	0.9818
BHA ^b	33.71 ± 0.11	0.9924	2.16 ± 0.09	0.9889
(+)-Catechin ^b	44.12 ± 0.09	0.9995	1.16 ± 0.05	0.9984

^aValues expressed are means \pm S.D. of 3 parallel measurements ($p < 0.05$)

^bReference compounds

Table 3. Anticholinesterase activity of the crude extract and compounds **2-8**.^a

Samples	Inhibition % against AChE				Inhibition % against BChE			
	25 μ M	50 μ M	100 μ M	200 μ M	25 μ M	50 μ M	100 μ M	200 μ M
ESP*	1.78 \pm 0.87	3.78 \pm 0.58	5.81 \pm 1.34	8.62 \pm 0.34	32.68 \pm 1.73	39.79 \pm 1.09	49.09 \pm 1.45	55.59 \pm 0.45
2	34.60 \pm 0.39	42.63 \pm 0.75	50.72 \pm 0.53	54.26 \pm 0.21	62.84 \pm 0.62	64.46 \pm 0.90	66.70 \pm 0.89	70.80 \pm 0.17
3	na	na	na	na	na	na	na	na
4	1.78 \pm 0.87	3.78 \pm 0.58	7.35 \pm 0.58	12.35 \pm 0.99	30.59 \pm 0.73	35.69 \pm 1.29	42.31 \pm 0.36	50.31 \pm 0.94
5	na	na	0.94 \pm 0.63	11.00 \pm 0.72	15.22 \pm 1.03	22.12 \pm 1.41	32.22 \pm 0.24	45.27 \pm 0.33
6	na	na	na	na	4.85 \pm 0.54	5.67 \pm 0.36	7.96 \pm 0.67	14.19 \pm 0.51
7	na	na	na	na	0.99 \pm 0.09	1.35 \pm 0.07	7.99 \pm 0.73	26.60 \pm 0.23
8	na	na	na	na	3.83 \pm 1.37	4.28 \pm 0.94	6.80 \pm 0.82	11.04 \pm 0.88
Galantamine ^b	68.36 \pm 1.10	74.38 \pm 0.65	78.59 \pm 0.47	81.41 \pm 0.03	40.59 \pm 0.88	48.73 \pm 0.90	65.02 \pm 0.44	75.54 \pm 1.05

^a Values expressed are means \pm S.D. of t3 parallel measurements (p < 0.05)

^b Standard drug
 * at 200 μ g/mL concentration

na: not active




Figure 2. Inhibition (%) of lipid peroxidation of ESP, compounds (**2-3**, **5**, **7-8**), (+)-catechin, BHT and trolox by the β -carotene bleaching method. Values are mean \pm S.D., $n=3$. $p < 0.05$, significantly different with Student's t test. (ESP: Extract of *S. pocolata*, BHT: butylatedhydroxytoluene).

The anticholinesterase activity against the enzymes acetylcholinesterase (AChE) and butyrylcholinesterase (BChE) of the crude extract and the isolated compounds (**2-8**) was investigated for the first time. While the crude extract was found to be almost inactive in the AChE inhibitory test, it exhibited high inhibition in the BChE inhibitory assay. Ursolic acid (**2**) possessed moderate AChE inhibitory activity, as seen in Table 3, which has been previously investigated by Chung et al.³⁵ However, ursolic acid (**2**) exhibited higher BChE inhibitory activity than the standard test compound galantamine, which is used in the treatment of mild to moderate AD. Flavonoid compounds, cirsimaritin (**4**) and eupatilin (**5**), were almost inactive against AChE, while they showed high BChE inhibition. Other isolated compounds, 5-hydroxy-7,4'-dimethoxyflavone (**3**), salvigenin (**6**), sclareol (**7**) and β -sitosterol (**8**), showed no inhibition against AChE and BChE.

Conclusion

Although sclareol (**7**) and β -sitosterol (**8**) showed high total antioxidant activity in the β -carotene bleaching method, they exhibited no anticholinesterase activity. Among the isolated compounds, although triterpenoid ursolic acid (**2**) showed the highest anticholinesterase activity, especially against BChE, it exhibited neither lipid inhibition nor radical scavenging activity.

Acknowledgements

This work was supported by the Research Fund of Istanbul University: Project number: BYP-1967 and by The Scientific and Technological Research Council of Turkey (TÜBİTAK-TBAG-107T592). One of us (A.U.) thanks

the Turkish Academy of Sciences (TUBA) for the partial support of this study.

References

1. Reische, D. W.; Lillard, D. A.; Eitenmiller, R. R. *Antioxidants*. In *Food Lipids* (Chemistry, nutrition and biotechnology). Akoh, C. C.; Min, D. B., Eds.; pp. 489-516: Marcel Dekker, New York, 2002.
2. Fraga, B. M.; Diaz, C. E.; Guadano, A.; Gonzalez-Coloma, A. *J. Agric. Food Chem.* **2005**, textit53, 5200-5206.
3. Mukherjee, P. K.; Kumar, V.; Mal, M.; Houghton, P. J. *Phytomedicine* **2007**, textit14, 289-300.
4. Howes, M-J. R.; Perry, N. S. L.; Houghton, P. J. *Phytother. Res.* **2003**, textit17, 1-18.
5. Scholey, A. B.; Tildesley, N. T. J.; Ballard, C. G.; Wesnes, K. A.; Tasker, A.; Perry, E. K.; Kennedy, D. O. *Psychopharmacology* **2008**, textit198, 127-139.
6. Topçu, G. *J. Nat. Prod.* **2006**, textit69, 482-487.
7. Savelev, S. U.; Okello, E. J.; Perry, E. K. *Phytother. Res.* **2004**, textit18, 315-324.
8. Delamare, A. P. L.; Moschen-Pistorello, I. T.; Artico, L.; Atti-Serafini, L.; Echeverrigaray, S. *Food Chem.* **2007**, textit100, 603-608.
9. Hedge, I. C. *Salvia* L. In *Flora of Turkey and the East Aegean Islands*; Davis, P.H., Ed.; vol. 7, p. 400-459, Edinburgh University Press, Edinburgh, 1982.
10. Baytop, T. *Therapy with Medicinal Plants in Turkey*, pp. 156-158: Istanbul University Press, İstanbul, 1984.
11. Ulubelen, A.; Topçu, G. Chemical and Biological Investigations of *Salvia* species growing in Turkey. In *Studies in Natural Products Chemistry*; Atta-ur-Rahman, Ed.; vol. 20, pp. 659-718, Elsevier Press, New York, 1998.
12. Gören, A. C.; Kılıç, T.; Dirmenci, T.; Bilsel, G. *Biochem. Syst. Ecol.* **2006**, textit34, 160-164.
13. Ballesta-Acosta, M. C.; Pascual-Villalobos, M. J.; Rodriguez, B. *J. Nat. Prod.* **2002**, textit65, 1513-1515.
14. Ogura, M.; Cordell, G. A.; Farnsworth, N. R. *Lloydia* **1977**, textit40, 157-168.
15. Ahmad, S. *Planta Med.* **1983**, textit 48, 62-63.
16. Mues, R.; Timmermann, B. N.; Ohno, N.; Mabry, T. J. *Phytochemistry* **1979**, textit18, 1379-1383.
17. Appendino, G.; Valle, M. G.; Nano, G. M. *Fitoterapia* **1987**, LVIII, 115-118.
18. Ulubelen, A.; Ozturk, S.; Isildatici, S. *J. Pharm. Sci.* **1968**, textit57, 1037-1038.
19. Ulubelen, A.; Topçu, G.; Eris, C.; Sönmez, U.; Kartal, M.; Kurucu, S.; Bozok-Johansson, C. *Phytochemistry* **1994**, textit36, 971-974.
20. DellaGreca, M. D.; Monaco, P.; Previtiera, L. *J. Nat. Prod.* **1990**, textit53, 1430-1435.
21. Slinkard, K.; Singleton, V. L. *Am. J. Enol. Viticult.* **1977**, textit28, 49-55.
22. Moreno, M. I. N.; Isla, M. I.; Sampietro, A. R.; Vattuone, M. A. *J. Ethnopharmacol.* **2000**, textit71, 109-114.
23. Miller, H. M. *J. Am. Oil Chem. Soc.* **1971**, textit45, 91.
24. Liu, F.; Ooi, V. E. C.; Chang, S. T. *Life Sci.* **1997**, textit60, 763-771.
25. Re, R.; Pellegrini, N.; Proteggente, A.; Pannala, A.; Yang, M.; Rice-Evans, C. *Free Radical Bio. Med.* **1999**, textit26, 1231-1237.

26. Ellman, G. L.; Courtney K. D.; Andres V.; Featherston, R. M. *Biochem. Pharmacol.* **1961**, textit7, 88-95.
27. D'Abrosca, B.; Fiorentino, A.; Monaco, P.; Oriano, P.; Pacifico, S. *Food Chem.* **2006**,textit 98, 285-290.
28. Yin, M-C; Chan, K-C. *J. Agr. Food Chem.* **2007**, textit55, 7177-7181.
29. Martini, N. D.; Katerere, D. R. P.; Eloff, J. N. *J. Ethnopharmacol.* **2004**, textit93, 207-212.
30. Ono, M.; Morinaga, H.; Masuoka, C.; Ikeda, T.; Okawa, M.; Kinjo, J.; Nohara, T. *Chem. Pharm. Bul.* **2005**, textit53, 1175-1177.
31. Zheng, W.; Wang, S. Y. *J. Agr. Food Chem.* **2001**, textit49, 5165-5170.
32. Ibanez, E.; Kubatova, A.; Senorans, F. J.; Caverro, S.; Reglero, G.; Hawthorne, S. B. *J. Agr. Food Chem.* **2003**,textit 51, 375-382.
33. Han, J.; Weng, X.; Bi, K. *Food Chem.* **2008**, textit106, 2-10.
34. Weng, X. C.; Wang, W. *Food Chem.* **2000**, textit71, 489-493.
35. Chung, Y. K.; Heo, H. J.; Kim, E. K.; Kim, H. K.; Huh, T. L.; Lim, Y.; Kim, S. K.; Shin, D. H. *Mol. Cells* **2001**, textit11, 137-143.