

1-1-2013

Quasinormability and diametral dimension

AHMET TOSUN TERZİOĞLU

Follow this and additional works at: <https://dctubitak.researchcommons.org/math>



Part of the [Mathematics Commons](#)

Recommended Citation

TERZİOĞLU, AHMET TOSUN (2013) "Quasinormability and diametral dimension," *Turkish Journal of Mathematics*: Vol. 37: No. 5, Article 11. <https://doi.org/10.3906/mat-1207-26>
Available at: <https://dctubitak.researchcommons.org/math/vol37/iss5/11>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Mathematics by an authorized editor of TÜBİTAK Academic Journals.

Quasinormability and diametral dimension

A. Tosun TERZİOĞLU*

Sabancı University, Faculty of Engineering and Natural Sciences
34956 Orhanlı, Tuzla, İstanbul, Turkey

Received: 18.07.2012 • Accepted: 05.09.2012 • Published Online: 26.08.2013 • Printed: 23.09.2013

Abstract: Two versions of diametral dimension are shown to coincide for quasinormable Fréchet spaces. The diametral dimension is determined by a single bounded subset in certain cases.

Key words and phrases: Diametral dimension, Fréchet spaces, Köthe spaces

1. Introduction

The set $\Delta(E)$ of sequences (ξ_n) such that for each neighborhood U of zero of a locally convex space E there is another such neighborhood with $\lim \xi_n d_n(V, U) = 0$, where $d_n(V, U)$ is the n -th diameter of V with respect to U , is called the *diametral dimension* of E . ([3], [6], [7], [8]). Another version is the set $\Delta_b(E)$ of all sequences (ξ_n) such that for each neighborhood U and each bounded subset B we have $\lim \xi_n d_n(B, U) = 0$. $\Delta_b(E)$ is less frequently used than $\Delta(E)$. We always have $c_0 \subset \Delta(E) \subset \Delta_b(E)$. In [6] Mitiagin claimed that $\Delta(E) = \Delta_b(E)$ holds for every Fréchet space (F -space) E , referring for the proof to a forthcoming joint paper. However, there is an example of a Köthe space $\lambda(A)$, which is a Montel space but fails to be a Schwartz space. In this case we have

$$\Delta(\lambda(A)) = c_0 \subset \ell_\infty \subset \Delta_b(\lambda(A)).$$

On the other hand, if E is a locally convex space with a bounded subset that is not precompact, we have $\Delta(E) = \Delta_b(E) = c_0$.

We recall that a Fréchet-Montel space (FM -space) is a Fréchet-Schwartz space (FS -space) if and only if E is quasinormable [3]. There is an extensive literature concerning quasinormability (cf. [1]). We want to single out a remarkable result of Meise and Vogt [5], which states that an F -space is quasinormable if and only if it is isomorphic to a quotient space of the complete tensor product $\ell^1(I) \tilde{\otimes}_\pi \lambda(A)$, where I is a suitable index set and $\lambda(A)$ a suitable Köthe-Schwartz space.

We recall the definition of the n -th diameter

$$d_n(A, B) = \inf \inf \{ \alpha > 0 : A \subset \alpha B + L \}$$

where A and B are subsets of a locally convex space E with $A \subset \rho B$ for some $\rho > 0$. The second infimum is taken over all subspaces L of E with dimension not exceeding $n \in \mathbb{N}$.

*Correspondence: tosun@sabanciuniv.edu

2010 AMS Mathematics Subject Classification: 46A04, 46A11, 46A45.

Proposition 1 *If E is a quasinormable metrisable space, then $\Delta(E) = \Delta_b(E)$.*

Proof For $(\xi_n) \in \Delta_b(E)$ we find $\delta_n \geq \delta_{n+1} > 0$ with $\lim \xi_n \delta_n = 0$. By quasinormability we can choose a base of absolutely convex, closed neighborhoods $U_1, \supset U_2 \supset \dots$ such that for each k and $\delta_n > 0$ there is a bounded subset $B_{k,n}$ with

$$U_{k+1} \subset B_{k,n} + \delta_n U_k.$$

In particular,

$$U_{n+1} \subset B_{k,n} + \delta_n U_n$$

holds for each $n \geq k$. To see how much of $B_{k,n}$ we need in the above inclusion we observe that for each $x \in U_{n+1}$ there is a $b \in B_{k,n}$ with $\|x - b\|_n \leq \delta_n$. This means if we replace $B_{k,n}$ with

$$B_{k,n} \cap (1 + \delta_0)U_n$$

the above inclusion still holds. Therefore, we will assume without loss of generality that

$$B_{k,n} \subset (1 + \delta_0)U_n$$

in the above inclusion.

This implies that

$$B_k = \bigcup_{n=k}^{\infty} B_{k,n}$$

is a bounded set and therefore for all $n \geq k$ we obtain

$$U_{k+1} \subset B_k + \delta_n U_k.$$

Using the definition of the n -th diameter, from the above inclusion for $n \geq k$ we get

$$d_n(U_{k+1}, U_k) \leq d_n(B_k, U_k) + \delta_n.$$

Hence

$$\lim \xi_n d_n(U_{k+1}, U_k) = 0.$$

□

Our result implies that for an F -space we have $\Delta(E) \neq \Delta_b(E)$ if and only if E is a Montel but not a Schwartz space.

In certain cases it is sufficient to consider a single bounded subset of E to determine $\Delta(E)$. We will call an absolutely convex bounded subset B of an F -space E a *prominent set* if $\lim \xi_n d_n(B, U_k) = 0$ for every k implies $(\xi_n) \in \Delta(E)$. If E has a prominent set B then since

$$\Delta(E) = \{(\xi_n) : \lim \xi_n d_n(B, U_k) = 0, \quad k = 1, 2, \dots\}$$

the diametral dimension as a space of sequences is an F -space itself. For an exponent sequence $0 < \alpha_n \leq \alpha_{n+1} \leq \dots$ with $\lim \alpha_n = \infty$ the unit ball B_1 of ℓ_1 is a prominent set of the finite-type power series space $\Lambda_1(\alpha)$ (cf. for example [6], [8]). We will generalize this result in what follows.

Following [10], we call a Banach space $(\ell, \|\cdot\|)$ of scalar sequences an *admissible space* if $\|e_n\| = 1$ and for $a \in \ell_\infty$, $x \in \ell$ we have $ax = (a_n x_n) \in \ell$ and $\|ax\| \leq \|a\|_\infty \|x\|$. As usual e_n is that sequence with 1 as the n -th term and zero elsewhere. The classical sequence spaces $\ell_p, 1 \leq p \leq \infty$ and c_0 are the most well-known examples of admissible spaces.

Let A be a Köthe set and $\lambda^\ell(A)$ be the space of all sequences $x = (x_n)$ such that $(x_n a_n^k) \in \ell$ for each k . Equipped with the seminorms, $\|x\|_k = \|a^k x\|$, $\lambda^\ell(A)$ is an F -space. $\lambda^{\ell_1}(A)$ is of course the usual Köthe space $\lambda(A)$. In fact the spaces $\lambda^{\ell_p}(A), \lambda^{c_0}(A), 1 \leq p \leq \infty$, are also quite well known.

A Köthe space $\lambda(A)$ is called a *smooth sequence space of finite type* [8] (or a G_1 -space) if $0 < a_{n+1}^k \leq a_n^k$ and for each k there is a j with $(a_n^k / (a_n^j)^2) \in \ell_\infty$.

Proposition 2 *Let $\lambda(A)$ be a G_1 -space and ℓ an admissible space with closed unit ball B_ℓ . Then B_ℓ is a prominent subset of $\lambda^\ell(A)$.*

Proof Let

$$U_k = \{(x_n) \in \lambda^\ell(A) : \|(x_n a_n^k)\| \leq 1\}.$$

By [10], Prop. 1, we have the basic inequality

$$\inf \left\{ \frac{a_i^k}{a_i^j} : i \leq n \right\} \leq d_n(U_j, U_k) \leq \sup \left\{ \frac{a_i^k}{a_i^j} : i \geq n \right\}$$

We note that both sides of this inequality are independent of ℓ . With the same argument we can easily show

$$d_n(B_\ell, U_k) = a_n^k.$$

Now for k given we choose j such that for some $\rho > 0$ we have $a_n^k \leq \rho (a_n^j)^2$ for all $n \in N$. From the above inequality we obtain

$$d_n(U_j, U_k) \leq \rho \sup\{a_i^j : i \geq n\} = \rho a_n^j.$$

Therefore,

$$d_n(U_j, U_k) \leq \rho d_n(B_\ell, U_j).$$

This shows

$$\Delta(\lambda^\ell(A)) = \Delta_b(\lambda^\ell(A)) = \{(\xi_n) : \lim \xi_n d_n(B_\ell, U_j) = 0 \text{ for all } j \in N\}.$$

□

Of course, a finite type power series space $\Lambda_1(\alpha)$ is a G_1 -space. In this special case the closed unit ball B_p of ℓ_p or B_0 of c_0 is a prominent subset of $\Lambda_1^{\ell_p}(\alpha)$, or of $\Lambda_1^{c_0}(\alpha)$.

We will now give a necessary and sufficient condition for a bounded subset to be prominent.

Proposition 3 *Let B be a bounded subset of an F -space E . B is a prominent set if and only if for each k there is a p and $\rho > 0$ such that*

$$d_n(U_p, U_k) \leq \rho d_n(B, U_p)$$

for all $n \in N$.

Proof Sufficiency follows immediately from definitions of $\Delta(E)$ and $\Delta_b(E)$. Let us now assume B is a prominent subset of E . In this case, the diametral dimension is

$$\Delta(E) = \lambda^{co}(B_E)$$

where

$$B_E = \{(d_n(B, U_k)) : k = 1, 2, \dots\}$$

and so $\Delta(E)$ is itself an F -space. On the other hand, from the definition of $\Delta(E)$, for a given k we have

$$\Delta(E) \subset \cup_{p \geq k} \{(\xi_n) : \lim \xi_n d_n(U_p, U_k) = 0\}$$

The right-hand side of the above inclusion is an LB -space and the canonical inclusion map is sequentially closed. Therefore, by the Grothendieck factorization theorem we can find $m > k$ such that

$$\Delta(E) \subset \{(\xi_n) : \lim \xi_n d_n(U_m, U_k) = 0\}.$$

This implies the existence of j and $\rho > 0$ with

$$d_n(U_m, U_k) \leq \rho d_n(B, U_j), \quad n \in N.$$

Finally we choose $p = \max\{m, j\}$. □

Let $\lambda(A)$ now be a *smooth sequence space of infinite type* [8]. This means $0 < a_n^k \leq a_{n+1}^k \leq \dots$ and for each k there is a_j with $((a_n^k)^2/a_n^j) \in \ell_\infty$. To avoid the trivial case $\lambda(A) = \ell_1$ we will also assume $\lim_{n \rightarrow \infty} a_n^k = \infty$ for every k . This implies that $\lambda(A)$ is a Schwartz space, and so

$$\Delta(\lambda(A)) = \Delta_b(\lambda(A))$$

by Proposition 1. Let B be a prominent subset of $\lambda(A)$. Since a bounded set, which contains a prominent subset, is itself prominent, we can assume without loss of generality [2] that

$$B = \{(\xi_n) : \sum |\xi_n| a_n \leq 1\}$$

where (a_n) is some sequence such that for each k there is a $\rho_k > 0$ with $a_n^k \leq \rho_k a_n$ for all $n \in N$. By Prop. 3. for each k there is a $\rho > 0$ and $p \geq k$ with

$$d_n(U_p, U_k) \leq \rho d_n(B, U_p).$$

We choose m so that $((a_n^p)^3/a_n^m) \in \ell_\infty$. By the basic inequality

$$d_n(B, U_p) \leq \inf\{a_i^p/a_i : i \geq n\}$$

but

$$\frac{a_i^p}{a_i} \leq c \frac{a_i^m}{(a_i^p)^2 a_i} \leq \frac{c \rho_m}{(a_i^p)^2}$$

for some constant $c > 0$.

Applying the left-hand side of the basic inequality we have

$$\frac{a_0^k}{a_n^p} \leq d_n(U_p, U_k).$$

Hence $(a_n^p) \in \ell_\infty$, which is a contradiction. So in contrast to Prop. 2 we have the following result.

Proposition 4 *A smooth sequence space of infinite type that is also a Schwartz space has no prominent subset.*

In particular, an infinite type power series space $\Lambda_\infty(\alpha)$ has no prominent subset although $\Delta(\Lambda_\infty(\alpha)) = \Delta_b(\Lambda_\infty(\alpha))$.

References

- [1] Bierstedt, K-D., Bonet, J.: Some aspects of the modern theory of Fréchet spaces. RACSAM. Rev. R. Acad. Cienc. Exactas Ser. A. 97, 159–188 (2003).
- [2]] Bierstedt, K-D., Meise, R., Summers, W.H.: Köthe sets and Köthe sequence spaces. North-Holland Math. Studies 71, 27–91 (1982).
- [3] Jarchow, H.: Locally convex spaces. Stuttgart, B.G. Teubner 1981.
- [4] Köthe, G.: Topological vector spaces I and II. Berlin-Heidelberg-New York, Springer 1969 and 1979.
- [5] Meise, R., Vogt, D.: A characterization of the quasinormable Fréchet spaces. Math. Nachr. 122, 141–150 (1985).
- [6] Mitiagin, B.S.: Approximative dimension and basis in nuclear spaces. Russian Math. Surveys, 16, 59–127 (1961).
- [7] Pietsch, A.: Nuclear locally convex spaces. Berlin-Heidelberg-New York, Springer 1972.
- [8] Terzioğlu, T.: Die diametrale Dimension von lokalkonvexen Räumen. Collect. Math., 20, 49–99 (1969).
- [9] Terzioğlu, T.: On the diametral dimension of some classes of F -spaces. J. Karadeniz Uni. Ser. Math-Physics 8, 1–13 (1985).
- [10] Terzioğlu, T.: Diametral dimension and Köthe spaces. Turk.J.Math. 32, 213-218 (2008).