

1-1-2013

Products of conjugacy classes and products of irreducible characters in finite groups

MOHAMMAD REZA DARAFSHEH

SAJJAD MAHMOOD ROBATI

Follow this and additional works at: <https://dctubitak.researchcommons.org/math>



Part of the [Mathematics Commons](#)

Recommended Citation

DARAFSHEH, MOHAMMAD REZA and ROBATI, SAJJAD MAHMOOD (2013) "Products of conjugacy classes and products of irreducible characters in finite groups," *Turkish Journal of Mathematics*: Vol. 37: No. 4, Article 8. <https://doi.org/10.3906/mat-1112-34>
Available at: <https://dctubitak.researchcommons.org/math/vol37/iss4/8>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Mathematics by an authorized editor of TÜBİTAK Academic Journals.

Products of conjugacy classes and products of irreducible characters in finite groups

Mohammad Reza DARAFSHEH,* Sajjad MAHMOOD ROBATI

School of Mathematics, Statistics and Computer Science, College of Science, University of Tehran,
Tehran, Iran

Received: 25.12.2011 • Accepted: 10.05.2012 • Published Online: 12.06.2013 • Printed: 08.07.2013

Abstract: Let G be a finite group. If A and B are two conjugacy classes in G , then AB is a union of conjugacy classes in G and $\eta(AB)$ denotes the number of distinct conjugacy classes of G contained in AB . If χ and ψ are two complex irreducible characters of G , then $\chi\psi$ is a character of G and again we let $\eta(\chi\psi)$ be the number of irreducible characters of G appearing as constituents of $\chi\psi$. In this paper our aim is to study the product of conjugacy classes in a finite group and obtain an upper bound for η in general. Then we study similar results related to the product of two irreducible characters.

Key words: Conjugacy classes, irreducible characters, products

1. Introduction

Let G be a finite group, $a \in G$ and $Cl(a) = a^G = \{a^g | g \in G\}$ be the conjugacy class of a in G . If the subset X of G is G -invariant, i.e. $X^g = \{x^g | x \in X\} = X$ for all $g \in G$, then X is the union of m distinct conjugacy classes of G , for some positive integer m . Set $\eta(X) = m$. Given any conjugacy classes $Cl(a)$ and $Cl(b)$, we can check that the product $Cl(a)Cl(b) = \{a^h b^k | k, h \in G\}$ is a G -invariant set. Hence it is a union of conjugacy classes of G . Let $\eta(Cl(a)Cl(b))$ denote the number of conjugacy classes contained in $Cl(a)Cl(b)$. In this note, we study the product $Cl(a)Cl(b)$ when G is a finite group.

Let $[a, G]$ be the set of all commutators $[a, g] = a^{-1}g^{-1}ag$ where $g \in G$. Our first result is the following.

Theorem A. Let G be a finite group and $a \in G$. Then the following conditions are equivalent:

- (i) $[a, G]$ is a subgroup of G ;
- (ii) $Cl(a)Cl(a^{-1}) = [a, G]$;
- (iii) $|Cl(a)Cl(a^{-1})| = |Cl(a)|$;
- (iv) $|(Cl(a))^n| = |Cl(a)|$ for all $n \in \mathbb{N}$.

An application of Theorem A is the following.

Theorem B. Let G be a finite group and $a, b \in G$ and $[a, G]$ be a subset of $Z(G)$. Then

- (i) $\eta(Cl(a)Cl(b)) = |Cl(a)||Cl(b)|/|[a, G] \cap (Cl(b^{-1})Cl(b))||Cl(ab)|$;
- (ii) If $Cl(a)Cl(b) \cap Z(G) \neq \emptyset$, then $\eta(Cl(a)Cl(b)) = |Cl(a)|$;

*Correspondence: darafsheh@ut.ac.ir

2000 AMS Mathematics Subject Classification: 20E45, 20C15.

(iii) If $|Cl(a)|$ is an odd number, then $\eta(Cl(a)Cl(a)) = 1$;

(iv) If $|Cl(a)|$ is an even number, then $\eta(Cl(a)Cl(a)) = 2^n$ (where n is the number of cyclic direct factors in the decomposition of the Sylow 2-subgroup of $[a, G]$).

In [1], Adan-Bante shows that if G is a finite p -group and $a \in G$ and $|Cl(a)| = p^n$, then $\eta(Cl(a)Cl(a^{-1})) \geq n(p - 1) + 1$. In this paper, we prove that if G is a finite group and $|Cl(a)|$ is an odd number, then $\eta(Cl(a)Cl(a)) \leq (|Cl(a)| + 1)/2$ for all $a \in G$. Also, in Proposition 2.5, we prove that if G is a finite group and $a \in G$ and $|a|$ is an odd number, then $[a, G]$ is a subgroup of G if and only if $\eta(Cl(a)Cl(a)) = 1$, where $|a|$ is the order of a .

The case of products of two irreducible characters of a finite group has been of interest to researchers as well. For a finite group G , let χ and φ be irreducible characters of G , then obviously $\chi\varphi$ is a character of G , hence $\chi\varphi = \sum_{i=1}^k m_i\chi_i$, $m_i \in \mathbb{N}$, and $\chi_i \in \text{Irr}(G)$, hence we define $\eta(\chi\varphi) = k$. Analogous results on the products of the irreducible characters of G may be true for the products of conjugacy classes of G . For example we may mention [4], which studies this analogy. But in some cases this analogy may not hold.

In [7] it is proved that, there are two non-identity irreducible characters χ and φ such that $\chi\varphi$ is an irreducible character of A_n if and only if n is a perfect square, while in [3] it is proved that the product of non-trivial conjugacy classes in A_n for $n \geq 5$ is never a conjugacy class. Concerning irreducible characters of a finite p -group G , it is proved in [2] that if $\chi, \varphi \in \text{Irr}(G)$ are faithful, then either $\eta(\chi\varphi) = 1$ or $\eta(\chi\varphi) \geq (p + 1)/2$.

Concerning irreducible characters of a finite group G , in this paper the following results are also proved.

Theorem C. Let G be a finite group of odd order and $\chi \in \text{Irr}(G)$. Then χ vanishes on $G - Z(\chi)$ if and only if $\eta(\chi^2) = 1$.

Theorem D. Let G be a finite group and $\chi, \varphi \in \text{Irr}(G)$ such that χ vanishes on $G - Z(\chi)$. Then

- (i) If $\chi\varphi \cap \text{Lin}(G) \neq \emptyset$, then $\eta(\chi\varphi) = |\text{Irr}(G/Z(G))|$;
- (ii) If $\chi(1)$ is an odd number, then $\eta(\chi^2) = 1$.

We also put forward the following conjecture.

Conjecture. Let G be a finite group of odd order and $\chi \in \text{Irr}(G)$, then $\eta(\chi^2) \leq (\chi(1) + 1)/2$.

2. Proof of Theorems A and C

Denote by $C_G(a) = \{g \in G | a^g = a\}$ the centralizer of a in G . Also let $[a, g] = a^{-1}a^g$ and $[a, G] = \{[a, g] | g \in G\}$. Since $a^g = a[a, g]$ for any $g \in G$, we have that $Cl(a) = a^G = \{ax | x \in [a, G]\}$. In particular, we get that $|Cl(a)| = |[a, G]|$. Through this note, we will use the well-known fact that $|Cl(a)| = |G : C_G(a)|$.

Lemma 2.1 *Let G be a finite group and $a \in G$ and $[a, G]$ be a subgroup of G . Then*

- (i) $[a, G]$ is a normal subgroup of G ;
- (ii) $[a, G] = [a^{-1}, G]$.

Proof (i) Observe that

$$[a, g]^h = (a^{-1})^h a^{gh} = [a, h]^{-1} [a, gh]$$

for all $g, h \in G$. Since $[a, G]$ is a subgroup of G , then $[a, h]^{-1} [a, gh] \in [a, G]$. Therefore $[a, G]$ is a normal subgroup of G .

(ii) We have that

$$[a^{-1}, g] = a(a^{-1})^g = (a^{-1})^{ga^{-1}} a = [a, ga^{-1}]^{-1}$$

for all $g \in G$. Also since $[a, G]$ is a subgroup of G , then $[a, ga^{-1}]^{-1} \in [a, G]$. Therefore $[a^{-1}, G] \subseteq [a, G]$. Since $|Cl(a)| = |Cl(a^{-1})|$ we have $[a, G] = [a^{-1}, G]$. \square

Proof [Proof of Theorem A]

(i \Rightarrow ii)

By Lemma 2.1, observe that

$$Cl(a)Cl(a^{-1}) = a[a, G]a^{-1}[a^{-1}, G] = [a, G][a^{-1}, G] = [a, G].$$

(ii \Rightarrow i)

Observe that, since $a^G a = aa^G$ we can write

$$Cl(a)Cl(a^{-1}) = a[a, G]a^{-1}[a^{-1}, G] = [a, G][a^{-1}, G].$$

Fix $l, h \in G$. Since $Cl(a)Cl(a^{-1}) = [a, G]$ we have $[a, G][a^{-1}, G] = [a, G]$ and $[a, h][a^{-1}, k] = [a, l]$ for some $k \in G$.

Since $[a^{-1}, k]^{-1} = [a, ka^{-1}]$ we have $[a, h] = [a, l][a, ka^{-1}]$ and $[a, h][a, l]^{-1} = [a, ka^{-1}]$. Thus $1_G \in [a, G]$, and hence $[a, G]$ is a subgroup of G .

(ii \Leftrightarrow iii)

Since $a^{-1}a^G = a^G a^{-1}$ we have $[a, G] \subseteq Cl(a)Cl(a^{-1})$. Therefore from $|[a, G]| = |Cl(a)|$, we obtain $Cl(a)Cl(a^{-1}) = [a, G]$ if and only if $|Cl(a)Cl(a^{-1})| = |Cl(a)|$.

(i \Rightarrow iv)

Observe that

$$(Cl(a))^n = (a[a, G])^n = a^n[a, G]$$

for all $n \in \mathbb{N}$. Therefore $|(Cl(a))^n| = |Cl(a)|$.

(iv \Rightarrow i)

Let $n = 2$. Observe that, since $a^G a = aa^G$, we obtain

$$Cl(a)Cl(a) = a[a, G]a[a, G] = a^2[a, G][a, G].$$

Thus if $|Cl(a)Cl(a)| = |Cl(a)|$, then $|[a, G][a, G]| = |[a, G]|$. Since $1_G \in [a, G]$ we obtain $[a, G] \subseteq [a, G][a, G]$ and $[a, G][a, G] = [a, G]$. Then we conclude that $[a, G]$ is a subgroup of G , since $xy \in [a, G]$ for any x, y in $[a, G]$ and $[a, G]$ is a nonempty finite set. \square

Proposition 2.2 *Let G be a finite group and $a, b \in G$ and $Cl(a)Cl(b) \cap Z(G) \neq \emptyset$. Then $[a, G]$ is a subgroup of G if and only if $|Cl(a)Cl(b)| = |Cl(a)|$.*

Proof Suppose that there exists some $z \in Cl(a)Cl(b) \cap Z(G)$. Then there exists some $g \in G$ such that $z = a^g b$ and $b = (a^{-1})^g z$. Thus we have

$$Cl(a)Cl(b) = Cl(a)Cl((a^{-1})^g z) = Cl(a)Cl((a^{-1})^g)z = Cl(a)Cl(a^{-1})z,$$

therefore $|Cl(a)Cl(b)| = |Cl(a)Cl(a^{-1})|$. Thus by Theorem A, $[a, G]$ is a subgroup of G if and only if $|Cl(a)Cl(b)| = |Cl(a)Cl(a^{-1})| = |Cl(a)|$.

□

Proposition 2.3 *Let G be a finite group and $\chi, \psi \in \text{Irr}(G)$ with $\text{Irr}(\chi\psi) \cap \text{Lin}(G) \neq \emptyset$. Then χ vanishes on $G - Z(G)$ if and only if $\widehat{\chi\psi} = \rho_{G/Z(G)}$, where $\rho_{G/Z(G)}$ is the regular character of $G/Z(G)$ and $\widehat{\chi\psi}(gZ(G)) := \chi\psi(g)$ for all $g \in G$.*

Proof Suppose that χ vanishes on $G - Z(G)$ and $\lambda \in \text{Lin}(G) \cap \text{Irr}(\chi\psi)$. Since χ and $\lambda\bar{\psi}$ are irreducible characters then $0 \neq [\chi\psi, \lambda] = [\chi, \lambda\bar{\psi}] = 1$ and $\chi = \lambda\bar{\psi}$. Therefore $\chi\psi = \lambda\bar{\psi}\psi$ and ψ vanishes on $G - Z(G)$. We can check that $\eta(\chi\psi) = \eta(\bar{\psi}\psi)$.

We can write $\bar{\psi}(g)\psi(g) = |\psi(g)|^2 = \psi(1)^2$ for all $g \in Z(G)$ and $\bar{\psi}(g)\psi(g) = 0$ for all $g \in G - Z(G)$, thus $Z(G) = \ker(\bar{\psi}\psi)$. Suppose that $\varphi_i \in \text{Irr}(\bar{\psi}\psi)$ for $i = 1, \dots, n$, then $Z(G) = \ker(\bar{\psi}\psi) = \bigcap_{i=1}^n \ker(\varphi_i) \subseteq \ker(\varphi_i)$ and

$$\begin{aligned} 0 \neq [\bar{\psi}\psi, \varphi_i] &= 1/|G| \sum_{g \in G} \bar{\psi}\psi(g) \overline{\varphi_i}(g) \\ &= 1/|G| \sum_{g \in Z(G)} \bar{\psi}(g)\psi(g) \overline{\varphi_i}(g) \\ &= \frac{|Z(G)|\psi(1)^2\varphi_i(1)}{|G|} = \varphi_i(1) \end{aligned} \tag{2.1}$$

because $\psi(1)^2 = |G : Z(G)|$. Thus $\bar{\psi}\psi = \sum_{i=1}^n \varphi_i(1)\varphi_i$ and $|G : Z(G)| = \psi(1)^2 = \sum_{i=1}^n \varphi_i(1)^2$. Since $\text{Irr}(G/Z(G)) = \{\theta \in \text{Irr}(G) | Z(G) \subseteq \ker\theta\}$ and $\rho_{G/Z(G)} = \sum_{\hat{\theta} \in \text{Irr}(G/Z(G))} \hat{\theta}(1)\hat{\theta}$ then $\hat{\varphi}_i \in \text{Irr}(G/Z(G))$ and $\widehat{\bar{\psi}\psi} = \rho_{G/Z(G)}$, where $\hat{\varphi}_i(gZ(G)) := \varphi_i(g)$ for all $g \in G$. It follows that $\widehat{\chi\psi} = \rho_{G/Z(G)}$.

Conversely suppose $\widehat{\chi\psi} = \rho_{G/Z(G)} = \sum_{\hat{\theta} \in \text{Irr}(G/Z(G))} \hat{\theta}(1)\hat{\theta}$, then $\chi\psi$ vanishes on $G - Z(G)$. On the other hand, we have $\text{Irr}(\chi\psi) \cap \text{Lin}(G) \neq \emptyset$, then there exists some $\lambda \in \text{Lin}(G)$ such that $\lambda \in \text{Irr}(\chi\psi)$ and $0 \neq [\chi\psi, \lambda] = [\chi, \lambda\bar{\psi}] = 1$. Thus we have $\chi = \lambda\bar{\psi}$ and $\chi\psi = \lambda\bar{\psi}\psi$. It follows that $\bar{\psi}\psi$ vanishes on $G - Z(G)$ and χ vanishes on $G - Z(G)$. □

Lemma 2.4 *Let G be a finite group and $a \in G$. If $[a, G]$ is not a subgroup of G , then $\eta(Cl(a)Cl(a)) \neq 1$.*

Proof We have that $C_G(a) \subseteq C_G(a^2)$, therefore $|Cl(a^2)| \leq |Cl(a)|$. Since $[a, G]$ is not a subgroup of G , then by proof of Theorem A, $|Cl(a^2)| \leq |Cl(a)| < |Cl(a)Cl(a)|$. Thus $Cl(a)Cl(a)$ is not a single conjugacy class. □

Proposition 2.5 *Let G be a finite group and $a \in G$ and the order of a be an odd number. Then $[a, G]$ is a subgroup of G if and only if $\eta(Cl(a)Cl(a)) = 1$.*

Proof We have $C_G(a) \subseteq C_G(a^2)$. Since $|a|$ is an odd number, we can check that $C_G(a) = C_G(a^2)$. Thus

$$|Cl(a)| = |G : C_G(a)| = |G : C_G(a^2)| = |Cl(a^2)|.$$

Then by Theorem A, $Cl(a)Cl(a) = Cl(a^2)$. Conversely, by Lemma 2.4, if $\eta(Cl(a)Cl(a)) = 1$ then $[a, G]$ is a subgroup of G and the proof is complete. \square

At this point we prove Theorem C, which is an analogue of Proposition 2.5.

Proof [Proof of Theorem C] First suppose χ vanishes on $G - Z(\chi)$. By Corollary 2.30 of [5] we have $\chi(1)^2 = |G : Z(\chi)|$. By Theorem 4.5 of [5] the alternating part of the character χ^2 which is denoted by χ_A is a character of G and for $g \in G$ we have $\chi_A(g) = 1/2(\chi(g)^2 - \chi(g^2))$. Let us define $\chi^{(2)}$ by $\chi^{(2)}(g) = \chi(g^2)$, for all $g \in G$, and χ^2 by $\chi^2(g) = \chi(g)^2$, for all $g \in G$. Then $\chi^{(2)} = \chi^2 - 2\chi_A$. Since $|G|$ is odd, by Problem 4.5 of [5], $\chi^{(2)} \in \text{Irr}(G)$, hence

$$\begin{aligned} 0 \neq [\chi^2, \chi^{(2)}] &= 1/|G| \sum_{g \in G} \chi^2(g) \overline{\chi^{(2)}(g)} \\ &= 1/|G| \sum_{g \in Z(\chi)} \chi^2(g) \overline{\chi^{(2)}(g)}. \end{aligned} \tag{2.2}$$

But $\chi|_{Z(\chi)} = \chi(1)\lambda_1$ and $\chi^{(2)}|_{Z(\chi)} = \chi^{(2)}(1)\lambda_2 = \chi(1)\lambda_2$ where $\lambda_1, \lambda_2 \in \text{Lin}(Z(\chi))$. Therefore continuing, (2.2), we can write:

$$\begin{aligned} 0 \neq [\chi^2, \chi^{(2)}] &= \chi(1)^3/|G| \sum_{g \in Z(\chi)} \lambda_1^2(g) \overline{\lambda_2(g)} \\ &= (\chi(1)^3|Z(\chi)|/|G|)[\lambda_1^2, \lambda_2]. \end{aligned} \tag{2.3}$$

But $[\lambda_1^2, \lambda_2] = 0$ or 1 , which implies $\lambda_1^2 = \lambda_2$ and $0 \neq [\chi^2, \chi^{(2)}] = \chi(1)$. Therefore $\chi^2 = \chi(1)\chi^{(2)}$, proving $\eta(\chi^2) = 1$.

Conversely suppose G has odd order and $\eta(\chi^2) = 1$. Since $\chi^{(2)} \in \text{Irr}(\chi^2)$ and $\eta(\chi^2) = 1$, then $\chi^2 = \chi(1)\chi^{(2)}$. If $\chi(g) \neq 0$, then

$$|\chi(g)| = \frac{\chi(1)}{|\chi(g)|} |\chi^{(2)}(g)| \geq |\chi^{(2)}(g)|. \tag{2.4}$$

Thus $1 = [\chi^{(2)}, \chi^{(2)}] \leq [\chi, \chi] = 1$ and it follows that $|\chi(g)| = |\chi^{(2)}(g)|$. Therefore by (2.4), we have that $\chi(1) = |\chi(g)|$ and $g \in Z(\chi)$. Thus if $\chi(g) \neq 0$ then $g \in Z(\chi)$, and therefore χ vanishes on $G - Z(\chi)$. \square

Example 2.6 Let D_8 be the dihedral group of order 2^3 and $a \in D_8 \setminus Z(D_8)$. We can check that $Cl(a)Cl(a) = Cl(a^{-1})Cl(a) = Z(D_8)$, therefore $\eta(Cl(a)Cl(a)) = \eta(Cl(a^{-1})Cl(a)) = 2$. Thus Proposition 2.5 may not remain true if $a \in G$ has even order. On the other hand, let χ be the faithful irreducible character of D_8 . We can check that χ vanishes on $G - Z(G)$, but $\eta(\chi^2) = 4$. Thus Theorem C may not remain true if G has even order.

3. Upper bound

Lemma 3.1 Let G be a group and $a, b \in G$. Then $\eta(Cl(a)Cl(b)) \leq |Cl(a)|$.

Proof For $c \in Cl(a)Cl(b)$ we have $c = g^{-1}agh^{-1}bh$ where $g, h \in G$, we deduce that c is conjugate to $hg^{-1}agh^{-1}b = a^{gh^{-1}}b$, and hence there is $d \in G$ such that $c^d \in a^G b$. Therefore $c \in \bigcup_{g \in G} Cl(a^g b)$, implying

that $Cl(a)Cl(b) \subseteq \bigcup_{g \in G} Cl(a^g b)$. Clearly $\bigcup_{g \in G} Cl(a^g b) \subseteq Cl(a)Cl(b)$, hence $Cl(a)Cl(b) = \bigcup_{g \in G} Cl(a^g b)$, thus $\eta(Cl(a)Cl(b)) = \eta(\bigcup_{g \in G} Cl(a^g b)) \leq |a^G b| = |Cl(a)|$. \square

Proposition 3.2 *Let G be a finite group and $a \in G$. If $|Cl(a)|$ is an odd number, then $Cl(a)Cl(a)$ is the union of at most $(|Cl(a)| + 1)/2$ distinct conjugacy classes, i.e. $\eta(Cl(a)Cl(a)) \leq (|Cl(a)| + 1)/2$.*

Proof By Lemma 3.1, we have that $Cl(a)Cl(a) = \bigcup_{g \in G} Cl(a^g a)$. It suffices to show that for any $x \in Cl(a)$ with $x \neq a$, there exists $y \in Cl(a)$, $y \neq x, a$, such that $Cl(xa) = Cl(ya)$. Observe that $a^g a$ and $a^{g^{-1}} a$ are always conjugate; thus, if $x = a^g$ and g^2 does not centralize a , one can take $y = a^{g^{-1}}$.

Suppose on the other hand that $g^2 \in C_G(a)$ whenever $x = a^g$. We claim that this happens only when $x = a$. Indeed, fix a, g with $x = a^g$. Then $(zg)^2 \in C_G(a)$ for all $z \in C_G(a)$ and taking into account that $(zg)^2 = zz^{g^{-1}} g^2$, it follows that g normalizes $C_G(a)$. Now let P be a Sylow 2-subgroup of $C_G(a)$ (which is also a Sylow 2-subgroup of $C_G(a)$ since $|Cl(a)| = n$ is odd). Thus $P^g \subseteq C_G(a)$, therefore $P^{z^g} = P$ for some $z \in C_G(a)$. By replacing g with zg , we have that g normalizes P . Thus the 2-part of g is in $P \subseteq C_G(a)$ and since $g^2 \in C_G(a)$ then $g \in C_G(a)$. Therefore we can suppose that g has odd order. But then $g \in C_G(a)$ and $x = a$. Thus since $Cl(a)Cl(a) = \bigcup_{i=1}^{n-1} Cl(a^{g^i} a) \cup Cl(a^2)$ for some $\{g_i\}_{i=1}^{n-1} \subseteq G \setminus C_G(a)$, then

$$\eta(Cl(a)Cl(a)) \leq \frac{|Cl(a)| - 1}{2} + 1 = \frac{|Cl(a)| + 1}{2}.$$

\square

Proposition 3.3 *Let G be a finite group and $a \in G$ and $|Cl(a)| = 2$. Then $\eta(Cl(a)Cl(a)) = \eta(Cl(a)Cl(a^{-1})) = |Cl(a)| = 2$.*

Proof Let $g \in G \setminus C_G(a)$. Observe that $Cl(a) = \{a, a^g\}$ and $a^2, aa^g \in Cl(a)Cl(a)$. By Lemma 3.1, we have $Cl(a)Cl(a) = Cl(a^2) \cup Cl(aa^g)$. Since $a^2 \neq aa^g \neq (a^2)^g$ then $aa^g \notin Cl(a^2)$. Therefore $Cl(a^2) \neq Cl(aa^g)$ and $\eta(Cl(a)Cl(a)) = |Cl(a)| = 2$.

Also, by Lemma 3.1, we have $\eta(Cl(a)Cl(a^{-1})) \leq 2$. Since $1_G \in Cl(a)Cl(a^{-1})$ and $a \notin Z(G)$, then $\eta(Cl(a)Cl(a^{-1})) = 2$

\square

Remark 3.4 *Let p and q be two prime numbers such that $p \mid q - 1$. Let G be a finite nonabelian group of order pq and $a \in G$ and $|a| = q$.*

We can check that $|Cl(a)| = p$ and $|Cl(a)Cl(a)| = p(p + 1)/2$. Also if $p \neq 2$, then all of the conjugacy classes contained in $Cl(a)Cl(a)$ are of size p and therefore $\eta(Cl(a)Cl(a)) = (p + 1)/2$. Thus the bound in Proposition 3.2 is optimal.

Otherwise, if $p = 2$, then $|Cl(a)Cl(a)| = 3$ and $Cl(a)Cl(a) = 1_G \cup Cl(a^2)$. Therefore $\eta(Cl(a)Cl(a)) = 2$.

4. Simple groups and symmetric groups

Proposition 4.1 *Let G be a finite nonabelian group and $a \in G$. Then*

- (i) If $a \in G'$, then $Cl(a) \subsetneq G'$;
- (ii) If $a \notin G'$, then $G' \cap Cl(a) = \emptyset$.

Proof (i) Let $a \in G'$. If $a = 1_G$ then the statement is true. Otherwise, since $a^{-1}a^G = [a, G] \subseteq G'$, therefore $Cl(a) = a^G \subseteq G'$. Since $1_G \notin Cl(a)$, thus $Cl(a) \subsetneq G'$.

(ii) Let $a \notin G'$. Since $a^{-1}a^G = [a, G]$, therefore $Cl(a) \cap G' = \emptyset$. □

Corollary 4.2 *Let G be a finite nonabelian simple group and $a \in G$. Then $[a, G]$ is a subgroup of G if and only if $a = 1_G$.*

Proof Assume that $a \neq 1_G$ and $[a, G]$ be a subgroup of G . Therefore by Lemma 2.1, $[a, G]$ is a normal subgroup of G . Since G is a finite nonabelian simple group and $[a, G] \neq 1_G$, then $[a, G] = G' = G$. By Proposition 4.1, $|Cl(a)| \neq G'$, we have that $[a, G] \neq G'$ and the proof is complete. □

Corollary 4.3 *Let G be a finite nonabelian simple group and $1_G \neq a \in G$. If $|Cl(a)|$ is an odd number, then $2 \leq \eta(Cl(a)Cl(a)) \leq (|Cl(a)| + 1)/2$.*

Proof It follows from Corollary 4.2 and Lemma 2.4 and Proposition 3.2. □

Proposition 4.4 *Let S_n be the symmetric group of degree n and $\alpha \in S_n$. Then $[\alpha, S_n]$ is a subgroup of S_n for $n > 4$ if and only if $\alpha = id$.*

Proof Assume that $\alpha \neq id$ and $[\alpha, S_n]$ be a subgroup of S_n . Since $[\alpha, S_n] \subseteq S'_n = A_n$ for $n > 4$, thus $[\alpha, S_n] = A_n$. Therefore, by Theorem A, $|Cl(\alpha)| = n!/2$, thus

$$|Cl(\alpha)| = \frac{n!}{1^{e_1}e_1!2^{e_2}e_2!\dots n^{e_n}e_n!} = \frac{n!}{2}$$

where $1^{e_1}, \dots, n^{e_n}$ is the cycle structure of $\alpha \in S_n$. It is easy to see that $|Cl(\alpha)| \neq n!/2$ for $n > 4$. Therefore $[\alpha, S_n]$ is not a subgroup of S_n for $n > 4$. □

Remark 4.5 *We can check that $[\alpha, S_4]$ is not a subgroup of S_4 for all $id \neq \alpha \in S_4$.*

Remark 4.6 *Observe that if $Cl(a)Cl(a^{-1})$ is a subgroup of G , then $Cl(a)Cl(a^{-1})$ is a normal subgroup of G , since $Cl(a)Cl(a^{-1})$ is G -invariant.*

The authors in [3] show that if $G \simeq J_1$ (where J_1 is the Janko group of order 175560) then $Cl(a)Cl(a^{-1}) = J_1$ for all $a \in J_1 \setminus \{1_G\}$. But in Corollary 4.2, we showed that if G is a finite nonabelian simple group, then $[a, G] \neq G$ for all $a \in G$.

Thus if G is a finite nonabelian simple group and $Cl(a)Cl(a^{-1})$ is a subgroup of G , then it is not necessary that $[a, G]$ is a subgroup of G .

5. Proof of Theorems B and D

Lemma 5.1 *Let G be a finite group and $a \in G$ and $[a, G]$ is a subset of $Z(G)$. Then $[a, G]$ is a subgroup of G .*

Proof Since $[a, G] \subseteq Z(G)$, thus $[a, g_1] = z_1$ and $[a, g_2] = z_2$ for all $g_1, g_2 \in G$ and for some $z_1, z_2 \in Z(G)$. It is easy to see that $[a, g_1][a, g_2] = z_1z_2 = [a, g_1g_2]$. Since $1_G \in [a, G]$ and G is a finite group, it follows that $[a, G]$ is a subgroup of G . □

Proposition 5.2 *Let G be a finite group and $a, b \in G$ and $[a, G] \subseteq Z(G)$. Then all of the conjugacy classes contained in $Cl(a)Cl(b)$ are of size $|Cl(ab)|$. Therefore $\eta(Cl(a)Cl(b)) = |Cl(a)Cl(b)|/|Cl(ab)|$.*

Proof Since $Cl(a)Cl(b)$ is a G -invariant set, then

$$Cl(a)Cl(b) = \{Cl(a^g b) | g \in G\} = \{Cl(abz) | z \in [a, G]\} = \{Cl(ab)z | z \in [a, G]\}.$$

Therefore all of the conjugacy classes contained in $Cl(a)Cl(b)$ are of size $|Cl(ab)|$ and the proof is complete. □

Corollary 5.3 *Let G be a finite group and $a, b \in G$ and $[a, G] \subseteq Z(G)$. If $Cl(a)Cl(b) \cap Z(G) \neq \emptyset$, then $\eta(Cl(a)Cl(b)) = |Cl(a)|$.*

Proof By proposition 5.2, we have that

$$Cl(a)Cl(b) = \{Cl(ab)z | z \in [a, G]\}.$$

Observe that since $Cl(a)Cl(b) \cap Z(G) \neq \emptyset$, then $(ab)^g \in Z(G)$ for some $g \in G$, therefore $ab \in Z(G)$. Thus $Cl(a)Cl(b) \subseteq Z(G)$ and $\eta(Cl(a)Cl(b)) = |Cl(a)Cl(b)| = |Cl(a)|$. □

Proof [Proof of Theorem B]

(i) Observe that

$$Cl(a)Cl(b) = a[a, G]Cl(b) = aCl(b)[a, G] \tag{5.1}$$

and $[a, G]$ is a subgroup of $Z(G)$.

Assume that $X = \{Cl(g) | g \in G\}$. $[a, G]$ acts on X by right multiplication in G . Thus the stabilizer of $Cl(b)$ in $[a, G]$ is

$$\begin{aligned} St_{[a,G]}(Cl(b)) &= \{z \in [a, G] | Cl(b)z = Cl(b)\} \\ &= \{z \in [a, G] | z \in Cl(b)Cl(b^{-1})\} \\ &= [a, G] \cap (Cl(b)Cl(b^{-1})) \end{aligned} \tag{5.2}$$

and the orbit of $Cl(b)$ is

$$Orb_{[a,G]}(Cl(b)) = \{Cl(b)z | z \in [a, G]\}. \tag{5.3}$$

Since $|Cl(b)| = |Cl(b)z|$ for all $z \in [a, G]$, then $|Cl(b)[a, G]| = |Cl(b)||Orb_{[a,G]}(Cl(b))|$. Thus

$$|Cl(b)[a, G]| = |Cl(b)||[a, G] : St_{[a,G]}(Cl(b))| = \frac{|Cl(b)||[a, G]|}{|[a, G] \cap (Cl(b)Cl(b^{-1}))|}. \tag{5.4}$$

Therefore by 5.1 and 5.4, we have that

$$|Cl(a)Cl(b)| = |Cl(b)[a, G]| = \frac{|Cl(a)||Cl(b)|}{|[a, G] \cap (Cl(b)Cl(b^{-1}))|}. \tag{5.5}$$

Thus Proposition 5.2 and 5.5 imply (i).

(ii) Follows from Corollary 5.3.

(iii) By Lemma 5.1 we have that $[a, G]$ is a subgroup $Z(G)$, thus

$$Cl(a)Cl(a) = a[a, G]a[a, G] = a^2[a, G][a, G] = a^2[a, G].$$

Therefore $|Cl(a)Cl(a)| = |a^2[a, G]| = |Cl(a)|$. Also there exists $n \in \mathbb{N}$ such that $[a, G] = \langle z_1 \rangle \times \cdots \times \langle z_n \rangle$ for some $z_i \in Z(G)$, $1 \leq i \leq n$ (where $|z_i| = p_i^{\alpha_i}$ for $1 \leq i \leq n$ and $\alpha_i \in \mathbb{N}$). Thus

$$Cl(a) = a[a, G] = a\langle z_1 \rangle \times \cdots \times \langle z_n \rangle.$$

Also we have

$$\begin{aligned} Cl(a^2) &= \{az_1^{i_1} \dots z_n^{i_n} az_1^{i_1} \dots z_n^{i_n} \mid i_j = 0, \dots, |z_j| - 1, j = 0, \dots, n\} \\ &= \{a^2 z_1^{2i_1} \dots z_n^{2i_n} \mid i_j = 0, \dots, |z_j| - 1, j = 0, \dots, n\} \\ &= a^2(\langle z_1^2 \rangle \times \cdots \times \langle z_n^2 \rangle). \end{aligned}$$

Next if $|Cl(a)|$ is an odd number, then $\langle z_j^2 \rangle = \langle z_j \rangle$ for $j = 1, \dots, n$. Thus $Cl(a)Cl(a) = a^2[a, G] = Cl(a^2)$ and ii) follows.

(iv) Otherwise, let $|Cl(a)|$ be an even number and P be a Sylow 2-subgroup of $[a, G]$. Assume that $P = \langle z_1 \rangle \times \cdots \times \langle z_m \rangle$ (where $|z_i| = 2^{\alpha_i}$ for $1 \leq i \leq m$ and $\alpha_i \in \mathbb{N}$). Thus we have that

$$\begin{aligned} Cl(a^2 z_1^{l_1} \dots z_m^{l_m}) &= Cl(a^2) z_1^{l_1} \dots z_m^{l_m} \\ &= a^2 (\langle z_1^2 \rangle \times \cdots \times \langle z_m^2 \rangle) z_1^{l_1} \dots z_m^{l_m} \end{aligned}$$

for $l_1, \dots, l_m = 0, 1$. Therefore

$$Cl(a)Cl(a) = \bigcup_{l_1, \dots, l_m=0}^1 Cl(a^2 z_1^{l_1} \dots z_m^{l_m}).$$

Thus $Cl(a)Cl(a)$ is the union of exactly 2^m distinct conjugacy classes of G of size $|Cl(a^2)|$ and (iv) follows. \square

Proof [Proof of Theorem D]

(i) Follows from Proposition 2.3.

(ii) We can assume that χ is a faithful irreducible character and then $Z(\chi) = Z(G)$. Since χ vanishes on $G - Z(G)$, by Corollary 2.30 of [5] we can write $\chi(1)^2 = [G : Z(G)]$, therefore $|G : Z(G)|$ is an odd number. If $|G|$ is an odd number then by Theorem C $\eta(\chi^2) = 1$. Otherwise, let $|G|$ be an even number. If $P \in Syl_2(G)$

then $P \subseteq Z(G)$ and $P \trianglelefteq G$, because $|G : Z(G)|$ is an odd number. By Theorem 7.41 of [6], there is a subgroup H of G such that $|H| = |G : P|$. Since $P \trianglelefteq G$ then we can check that $G = HP = H \times P$. By problem 4.5 of [5], $\chi^{(2)} \in \text{Irr}(G)$ because $G = H \times P$ and P is an abelian subgroup of G and $(|H|, 2) = 1$. But by Theorem 4.5 of [5], $\chi^2 = \chi^{(2)} + 2\chi_A$ where $\chi_A \in \text{Char}(G)$. By the Proof of Theorem C we can write $0 \neq [\chi^2, \chi^{(2)}] = \chi(1)$ and $\chi^2 = \chi(1)\chi^{(2)}$. It follows that $\eta(\chi^2) = 1$. \square

References

- [1] Adan-Bante, E.: Conjugacy classes and finite p-groups. Arch. Math 85, 297-303 (2005).
- [2] Adan-Bante, E.: Products of characters and finite p-groups. J. Algebra 277, 236-255 (2004).
- [3] Arad, Z., Chillag, D. and Moran, G.: Groups with a small covering number. In: Products of Conjugacy Classes in Groups 222-244, Lecture Notes in Math. 1112. Berlin. Springer-Verlag (1985).
- [4] Arad, Z. and Fisman, E.: An analogy between products of two conjugacy classes and products of two irreducible characters in finite groups. Proc. Edin. Math. Soc. 30, 7-22 (1987).
- [5] Isaacs, I. M.: Character theory of finite groups. New York-San Francisco-London. Academic Press 1976.
- [6] Rotman, J. J.: An introduction to the theory of groups. New York. Springer-Verlag 1994.
- [7] Zisser, I.: Irreducible products of characters in A_n . Israel J. Math 84 no. 1-2, 147-151 (1993).