

1-1-2002

A New Record for the Myxomycetes Flora of Turkey: *Comatricha pulchella* (C.Bab.) Rost. var. *pulchella*

C. CEM ERGÜL

BAŞARAN DÜLGER

Follow this and additional works at: <https://journals.tubitak.gov.tr/botany>



Part of the [Botany Commons](#)

Recommended Citation

ERGÜL, C. CEM and DÜLGER, BAŞARAN (2002) "A New Record for the Myxomycetes Flora of Turkey: *Comatricha pulchella* (C.Bab.) Rost. var. *pulchella*," *Turkish Journal of Botany*: Vol. 26: No. 2, Article 7. Available at: <https://journals.tubitak.gov.tr/botany/vol26/iss2/7>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Botany by an authorized editor of TÜBİTAK Academic Journals. For more information, please contact academic.publications@tubitak.gov.tr.

A New Record for the Myxomycetes Flora of Turkey: *Comatricha pulchella* (C.Bab.) Rost. var. *pulchella*

C. Cem ERGÜL, Başaran DÜLGER
Uludağ University, Faculty of Arts & Science, Department of Biology, Bursa-TURKEY

Received: 19.02.2001

Accepted: 08.11.2001

Abstract: *Comatricha pulchella* (C.Bab.) Rost. var. *pulchella* is recorded for the first time from Turkey.

Key Words: Myxomycetes, *Comatricha*, Turkey

Türkiye Miksomiset Florası için Yeni Bir Kayıt: *Comatricha pulchella* (C.Bab.) Rost. var. *pulchella*

Özet: *Comatricha pulchella* (C.Bab.) Rost. var. *pulchella* Türkiye'den ilk defa kaydedilmektedir.

Anahtar Sözcükler: Myxomycetes, *Comatricha*, Türkiye

Introduction

The samples were collected from the edge of the Lake Abant (Bolu) in September, 1995. Samples were identified as *Comatricha pulchella* (C.Bab.) Rost. var. *pulchella* using the diagnostic literature (Martin & Alexopoulos, 1969; Farr, 1976; Nannenga-Bremekamp, 1991). This taxon is a new record for the Myxomycetes Flora of Turkey (Harkonen & Uotila, 1983; Harkonen, 1988; Lado, 1994; Ergül & Dülger, 2000). The samples are kept in the Herbarium of Uludağ University (Bursa).

Description of Species

Comatricha pulchella (C.Bab.) Rost. var. *pulchella*, Mon. App. 27. 1876 ; Syn: *Stemonitis pulchella* C.Bab., Proc. Linn. Soc. 1:32, 1839.

Sporangia gregarious, stipitate, ovate to cylindric, pale brown, 0.5-1.2 mm tall. Stalk black, shorter than the sporangium, 0.2-0.4 mm. Hypothallus usually discoid and small, rarely continuous under a group. Columella black, straight, tapering, becoming tortuous and coiled towards the apex. Capillitium dense, arising from the

entire columella, dark brown, primary branches darker and stouter, successive branches paler and slender, freely anastomosing branches and with few free ends. Spores brown in mass, paler in transmitted light, minutely punctate, 6.5-8.5 (-10) μ m in diameter (Figures 1-2).

Locality: Bolu – the edge of Lake Abant, Ergül 158-3, Fallen twigs and logs, 19.09.1995, Altitude Approx. 1600 m.

Known World Distribution: India, Ceylon, Japan, Europe, North America, Nigeria (Farr, 1976; Thind, 1977)

Result and Discussion

Comatricha pulchella can be distinguished from other similar-looking species of *Comatricha* as follows: This species often forms extensive fructification on small twigs; the reddish tint of the small, ovoid to cylindric sporangia, and the pale brown spores are its distinguishing marks (Farr, 1976). *Comatricha tenerrima* (Curtis) G.Lister is morphologically close to *Comatricha pulchella*, from which it is distinguished by




Figure 1. Stereomicroscopic image of the sporangia of *Comatricha pulchella* (C.Bab.) Rost. var. *pulchella*




Figure 2. A view of capillitial threads and spores of *Comatricha pulchella* (C.Bab.) Rost. var. *pulchella*

the pink colour, longer stipes, and smaller, more fusiform sporangia (Nannenga-Bremekamp, 1991).

Nannenga-Bremekamp (1991) recognized two varieties: *C. pulchella* var. *fusca* and *C. pulchella* var. *pulchella*. *Comatricha pulchella* var. *fusca* can be distinguished from var. *pulchella* by the usually somewhat longer sporangia, which are always dark brown (much darker than those of *C. pulchella* var. *pulchella*) and usually closer together; by the relatively short stalks; by the dark purple-brown capillitium, which often exhibits an interrupted surface net; and by the pale grey-brown spores in transmitted light. In addition, Lister (1966) described a variety as *gracilis*; but this has been transferred to the genus *Stemonitopsis* because of its

stalk structure, cylindrical sporangia and almost complete surface net.

The spore diameter of *Comatricha pulchella* var. *pulchella* given in the literature varies: Nannenga-Bremekamp (1991), 6.5-8.0 µm; Farr (1976), 6-8 µm; Martin & Alexopoulos (1969), 6.5-8.0 µm; Lakhanpal & Mukerji (1981), 7-8(-9) µm; Thind (1977), 7-9(-9.8) µm. The spores of our specimen were 6.5-8.5(-10) µm in diam. According to Nannenga-Bremekamp (1991), the total size is 0.7-1.5 mm, whereas Lakhanpal & Mukerji (1981) reported it to be 1.5-3.5 mm. The total size of our specimen was 0.5-1.2 mm tall (Table 1).

This record brings the species count for the genus *Comatricha* in Turkey to seven (Ergül & Dülger, 2000).

Table 1. Values of *Comatricha pulchella* var. *pulchella* from various studies.

	Total colour	Total size (mm)	Spore size (µm)	Spore colour By transmitted light
Nannenga-Bremekamp (1991)	Pale lilac-brown or cinnamon	0.7-1.5	6.5-8.0	Rosy-brown
Farr (1976)	Pale brown or ferruginous	0.7-1.5(-3.0)	6-8	Pale lilaceous brown
Martin & Alexopoulos (1969)	Pale brown or ferruginous	0.7-1.5(-3.0)	6.5-8.0	Pale lilac-brown
Lakhanpal & Mukerji (1981)	Bright ferruginous	1.5-3.5	7-8(-9)	Paler
Thind (1977)	Dark brown	0.5-1.3	7-9(-9.8)	Violaceous brown
Present specimen	Pale brown	0.5-1.2	6.5-8.5(-10)	Paler

References

- Ergül CC & Dülger B (2000). Myxomycetes of Turkey. *Karstenia* 40: 39-41.
- Farr ML (1976). *Flora Neotropica*. New York: The New York Botanical Garden.
- Harkonen M & Uotila P (1983). Turkish Myxomycetes Developed in Moist Chamber Cultures. *Karstenia* 23: 1-9.
- Harkonen M (1988). Some Additions to the Knowledge of Turkish Myxomycetes. *Karstenia* 27: 1-7.
- Lado C (1994) A Checklist of Myxomycetes of Mediterranean Countries. *Mycotaxon* LII(1): 117-185.
- Lakhanpal TN & Mukerji KG (1981). *Taxonomy of the Indian Myxomycetes*. AR Gantner Verlag KG Press Germany.
- Lister G (1966). The Genera of Myxomycetes. *Univ Iowa Stud Nat Hist* 20(8): 1-32.
- Martin GW & Alexopoulos CJ (1969). *The Myxomycetes*. Iowa City: Univ Iowa Press.
- Nannenga-Bremekamp NE (1991). *A Guide to Temperate Myxomycetes*. Bristol: Biopress Limited.
- Thind KS (1977). *The Myxomycetes of India*. New Delhi: Indian Council of Agricultural Research.