

1-1-2007

Gastropod Shell Species Occupied by Hermit Crabs (Anomura: Decapoda) along the Turkish Coast of the Aegean Sea

A. SUAT ATEŞ

TUNCER KATAĞAN

AHMET KOCATAŞ

Follow this and additional works at: <https://journals.tubitak.gov.tr/zoology>



Part of the [Zoology Commons](#)

Recommended Citation

ATEŞ, A. SUAT; KATAĞAN, TUNCER; and KOCATAŞ, AHMET (2007) "Gastropod Shell Species Occupied by Hermit Crabs (Anomura: Decapoda) along the Turkish Coast of the Aegean Sea," *Turkish Journal of Zoology*. Vol. 31: No. 1, Article 3. Available at: <https://journals.tubitak.gov.tr/zoology/vol31/iss1/3>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Zoology by an authorized editor of TÜBİTAK Academic Journals. For more information, please contact academic.publications@tubitak.gov.tr.

Gastropod Shell Species Occupied by Hermit Crabs (Anomura: Decapoda) along the Turkish Coast of the Aegean Sea

A. Suat ATEŞ^{1,*}, Tuncer KATAĞAN², Ahmet KOCATAŞ²

¹Çanakkale Onsekiz Mart University, Fisheries Faculty 17020 Çanakkale - TURKEY

²Ege University, Fisheries Faculty 35100 Bornova-İzmir - TURKEY

Received: 19.07.2005

Abstract: This study aimed to describe the gastropod shell species inhabited by hermit crabs (Anomura: Decapoda) collected during investigations of the decapod fauna of the Turkish Aegean Sea coast. We documented 12 hermit crab species inhabiting 16 species of gastropod shells. The hermit crab, *Pagurus anachoretus* Risso, 1827, had the highest inhabitation frequency (IF), inhabiting 37.5% of all the samples collected. The gastropod shell most inhabited was *Bittium latreillii* (Payraudeau, 1826), which accounted for 66.7% of inhabited shells.

Key Words: Shell utilization, Anomura Decapoda, Eastern Aegean Sea, Turkey

Ege Denizi Türkiye Kıyılarındaki Paguroid (Anomura: Decapoda)'ler Tarafından İşgal Edilen Gastropod Kabuğu Türleri

Özet: Bu çalışma Ege Denizi Türkiye Kıyıları dekapod faunası üzerine yürütülen araştırmalar sırasında toplanan Paguroid (Anomura: Decapoda)'ler tarafından kullanılan gastropod kabuklarını tanımlamayı amaçlamaktadır. 16 farklı gastropod kabuğunu kullanan toplam 12 Paguroid türü türü kaydedilmiştir. Paguroid türü *Pagurus anachoretus* Risso, 1827 % 37,5'lik bir değerle en yüksek yerleşme frekansına sahiptir. En fazla işgal edilen gastropod kabuğu % 66,7'lik frekans değeriyle *Bittium latreillii* (Payraudeau, 1826)'dir.

Anahtar Sözcükler: Kabuk kullanımı, Anomura Decapoda, Doğu Ege Denizi, Türkiye

Introduction

According to Reese (1969), hermit crabs are typically marine organisms, although some species have physiological and behavioral adaptations for living in various habitats. Most are small decapod crustaceans that occupy empty gastropod shells for protection against predation (Leonard et al., 2001) and depend on gastropod shells throughout their lifespan (Turra and Leita, 2000). These shells act as shelters from biotic factors, including predation (Angel, 2000). Hermit crabs need increasingly larger shells during their life cycle, a fact that keeps them in constant activity searching for suitable shells (Bertness, 1981). The life cycle of hermit crabs depends mostly on the processes that make suitable gastropod shells (Hazlett, 1981). According to the

hypothesis more commonly used, in their natural habitat, hermit crabs inhabit smaller shells than in laboratory experiments (Vance, 1972; Scully, 1979). Hermit crabs seem to select among the available empty gastropod shells the most suitable one for their size and shape (Koutsoubas et al., 1993). Studies on the utilization of gastropod shells by hermit crabs have been conducted by many authors (Fotheringham, 1976; Bertness, 1980, 1981, 1982; Blackstone, 1985, 1989; Gherardi and Vannini, 1989; Hazlett, 1989, 1990, 1992; Gherardi, 1991; Manjón-Cabeza and García Raso, 1999).

Previous studies on the decapod fauna of the Aegean Sea can be considered preliminary work. Of the 35 anomuran species listed by Koukouras et al. (1992), 21 were identified as hermit crabs, as were 21 of the 36

*Corresponding author: asuatates@yahoo.com

anomuran species reported by Kocataş and Katağan (2003). The only study on the occupation of gastropoda shells by hermit crabs was carried out by Koutsoubas et al. (1993) in the Greek waters of the northern Aegean Sea. To the best of our knowledge, the present study is the first to document hermit crab use of gastropod shells along the Turkish coast of the Aegean Sea.

Materials and Methods

In present study, the sublittoral zones of the region between Saros Bay in the north and Marmaris Bay in the south of the Turkish Aegean Sea coast (between lat 40° 35' 50" N, long 26° 22' 40" E and lat 36° 47' 30" N, long 28° 37' 00" E) were chosen as the investigation area (Figure 1).

Samples of hermit crabs were collected by beam-trawl, dredge, or grab from a total of 52 stations in July, August, and September 2000, at depths between 5 and 109 m. The sediment was composed mostly of *Posidonia oceanica* (L.) Delile meadows. The biotopes of the stations can be characterized as follows: 2 of the stations (38 and 48) were fine gravel; 5 of the stations (10, 14, 18, 23, and 52) were sandy; 5 of the stations (11, 26, 29, 32, and 35) were muddy; 9 of the stations (4, 13, 20, 25, 28, 30, 41, 44, and 46) were sandy-muddy; 10 of the stations (16, 17, 33, 34, 36, 37, 39, 43, 49, and 50) were covered by photophilic algae; and 21 of the stations (1, 2, 3, 5, 6, 7, 8, 9, 12, 15, 19, 21, 22, 24, 27, 31, 40, 42, 45, 47, and 51) were covered by *Posidonia oceanica* (L.) Delile meadows. Distribution of the hermit crab species at the 52 stations is shown in Table 1.

The hermit crab and gastropod species were identified based on the works of Ingle (1993), Falciai and Minervini (1996), and Pope and Goto (1991), using the European Register of Marine Species (ERMS) nomenclature (2003).

Results

We documented 12 hermit crab species inhabiting 16 different gastropod shell species (Table 2).

The gastropod families, Barleidae, Cerithiidae, Muricidae, Nassariidae, Rissoidae, Trochidae, and Turritellidae were inhabited by hermit crabs. Frequencies (Fr) of gastropod shells occupied by hermit crabs are shown in Figure 2. The main gastropod species found in the studied area and used by hermit crabs was *Bittium*



Figure 1. Location of the study area showing sampling stations.

latreillii, which was inhabited by 66.7% of the collected hermit crabs (8 different species). This species was followed by *Cerithium vulgatum* (Fr = 41.7%) inhabited by 5 different hermit crabs; *Cerithium protractum*, *Gibbula fanulum*, *Pollia dorbignyi*, and *Turritella communis* (Fr = 16.7%) by 2 hermit crabs; and *Bela nebula*, *Gibbula ardens*, *Gibbula albida*, *Jujubinus striatus*, *Jujubinus exasperatus*, *Muricopsis cristata*, *Nassarius incrassatus*, *Ocenebra aciculata*, *Pusilina* sp., and *Rissoa auriscalptum* (Fr = 8.33%) by 1 hermit crab.

Pagurus anachoretus occupied shells of 8 different gastropod species and had the highest inhabitation frequency (IF = 37.5%) of gastropod shells, followed by *Pagurus prideauxi* (IF = 31.25%). *Pagurus cuanensis* had an IF of 25.0%, and *Anapagurus petiti* and *Pagurus forbesii* had an IF of 18.8%. The IF of *Diogenes pugilator*, *Pagurus alatus*, and *Pagurus chevreuxi* were

Table 1. Distribution of hermit crabs at the sampling stations.

Hermit crab species	Stations
<i>Dardanus arrosor</i> (Herbst, 1796)	34
<i>Diogenes pugilator</i> (Roux, 1829)	34, 35
<i>Paguristes eremita</i> (Linnaeus, 1767)	15, 22, 49
<i>Paguristes syrtensis</i> De Saint Laurent, 1971	6, 15, 19, 34, 36, 39, 40, 41, 42, 43, 44, 45, 47, 48
<i>Anapagurus laevis</i> (Bell, 1845)	13, 28
<i>Anapagurus petiti</i> Dechancé & Forest, 1962	2,3, 4, 18, 28, 31, 32, 44, 51
<i>Pagurus alatus</i> Fabricius, 1775	10, 19, 23, 24, 29, 36, 42, 43, 44, 48
<i>Pagurus anachoretus</i> Risso, 1827	2, 8, 12, 14, 15, 20, 22, 25, 27, 30, 33, 36, 37, 38, 44, 45, 47
<i>Pagurus chevreuxi</i> (Bouvier, 1896)	21, 23, 25, 26, 33, 37, 39, 40, 42, 43, 45, 51, 52
<i>Pagurus cuanensis</i> Bell, 1845	1, 4, 5, 7, 9, 11, 15, 16, 17, 23, 24, 29, 33, 34, 38, 40, 44, 45, 47, 51, 52
<i>Pagurus forbesii</i> Bell, 1845	7, 10, 13, 34, 45, 50
<i>Pagurus prideaux</i> Leach, 1815	19, 26, 38, 45, 46, 49, 52

Table 2. Gastropod shells used by 12 species of hermit crabs collected in the Aegean Sea (1. *Bela nebula* (Montagu, 1803), 2. *Bittium latreilli* (Payraudeau, 1826), 3. *Cerithium vulgatum* (Bruguière, 1792), 4. *C. protractum* Ant. Bivona, in and Bivona 1838, 5. *Gibbula fanulum* (Gmelin, 1791), 6. *G. ardens* (von Salis, 1793), 7. *G. albida* (Gmelin, 1791), 8. *Jujubinus striatus* (Linnaeus, 1767), 9. *J. exasperatus* (Pennant, 1777), 10. *Muricopsis cristata* (Brocchi, 1814), 11. *Nassarius incrassatus* (Ström, 1768), 12. *Ocenebra aciculata* (Lamarck 1822), 13. *Pollia dorbignyi* (Payraudeau, 1826), 14. *Pusilina* sp., 15. *Rissoa auriscalpium* (Linnaeus, 1758), 16. *Turritella communis* Risso, 1826).

Hermit crab species	Gastropod shell species															
	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16
<i>Dardanus arrosor</i>		+														
<i>Diogenes pugilator</i>			+		+											
<i>Paguristes eremita</i>			+													
<i>Paguristes syrtensis</i>		+														
<i>Anapagurus laevis</i>		+														
<i>Anapagurus petiti</i>				+									+		+	
<i>Pagurus alatus</i>		+														
<i>Pagurus anachoretus</i>		+		+		+		+	+	+						+
<i>Pagurus chevreuxi</i>		+			+											
<i>Pagurus cuanensis</i>		+	+										+			
<i>Pagurus forbesii</i>			+								+	+				+
<i>Pagurus prideauxi</i>	+	+	+					+						+		




Figure 2. Gastropod shells used by hermit crabs and inhabitation frequencies.

12.5%, whereas *Dardanus arrosor*, *Paguristes eremita*, *P. syrtensis*, and *Anapagurus laevis* had the lowest IF, 6.23%. Inhabitation frequencies (IF) of gastropod shells by hermit crabs are given in Figure 3.

IF = Inhabitation Frequency (%)

Discussion

In the study area, 12 hermit crab species inhabited 16 different gastropod shells in the Aegean Sea. The majority

of hermit crabs recorded in the study area inhabited the biotope of *Posidonia oceanica* meadows. The hermit crab species most observed on *Posidonia oceanica* beds in the study area was *Pagurus cuanensis*.

The gastropod species occupied by the highest number of hermit crab species was *Bittium latreillii*, with a frequency value of 66.7%, whereas the hermit crab species that inhabited the highest number of gastropod species was *Pagurus anachoretus*, with an IF of 37.5%. Koutsoubas et al. (1993) reported a total of 11 hermit crab species (*Anapagurus laevis*, *Cestopagurus timidus*, *Clibanarius erythropus*, *Dardanus calidus*, *Diogenes pugilator*, *Paguristes eremita*, *Pagurus anachoretus*, *Pagurus cuanensis*, *Pagurus excavatus*, *Pagurus forbesii*, and *Pagurus prideauxi*) inhabited 56 different gastropod shell species collected from 2 to 300 m. The species *C. timidus* (Roux, 1830), *C. erythropus* (Latreille, 1818), *D. calidus* (Risso, 1827), and *P. excavatus* (Herbst, 1791), reported by Koutsoubas et al. (1993), were not recorded in the present study. On the other hand, the hermit crabs, *D. arrosor*, *P. syrtensis*, *A. petiti*, *P. alatus*, and *P. chevreuxi*, reported in the present study were not recorded by Koutsoubas et al. (1993).

According to Kocataş and Katağan (2003), the hermit crab community of the Turkish coasts of the Aegean Sea is composed of 21 species. The present taxocoenosis in this area was controlled by *Pagurus cuanensis*, recorded from 21 stations of the 52 different stations, with 94 specimens. Although *Pagurus cuanensis*




Figure 3. Hermit crab species inhabiting gastropod shell species and inhabitation frequencies.

was the most observed species in the area, it occupied only 3 different gastropod species. Similar studies related to the utilization of gastropod shells by hermit crab populations have been carried out previously in the Mediterranean. Manjón-Cabeza and García Raso (1999) studied the gastropod shell utilization by hermit crabs *Diogenes pugilator*, *Paguristes eremite*, and *Pagurus forbesii* along the Atlantic Ocean shores of Spain (Cadiz) and reported that these hermit crab species used many different gastropod shells, with *Paguristes eremite* being the largest species inhabiting heavier gastropod shell species belonging to the family Muricidae. The same authors reported that *Diogenes pugilator* and *Pagurus forbesii* inhabit smaller turrids. Also in this study, the gastropod families used by *Diogenes pugilator*, *Paguristes eremite*, and *Pagurus forbesii* in the area were Cerithiidae, Muricidae, Nassariidae, and Turriteliidae. In addition, in the present work, relationships between some morphological parameters of hermit crab species

(i.e. relative growth, cephalothoracic shield length, maximum width, cheliped length, and male/female ratio) and gastropod shells (i.e. aperture widths) were not analyzed.

In summary, 12 of the most common hermit crab species of the Turkish coast of the Aegean Sea have been found to occupy 16 different gastropod shell species. Although the species compositions of gastropods and hermit crabs reported previously differ from one study to another, in general, a great number of hermit crab species use a relatively limited number of gastropoda species in particular areas.

Acknowledgments

The authors would like to thank Dr. Bilal Öztürk (Ege University, Fisheries Faculty) for his help in identifying shells.

References

- Angel, J.E. 2000. Effects of shell fit on the biology of the hermit crab *Pagurus longicarpus* (Say). J. Exp. Mar. Biol. Ecol., 243: 169-184.
- Bertness, M.D. 1980. Shell preference and utilization patterns in littoral hermit crabs of the Bay of Panama. J. Exp. Biol. Ecol., 48: 1-16.
- Bertness, M.D. 1981. Conflicting advantages in resource utilization: The hermit crab housing dilemma. Am. Nat., 118: 432-437.
- Bertness, M.D. 1982. Shell utilization, predation pressure and thermal stress in Panamanian hermit crabs: an interoceanic comparison. J. Exp. Mar. Biol. Ecol., 64: 159-187.
- Blackstone, N.W. 1985. The effects of shell size and shape on growth and form in the hermit crab *Pagurus longicarpus*. Biol. Bull., 171: 379-390.
- Blackstone, N.W. 1989. Size, shell-living and carcinization in geographic populations of a hermit crab, *Pagurus hirsutiussculus*. J. Zool., 217: 477-790.
- European register of marine species, 2003. <http://www.erms.biol.soton.ac.uk/>
- Falciai, L. and Minervini, R. 1996. Guide des Homards, Crabes, Longoustes, Crevettes et Autres Crustacés Décapodes d'Europe. Delachaux et Niestle SA, Lausanne-Paris.
- Fotheringham, N. 1976. Population consequences of shell utilization by hermit crabs. Ecology, 57: 570-578.
- Gherardi, F. 1991. Relative growth, population structure and shell utilization of the hermit crab *Clibanarius erythropus* in the Mediterranean. Obelia, 17: 181-196.
- Gherardi, F. and Vannini, M. 1989. Field observations on activity and clustering in two intertidal hermit crabs, *Clibanarius virescens* and *Calcinus laevimanus* (Decapoda, Anomura). Mar. Behav. Physiol., 14: 145-159.
- Hazlett, B.A. 1981. The behavioral ecology of hermit crab. Ann. Rev. Ecol. Syst., 12: 1-22.
- Hazlett, B.A. 1989. Shell exchanges in the hermit crab *Calcinus tibicen*. Animal Behav., 37: 104-111.
- Hazlett, B.A. 1990. Shell exchange in Hawaiian hermit crabs. Pacific Sci., 44: 401-406.
- Hazlett, B.A. 1992. The effect of past experience on the size of shells selected by hermit crabs. Animal Behav., 44: 203-205.
- Ingle, R. 1993. Hermit crabs of the Northeastern Atlantic Ocean and Mediterranean Sea. An Illustrated key. Nat. Hist. Mus. Publ., Chapman and Hall. London.
- Kocataş, A. and Katağan, T. 2003. The Decapod Crustacean fauna of the Turkish Seas. Zool. in the Middle East, 29: 63-74.
- Koukouras, A., Dounas, C., Turkay, M. and Voultziadou, E. 1992. Decapod Crustacean fauna of the Aegean Sea: New information, check list, affinities. Senckenbergiana marit., 22: 217-244.
- Koutsoubas, D., Labadariou, N. and Koukouras, A. 1993. Gastropod shells inhabited by Anomura Decapoda in the North Aegean Sea. Bios, 1: 247-249.
- Leonard, M., Gainess, K.H. and Sandoval, C.M. 2001. Gastropod shell distribution and factors affecting their utilization by marine hermit crabs in Bahia Kino, Sonora, Mexico, Aquatic Sciences Meeting, Albuquerque.

- Manjón-Cabeza, M.E. and García Raso, J.E. 1999. Shell utilization by the hermit crabs *Diogenes pugilator* (Roux, 1829), *Paguristes eremita* (Linnaeus, 1767) and *Pagurus forbesii* Bell, 1845 (Crustacea: Decapoda: Anomura), in a shallow-water community from Southern Spain. *Bull. Mar. Scien.*, 65: 391-405.
- Pope, G.T. and Goto, Y. 1991. European seashells. Vol. I Polyplacophora, Caudofoveata, Solenogastrea and Gastropoda. Verlag Christa Hemmen. Germany, 352 pp.
- Reese, E.S. 1969. Behavioral adaptations of intertidal hermit crabs. *Am. Zool.*, 9: 343-355.
- Turra, A. and Leite, F.P.P. 2000. Shell utilization patterns of a tropical rocky intertidal hermit crab assemblage: I. The case of Grande Beach, J. *Crustacean Biol.*, 21: 393-406.
- Scully, E.P. 1979. The effects of gastropod shell availability and habitat characteristics on shell utilization by the intertidal hermit crab *Pagurus longicarpus* Say. *Jr. Exp. Biol. Ecol.*, 37: 139-152.
- Vance, R.R. 1972. Competition and mechanisms of coexistence in three sympatric species of intertidal hermit crabs. *Ecology*, 53: 1062-1074.