

1-1-2013

A note on unmixed ideals of Veronese bi-type

MONICA LA BARBIERA

Follow this and additional works at: <https://journals.tubitak.gov.tr/math>



Part of the [Mathematics Commons](#)

Recommended Citation

BARBIERA, MONICA LA (2013) "A note on unmixed ideals of Veronese bi-type," *Turkish Journal of Mathematics*: Vol. 37: No. 1, Article 1. <https://doi.org/10.3906/mat-1101-23>

Available at: <https://journals.tubitak.gov.tr/math/vol37/iss1/1>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Mathematics by an authorized editor of TÜBİTAK Academic Journals. For more information, please contact academic.publications@tubitak.gov.tr.

A note on unmixed ideals of Veronese bi-type

Monica LA BARBIERA*

University of Messina, Department of Mathematics Viale Ferdinando Stagno d'Alcontres, 31,
98166 Messina, Italy

Received: 06.01.2011 • Accepted: 12.04.2011 • Published Online: 17.12.2012 • Printed: 14.01.2013

Abstract: We classify the unmixed ideals of Veronese bi-type and in some cases we give a description of their associated prime ideals.

Key words: Unmixed ideals. Veronese bi-type ideals

1. Introduction

Let $R = K[X_1, \dots, X_n; Y_1, \dots, Y_m]$ be a polynomial ring in two sets of variables over a field K . In recent papers, monomial ideals of R are introduced and their connection to bipartite complete graphs is studied ([4], [6]). In this paper we study a class of monomial ideals of R , so-called Veronese bi-type ideals. They are an extension of the ideals of Veronese type ([5]) in a polynomial ring in two sets of variables. More precisely, the ideals of Veronese bi-type are monomial ideals of R generated in the same degree: $L_{q,s} = \sum_{k+r=q} I_{k,s} J_{r,s}$, with $k, r \geq 1$, where $I_{k,s}$ is the Veronese-type ideal generated on degree k by the set $\{X_1^{a_{i_1}} \cdots X_n^{a_{i_n}} \mid \sum_{j=1}^n a_{i_j} = k, 0 \leq a_{i_j} \leq s, s \in \{1, \dots, k\}\}$ and $J_{r,s}$ is the Veronese-type ideal generated on degree r by the set $\{Y_1^{b_{i_1}} \cdots Y_m^{b_{i_m}} \mid \sum_{j=1}^m b_{i_j} = r, 0 \leq b_{i_j} \leq s, s \in \{1, \dots, r\}\}$ ([2], [3]). For $s = 2$ the Veronese bi-type ideals are the ideals associated to bipartite graphs with loops ([2]).

In this paper some properties of these class of monomial ideals are discussed. In particular, our aim is to classify the unmixed Veronese bi-type ideals.

Establishing whenever an ideal is unmixed in general is a difficult problem because it is necessary to know all its associated prime ideals. In [8] equidimensional and unmixed ideals of Veronese type are characterized. Now we are able to classify the unmixed Veronese bi-type ideals and in some cases we can give a description of the associated prime ideals.

This paper is organized as follows. In Section 1, unmixed ideals of Veronese bi-type are classified and the generalized ideals associated to the walks of special bipartite graphs, described by the Veronese bi-type ideals $L_{q,2} = \sum_{k+r=q} I_{k,2} J_{r,2}$, are considered. In Section 2, the toric ideal $I(L_{q,s})$ of the monomial subring $K[L_{q,s}] \subset R$ is studied. Let $L_{q,s} = (f_1, \dots, f_p)$ and $K[L_{q,s}]$ be the K -algebra spanned by f_1, \dots, f_p . There is a graded epimorphism of K -algebras: $\varphi : S = K[T_1, \dots, T_p] \rightarrow K[L_{q,s}]$ induced by $\varphi(T_i) = f_i$, where S is a polynomial ring graded by $\deg(T_i) = \deg(f_i)$. Let $I(L_{q,s})$ be the toric ideal of $K[L_{q,s}]$, that is the kernel of φ .

*Correspondence: monicalb@dipmat.unime.it

2000 AMS Mathematics Subject Classification: 05B35, 13P10, 13C15.

We are able to prove that $I(L_{q,s})$ has a quadratic Groebner basis and as a consequence the K -algebra $K[L_{q,s}]$ is Koszul. In order to formulate these results we have to recall the notion of sortability ([5]), and we apply it to the monomial ideals $L_{q,s}$.

2. Unmixed ideals of Veronese bi-type

Let $R = K[X_1, \dots, X_n; Y_1, \dots, Y_m]$ be the polynomial ring over a field K in two sets of variables with each $\deg X_i = 1$, $\deg Y_j = 1$, for all $i = 1, \dots, n$, $j = 1, \dots, m$.

We define the *ideals of Veronese bi-type* of degree q as the monomial ideals of R

$$L_{q,s} = \sum_{r+k=q} I_{k,s} J_{r,s}, \quad r, k \geq 1,$$

where $I_{k,s}$ is the ideal of Veronese-type of degree k in the variables X_1, \dots, X_n and $J_{r,s}$ is the ideal of Veronese-type of degree r in the variables Y_1, \dots, Y_m .

$L_{q,s}$ is not trivial for $2 \leq q \leq s(n+m) - 1$, $s \leq q$.

Remark 2.1 In general, $I_{k,s} \subseteq I_k$, where I_k is the Veronese ideal of degree k generated by all the monomials in the variables X_1, \dots, X_n of degree k ([6]).

One has $I_{k,s} = I_k$ for any $k \leq s$. If $s = 1$, $I_{k,1}$ is the square-free Veronese ideal of degree k generated by all the square-free monomials in the variables X_1, \dots, X_n of degree k . Similar considerations hold for $J_{r,s} \subset K[Y_1, \dots, Y_m]$.

Example 2.2 Let $R = K[X_1, X_2; Y_1, Y_2]$ be a polynomial ring.

- 1) $L_{2,2} = I_{1,2} J_{1,2} = I_1 J_1 = (X_1 Y_1, X_1 Y_2, X_2 Y_1, X_2 Y_2)$;
- 2) $L_{4,2} = I_{3,2} J_{1,2} + I_{1,2} J_{3,2} + I_{2,2} J_{2,2} = I_{3,2} J_1 + I_1 J_{3,2} + I_2 J_2 = (X_1^2 X_2 Y_1, X_1^2 X_2 Y_2, X_1 X_2^2 Y_1, X_1 X_2^2 Y_2, X_1 Y_1^2 Y_2, X_2 Y_1^2 Y_2, X_1 Y_1 Y_2^2, X_2 Y_1 Y_2^2, X_1^2 Y_1^2, X_1^2 Y_1 Y_2, X_1^2 Y_2^2, X_2^2 Y_1^2, X_2^2 Y_2^2, X_2^2 Y_1 Y_2, X_1 X_2 Y_1^2, X_1 X_2 Y_2^2, X_1 X_2 Y_1 Y_2)$.

In this section we classify the unmixed Veronese bi-type ideals. First, we recall some preliminary notions.

Definition 2.3 Let $G(L_{q,s})$ be the unique minimal set of monic monomial generators of $L_{q,s}$. A *vertex cover* of $L_{q,s}$ is a subset W of $\{X_1, \dots, X_n; Y_1, \dots, Y_m\}$ such that each $u \in G(L_{q,s})$ is divided by some variables of W . Such a vertex cover W is called *minimal* if no proper subset of W is a vertex cover.

Denote by $h(L_{q,s})$ the minimal cardinality of the vertex covers of $L_{q,s}$.

Definition 2.4 A monomial ideal is said to be *unmixed* if all its minimal vertex covers have the same cardinality.

Remark 2.5 We recall the one-to-one correspondence between the minimal vertex covers of an ideal and its minimal primes. Hence \wp is a minimal prime ideal of $L_{q,s}$ if and only if $\wp = (\mathcal{A})$ for some minimal vertex cover \mathcal{A} of $L_{q,s}$.

Now we are able to classify the unmixed Veronese bi-type ideals and in some cases we can give a description of the associated prime ideals.

Proposition 2.6 Let $R = K[X_1, \dots, X_n; Y_1, \dots, Y_m]$. Let $2 \leq q < s(n+m) - i$ for $i = 1, \dots, s-1$. $L_{q,s}$ is unmixed if and only if $n = m$.

Proof By the structure of $G(L_{q,s})$ the minimal vertex covers of $L_{q,s}$ are $W_1 = \{X_1, \dots, X_n\}$, $W_2 = \{Y_1, \dots, Y_m\}$. The minimal cardinality of the vertex covers of $L_{q,s}$ is $h(L_{q,s}) = \min\{n, m\}$. Hence all the minimal vertex covers have the same cardinality if and only if $n = m$. \square

Example 2.7 $R = K[X_1, X_2; Y_1, Y_2]$

$$L_{3,2} = (X_1^2 Y_1, X_1^2 Y_2, X_1 X_2 Y_1, X_1 X_2 Y_2, X_2^2 Y_1, X_2^2 Y_2, X_1 Y_1^2, X_1 Y_1 Y_2, X_1 Y_2^2, X_2 Y_1^2, X_2 Y_1 Y_2, X_2 Y_2^2).$$

The minimal vertex covers are: $W_1 = \{X_1, X_2\}$; $W_2 = \{Y_1, Y_2\}$.

$$h(L_{3,2}) = |W_1| = |W_2| = 2 \Rightarrow L_{3,2} \text{ is unmixed.}$$

Proposition 2.8 Let $R = K[X_1, \dots, X_n; Y_1, \dots, Y_m]$. If $q = s(n+m) - i$ for $i = 1, \dots, s-1$, then $L_{q,s}$ is unmixed.

Proof One has $W_i = \{X_i\}$, $i = 1, \dots, n$, $W_j = \{Y_j\}$, $j = 1, \dots, m$, are the minimal vertex covers of $L_{q,s}$ by construction. \square

Example 2.9 $R = K[X_1, X_2; Y_1, Y_2]$,

$$L_{11,3} = (X_1^3 X_2^3 Y_1^3 Y_2^2, X_1^3 X_2^3 Y_1^2 Y_2^3, X_1^3 X_2^2 Y_1^3 Y_2^3, X_1^2 X_2^3 Y_1^3 Y_2^3).$$

The minimal vertex covers are:

$$W_1 = \{X_1\}; W_2 = \{X_2\}; W_3 = \{Y_1\}; W_4 = \{Y_2\}.$$

$$h(L_{11,3}) = |W_i| = 1 \text{ for all } i = 1, 2, 3, 4 \Rightarrow L_{11,3} \text{ is unmixed.}$$

Let $\mathcal{A} \subseteq \{1, 2, \dots, n+m\}$, where $n+m$ is the number of the variables of the polynomial ring R . For a subset \mathcal{A} we denote by $\mathcal{P}_{\mathcal{A}}$ the prime ideal of R generated by the variables whose index is in \mathcal{A} .

Theorem 2.10 Let $R = K[X_1, \dots, X_n; Y_1, \dots, Y_m]$, $L_{q,s} \subset R$.

$$\mathcal{P}_{\mathcal{A}} \in \text{Ass}_R(L_{q,s}) \Leftrightarrow |\mathcal{A}| \leq i+1,$$

for $i = s(n+m) - q$, $i = 1, \dots, s-1$.

Proof In the following replace the set of variables $\{X_1, \dots, X_n\}$ with $\{z_1, \dots, z_n\}$ and $\{Y_1, \dots, Y_m\}$ with $\{z_{n+1}, \dots, z_{n+m}\}$.

Assume that $\mathcal{P}_{\mathcal{A}} \in \text{Ass}_R(L_{q,s})$. Then there exists a monomial $f \notin L_{q,s}$ such that $L_{q,s} : f = \mathcal{P}_{\mathcal{A}}$. Now we show that we can choose such a monomial f of degree $q-1$ such that $L_{q,s} : f = \mathcal{P}_{\mathcal{A}}$. Suppose that $f \notin L_{q,s}$, $L_{q,s} : f = \mathcal{P}_{\mathcal{A}}$, $\deg(f) \geq q$ and $f = z_1^{a_1} \dots z_{n+m}^{a_{n+m}}$. Then there exists $j_0 \in \{1, 2, \dots, n+m\}$ such that $a_{j_0} > s$. Since $L_{q,s} : f = \mathcal{P}_{\mathcal{A}}$, we have $z_i f \in L_{q,s}$ for all $i \in \mathcal{A}$ and $z_i f \notin L_{q,s}$ for all $i \notin \mathcal{A}$. Moreover, for all $i \in \mathcal{A}$ there exists a monomial $u_i \in G(L_{q,s})$ such that $u_i | (z_i f)$. Being $f \notin L_{q,s}$, this fact means that, for all $i \in \mathcal{A}$, the variable z_i appears in u_i with exponent $a_i + 1$. Therefore $a_i < s$ for all $i \in \mathcal{A}$. It follows that $j_0 \notin \mathcal{A}$. Now we claim that: I) $\bar{f} = f/z_{j_0} \notin L_{q,s}$ and II) $L_{q,s} : \bar{f} = \mathcal{P}_{\mathcal{A}}$.

The first fact follows from that $f \notin L_{q,s}$ and $a_{j_0} - 1 \geq s$. For the second assertion we proceed as follows. $L_{q,s} : \bar{f} \subseteq L_{q,s} : f$ because \bar{f} divides f . Then $L_{q,s} : \bar{f} \subseteq \mathcal{P}_{\mathcal{A}}$, being $\mathcal{P}_{\mathcal{A}} = L_{q,s} : f$. Moreover, since $a_{j_0} - 1 \geq s$

then u_i divides $z_i f / z_{j_0}$ for all $i \in \mathcal{A}$, then $z_i \in L_{q,s} : (f/z_{j_0})$ for all $i \in \mathcal{A}$. Hence $\mathcal{P}_{\mathcal{A}} \subseteq L_{q,s} : (f/x_{j_0})$. It follows the other inclusion $\mathcal{P}_{\mathcal{A}} \subseteq L_{q,s} : \bar{f}$. Hence $\mathcal{P}_{\mathcal{A}} = L_{q,s} : \bar{f}$. After a finite number of these reductions, we find $f \notin L_{q,s}$ of degree $q-1$ such that $\mathcal{P}_{\mathcal{A}} = L_{q,s} : f$. From this fact follows that $fz_i \in L_{q,s}$ for all $i \in \mathcal{A}$ and $fz_i \notin L_{q,s}$ for all $i \notin \mathcal{A}$. In particular $a_i + 1 \leq s$ for all $i \in \mathcal{A}$, and $a_i \leq s$ for all $i \notin \mathcal{A}$. Then $a_i = s$ for all $i \notin \mathcal{A}$. Therefore $f = \prod_{i \in \mathcal{A}} z_i^{a_i} \prod_{i \notin \mathcal{A}} z_i^s$ with $0 \leq a_i < s$ for all $i \in \mathcal{A}$. We have $\deg(\prod_{i \notin \mathcal{A}} z_i^s) = s(n+m-|\mathcal{A}|) = t$. Then we obtain: $s(n+m) \geq (\sum_{i \in \mathcal{A}} a_i + 1) + t = \sum_{i \in \mathcal{A}} a_i + |\mathcal{A}| + t = \sum_{i \in \mathcal{A}} a_i + t + |\mathcal{A}| = \deg(f) + |\mathcal{A}| = q-1 + |\mathcal{A}|$.

Conversely, let $|\mathcal{A}| \leq i+1$, for $i = s(n+m) - q, i = 1, \dots, s-1$, that is $|\mathcal{A}| \leq s(n+m) - q + 1$. Moreover, in these hypotheses one has $s(n+m-|\mathcal{A}|) \leq q-1$. In fact, $s(n+m-|\mathcal{A}|) \leq s(n+m) - i - 1$; then $s|\mathcal{A}| \geq i+1$ that is true for $i = 1, \dots, s-1$. Being $q = s(n+m) - i$ for $i = 1, \dots, s-1$, then by the definition of $L_{q,s}$ it follows that for any monomial $u \in G(L_{q,s})$ there exists an integer $j \in \mathcal{A}$ such that z_j divides u . Therefore $L_{q,s} \subset \mathcal{P}_{\mathcal{A}}$. The condition $s(n+m) \geq q-1 + |\mathcal{A}|$ implies that $(s-1)|\mathcal{A}| + s(n+m-|\mathcal{A}|) \geq q-1$, which together with $s(n+m-|\mathcal{A}|) \leq q-1$ shows that there exists an integer $c_i < s$, for all $i \in \mathcal{A}$ such that $c_i|\mathcal{A}| + s(n+m-|\mathcal{A}|) = q-1$. Then the monomial $f = \prod_{i \in \mathcal{A}} z_i^{c_i} \prod_{i \notin \mathcal{A}} z_i^s$ has degree $q-1$. Hence $f \notin L_{q,s}$ and as a consequence $\mathcal{P}_{\mathcal{A}} \subseteq L_{q,s} : f$. Now we prove that $\mathcal{P}_{\mathcal{A}} = L_{q,s} : f$. Assume that $\mathcal{P}_{\mathcal{A}}$ is a proper subset of $L_{q,s} : f$. Then there exists a monomial f' , in the variables z_i with $i \notin \mathcal{A}$, of degree at least 1 such that $ff' \in L_{q,s}$. This means that there exists a monomial $u = z_1^{a_1} \dots z_{n+m}^{a_{n+m}} \in G(L_{q,s})$ such that u divides ff' . Therefore $a_i \leq c_i$ for any $i \in \mathcal{A}$ because $f' \in K[z_i | i \notin \mathcal{A}]$. It follows that $q = \deg(u) = \sum_{i=1}^{n+m} a_i \leq \sum_{i \in \mathcal{A}} c_i + s(n+m-|\mathcal{A}|) = \deg(f) = q-1$, which is a contradiction. Hence $\mathcal{P}_{\mathcal{A}}$ is not a proper subset of $L_{q,s} : f$, but $\mathcal{P}_{\mathcal{A}} = L_{q,s} : f$. This equality means that $\mathcal{P}_{\mathcal{A}} \in \text{Ass}_R(L_{q,s})$. \square

Example 2.11 $R = K[X_1, X_2; Y_1, Y_2]$,

$$L_{15,4} = (X_1^4 X_2^4 Y_1^4 Y_2^3, X_1^4 X_2^4 Y_1^3 Y_2^4, X_1^4 X_2^3 Y_1^4 Y_2^4, X_1^3 X_2^4 Y_1^4 Y_2^4).$$

By Theorem 2.10 $\text{Ass}_R(L_{15,4}) = \{(X_1), (X_2), (Y_1), (Y_2), (X_1, X_2), (X_1, Y_1), (X_1, Y_2), (X_2, Y_1), (X_2, Y_2), (Y_1, Y_2)\}$.

As an application, we observe that the ideals of Veronese bi-type can be associated to graphs with loops. In fact for $s = 2$, the ideals $L_{q,s}$ are associated to the walks of length $q-1$ of the strong quasi-bipartite graphs with loops ([2]).

Definition 2.12 A graph G with loops is a strong quasi-bipartite if all vertices of V_1 are joined to all vertices of V_2 and for each vertex of V there is a loop.

Definition 2.13 Let G be a strong quasi-bipartite graph on the vertex set $V = \{v_1, \dots, v_n\}$. A walk of length q in G is an alternating sequence $w = \{v_{i_0}, l_{i_1}, v_{i_1}, l_{i_2}, \dots, v_{i_{q-1}}, l_{i_q}, v_{i_q}\}$, where v_{i_j} is a vertex of G and $l_{i_j} = \{v_{i_{j-1}}, v_{i_j}\}$ is the edge joining $v_{i_{j-1}}$ and v_{i_j} or a loop if $v_{i_{j-1}} = v_{i_j}$, $1 \leq i_1 \leq i_2 \leq \dots \leq i_q \leq n$.

Example 2.14 Let G be a strong quasi-bipartite graph on vertices $\{x_1, x_2; y_1, y_2\}$. A walk of length 2 is

$$w = \{x_1, l_1, x_1, l_2, y_1\},$$


where $l_1 = \{x_1, x_1\}$ is the loop on x_1 and $l_2 = \{x_1, y_1\}$ is the edge joining x_1 and y_1 .

Let G be a strong quasi-bipartite graph on vertex set $\{x_1, \dots, x_n; y_1, \dots, y_m\}$.

The *generalized ideal* $I_q(G)$ associated to G is the ideal of the polynomial ring $R = K[X_1, \dots, X_n; Y_1, \dots, Y_m]$ generated by the monomials of degree q corresponding to the walks of length $q - 1$. Hence the generalized ideal $I_q(G)$ is generated by all monomials of degree $q \geq 3$ corresponding to the walks of length $q - 1$ and the variables in each generator of $I_q(G)$ have at most degree 2. Therefore:

$$I_q(G) = L_{q,2} = \sum_{k+r=q} I_{k,2} J_{r,2}, \text{ for } q \geq 3 \text{ ([2])}.$$

Example 2.15 Let $R = K[X_1, X_2; Y_1, Y_2]$ be a polynomial ring over a field K and G be the strong quasi-bipartite graph on vertices x_1, x_2, y_1, y_2 :



$$I_3(G) = I_1 J_2 + I_2 J_1 = (X_1 Y_1 Y_2, X_2 Y_1 Y_2, X_1 Y_1^2, X_2 Y_1^2, X_1 Y_2^2, X_2 Y_2^2, X_1 X_2 Y_1, X_1 X_2 Y_2, X_1^2 Y_1, X_1^2 Y_2, X_2^2 Y_1, X_2^2 Y_2).$$

$$I_4(G) = I_{3,2} J_1 + I_1 J_{3,2} + I_2 J_2 = (X_1^2 X_2 Y_1, X_1^2 X_2 Y_2, X_1 X_2^2 Y_1, X_1 X_2^2 Y_2, X_1 Y_1^2 Y_2, X_2 Y_1^2 Y_2, X_1 Y_1 Y_2^2, X_2 Y_1 Y_2^2, X_1^2 Y_1^2, X_1^2 Y_1 Y_2, X_2^2 Y_1^2, X_2^2 Y_1 Y_2, X_2^2 Y_2^2, X_2^2 Y_2 Y_1, X_1 X_2 Y_1^2, X_1 X_2 Y_2^2, X_1 X_2 Y_1 Y_2).$$

The following result classifies the ideals $I_q(G)$ that are unmixed.

Proposition 2.16 Let $R = K[X_1, \dots, X_n; Y_1, \dots, Y_m]$.

- 1) If $2 \leq q < 2(n + m) - 1$ then $L_{q,2}$ is unmixed if and only if $n = m$.
- 2) If $q = 2(n + m) - 1$, then $L_{q,2}$ is unmixed.

Proof By Propositions 2.6 and 2.8. □

We consider the case $q \geq 3$. In fact, for $q = s = 2$, the ideal $L_{2,2} = (\{X_i Y_j | i = 1, \dots, n, j = 1, \dots, m\})$ doesn't describe the edge ideal $I(G) = I_2(G)$ of a strong quasi-bipartite graph, but it is the edge ideal of a complete bipartite graph (with no loops) on the vertex set $\{x_1, \dots, x_n; y_1, \dots, y_m\}$. Then $L_{2,2}$ satisfies the characterization of unmixed bipartite graphs given in [7].

3. The toric ideal of $K[L_{q,s}]$

Let $R = K[X_1, \dots, X_n; Y_1, \dots, Y_m]$ and $L_{q,s} = (f_1, \dots, f_p)$ be the ideal of Veronese bi-type. The monomial subring of R spanned by $F = \{f_1, \dots, f_p\}$ is the K -algebra $K[L_{q,s}] = K[F] = K[f_1, \dots, f_p]$. Note that $K[F]$ is a graded algebra with the grading $K[F]_i = K[F] \cap R_i$. There is a graded epimorphism of K -algebras: $\varphi : S = K[T_1, \dots, T_p] \rightarrow K[L_{q,s}]$ induced by $\varphi(T_i) = f_i$, where S is a polynomial ring graded by $\deg(T_i) = \deg(f_i)$. Note that the map φ is given by $\varphi(h(T_1, \dots, T_p)) = h(f_1, \dots, f_p)$ for all $h \in S$.

Let $I(L_{q,s})$ be the kernel of the K -algebra epimorphism, called the *toric ideal* of $K[L_{q,s}]$. It is known that the toric ideal of a monomial K -algebra is a graded prime ideal generated by a finite set of binomials.

Now we prove that $I(L_{q,s})$ has a quadratic Groebner basis. In order to formulate this result we have to recall the notion of sortability, introduced in [5], and we apply it to the monomial ideal $L_{q,s}$.

Let $A = K[z_1, \dots, z_t]$ be a polynomial ring and L be a monomial ideal of A generated in degree q . Let B be the set of the exponent vectors of the monomials of $G(L)$. If $u = (u_1, \dots, u_t), v = (v_1, \dots, v_t) \in B$, then $\underline{z}^u = \prod_i z_i^{u_i}, \underline{z}^v = \prod_i z_i^{v_i} \in L$. We write $\underline{z}^u \underline{z}^v = z_{i_1} \cdots z_{i_{2q}}$ with $i_1 \leq i_2 \leq \dots \leq i_{2q}$. Then we set $\underline{z}^{u'} = \prod_{j=1}^q z_{2j-1}$ and $\underline{z}^{v'} = \prod_{j=1}^q z_{2j}$. This defines a map

$$\text{sort} : B \times B \rightarrow M_q \times M_q, \quad (u, v) \mapsto (u', v'),$$

where M_q is the set of all integer vectors (a_1, \dots, a_t) such that $\sum_{i=1}^t a_i = q$.

The set B is called *sortable* if $\text{Im}(\text{sort}) \subseteq B \times B$.

The ideal L is called *sortable* if the set of exponent vectors of the monomials of $G(L)$ is sortable. In other words, let $\underline{z}^u, \underline{z}^v \in L$, then L is said sortable if $\underline{z}^{u'}, \underline{z}^{v'} \in L$, where $(u', v') = \text{sort}(u, v)$.

Theorem 3.1 *Let $R = K[X_1, \dots, X_n; Y_1, \dots, Y_m]$. $L_{q,s}$ is sortable.*

Proof Let $L_{q,s} = (\{X_1^{a_1} \cdots X_n^{a_n} Y_1^{b_1} \cdots Y_m^{b_m} \mid \sum_{i=1}^n a_i + \sum_{j=1}^m b_j = q, 0 \leq a_i, b_j \leq s\})$ and B be the set of the exponent vectors of the monomials of $G(L_{q,s})$.

Let $f_i = X_1^{a_1} \cdots X_n^{a_n} Y_1^{b_1} \cdots Y_m^{b_m}, f_j = X_1^{c_1} \cdots X_n^{c_n} Y_1^{d_1} \cdots Y_m^{d_m} \in G(L_{q,s})$, then $u = (a_1, \dots, a_n; b_1, \dots, b_m), v = (c_1, \dots, c_n; d_1, \dots, d_m) \in B$. One has that $f_i f_j = \underbrace{X_1 \cdots X_1}_{a_1+c_1\text{-times}} \cdots \underbrace{X_n \cdots X_n}_{a_n+c_n\text{-times}} \underbrace{Y_1 \cdots Y_1}_{b_1+d_1\text{-times}} \cdots \underbrace{Y_m \cdots Y_m}_{b_m+d_m\text{-times}}$ is

a monomial of degree $2q$. If one replaces the set of variables $\{X_1, \dots, X_n\}$ with $\{z_1, \dots, z_n\}$ and $\{Y_1, \dots, Y_m\}$ with $\{z_{n+1}, \dots, z_{n+m}\}$, then $f_i f_j = z_{i_1} \cdots z_{i_{2q}}$ with $i_1 \leq \dots \leq i_{2q}$. Then we consider $f'_i = \underline{z}^{u'} = \prod_{l=1}^q z_{2l-1}$ and $f'_j = \underline{z}^{v'} = \prod_{l=1}^q z_{2l}$. We must prove that $f'_i, f'_j \in L_{q,s}$. We have that f'_i is of degree q and we write

$f'_i = \prod_{l=1}^q z_{2l-1} = X_1^{a'_1} \cdots X_n^{a'_n} Y_1^{b'_1} \cdots Y_m^{b'_m}$. If $a_i + c_i$ is even then $a'_i = \frac{a_i+c_i}{2} \leq s$ and if $a_i + c_i$ is odd then $a'_i = \frac{a_i+c_i+1}{2} < s$. Similarly, if $b_j + d_j$ is even then $b'_j = \frac{b_j+d_j}{2} \leq s$. If $b_j + d_j$ is odd then $b'_j = \frac{b_j+d_j+1}{2} < s$.

Moreover, because f'_i is of degree q and there exist $a'_i \neq 0, b'_j \neq 0$ with $0 \leq a'_i, b'_j \leq q$ for all i, j , then $X_1^{a'_1} \cdots X_n^{a'_n} \neq X_i^q$ and $Y_1^{b'_1} \cdots Y_m^{b'_m} \neq Y_j^q$. It follows that $X_1^{a'_1} \cdots X_n^{a'_n} \in I_{k,s}$ and $Y_1^{b'_1} \cdots Y_m^{b'_m} \in J_{r,s}$ with $k+r=q$. Hence $f'_i \in L_{q,s}$. In the same way the argument holds for f'_j . Hence $L_{q,s}$ is sortable. \square

Corollary 3.2 *Let $R = K[X_1, \dots, X_n; Y_1, \dots, Y_m]$ and $L_{q,s} \subset R$. Then:*

- 1) $I(L_{q,s})$ has a quadratic Groebner basis.
- 2) $K[L_{q,s}]$ is Koszul.

Proof 1) By Theorem 3.1 and [1](Lemma 5.2).

2) The conclusion follows by 1). \square

References

- [1] Herzog, J., Hibi, T.: Discrete Polymatroids, Journal of Alg. Comb., 16, 239–268 (2002).
- [2] La Barbiera, M.: Normalization of Veronese bi-type ideals, Italian Journal of Pure and Applied Mathematics, 26, 79-92 (2009).

- [3] La Barbiera, M.: On a class of monomial ideals generated by s -sequences, *Mathematical Reports*, 12(62), N.3, 201-216 (2010).
- [4] Restuccia, G., Villarreal, R.H.: On the normality of monomial ideals of mixed products, *Communications in Algebra*, 29(8), 3571-3580 (2001).
- [5] Sturmfels, B.: *Groebner bases and convex polytopes*, Amer. Math. Soc., Providence, RI, 1995.
- [6] Villarreal, R.H.: *Monomial Algebras, Monographs and Textbooks in Pure and Applied Mathematics*, 238, Marcel Dekker, Inc., New York, 2001.
- [7] Villarreal, R.H.: Unmixed bipartite graphs, *Revista Colombiana de Matematicas*, 41, 393-395 (2007).
- [8] Vladoiu, M.: Equidimensional and unmixed ideals of Veronese type, *Communication in Algebra*, 36(9), 3378-3392 (2008).