

1-1-2006

New *Torrenticola* Piersig (Acari: Hydrachnidia: Torrenticolidae) Species for the Turkish Fauna

VLADIMIR PESIC

ORHAN ERMAN

YUNUS ESEN

Follow this and additional works at: <https://journals.tubitak.gov.tr/zoology>



Part of the [Zoology Commons](#)

Recommended Citation

PESIC, VLADIMIR; ERMAN, ORHAN; and ESEN, YUNUS (2006) "New *Torrenticola* Piersig (Acari: Hydrachnidia: Torrenticolidae) Species for the Turkish Fauna," *Turkish Journal of Zoology*: Vol. 30: No. 3, Article 10. Available at: <https://journals.tubitak.gov.tr/zoology/vol30/iss3/10>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Zoology by an authorized editor of TÜBİTAK Academic Journals. For more information, please contact academic.publications@tubitak.gov.tr.

New *Torrenticola* Piersig (Acari: Hydrachnidia: Torrenticolidae) Species for the Turkish Fauna

Vladimir PESIC¹, Orhan ERMAN², Yunus ESEN²

¹Department of Biology, University of Crna Gora, Cetinjski put b.b., 81000 Podgorica, Montenegro, Serbia and Montenegro

²Department of Biology, Science and Arts Faculty, Fırat University, 23119 Elazığ - TURKEY

Received: 29.09.2005

Abstract: In this study, the morphological characteristics, measurements, habitats and distribution of *Torrenticola nana* Di Sabatino & Gerecke, 2003 and *Torrenticola jasmineae* Bader, 1988, which are new records for the Turkish fauna, are presented.

Key Words: Water mites, *Torrenticola*, new records, running water, Turkey

Türkiye Faunası İçin Yeni *Torrenticola* Piersig (Acari: Hydrachnidia: Torrenticolidae) Türleri

Özet: Bu çalışmada, Türkiye faunası için yeni kayıt olarak belirlenen, *Torrenticola nana* Di Sabatino ve Gerecke, 2003 ve *Torrenticola jasmineae* Bader, 1988'in morfolojik özellikleri, ölçümleri, habitatları ve dağılımları verilmiştir.

Anahtar Sözcükler: Su keneleri, *Torrenticola*, Yeni kayıtlar, Akarsu, Türkiye

Introduction

Water mites of the genus *Torrenticola* Piersig have been found in all biogeographic regions, except for Australia and Antarctica (Wiles, 1997). The aim of this paper was to contribute elements for an extended revision of the diversity, distribution, and ecology of torrenticolid water mites in Turkey.

At present, only 6 species of the genus *Torrenticola* Piersig are known from Turkey (Smit, 1995; Özkan, 1982; Turan and Pesic, 2005): *Torrenticola elliptica* (Maglio, 1909), *T. anomala* (Koch, 1837), *T. brevis* (Halbert, 1911), *T. barsica* (Szalay, 1933), *T. ungeri* (Szalay, 1927) and *T. amplexa* (Koenike, 1908).

During a survey of the freshwater fauna of Turkey, several species of the water mite were collected, including 2 species of the genus *Torrenticola* Piersig, which are new for the fauna of Turkey.

Material and Methods


Water mites were collected by hand netting, sorted on the spot from the living material, conserved in Koenike's fluid and dissected as described elsewhere (e.g., Gerecke, 1991). Slide-mounted specimens and material preserved in fluid were lodged in the collection of the first author; further material will be deposited in the collection of the second author.

The composition of the material is given as males/females/deutonymphs and the following abbreviations are used: Cx-1 = first coxae, L = length, P-1 = palp, first segment and W = width. All measurements are given in µm.

Results

Torrenticola nana Di Sabatino & Gerecke, 2003
(Figures 1-2)

* E-mail: oerman@firat.edu.tr



Figures 1-2. *Torrenticola nana* Di Sabatino & Gerecke, 2003; male: 1 = idiosoma, ventral view; 2 = palp. Scale bars = 0.1 mm.

Material examined: Malatya, Darende, Tohma Stream, 03.07.2004, leg. Y. Esen (3/10/0).

Description: Male: Idiosoma (Figure 1) L 659, W 531, dorsal shield L 587, W 400, L/W ratio 1.47; dorsal plate L 544; shoulder plate L/W 186/57, L/W ratio 3.26; frontal plate L/W 112/52, L/W ratio 2.15; shoulder/frontal plate L ratio 1.66; gnathosomal bay L 111, Cx-1 L 231, median L 159, Cx-2+3 median L 141; ratio Cx-1 L/Cx-2+3 median L 1.45, Cx-1 median L/Cx-2+3 median L 1.64; genital field L/W 100/91, L/W ratio 1.1, with roundish anterior angles; ejaculatory complex L 106; distance genital field–excretory pore 89, genital field–caudal body margin 134; gnathosoma ventral L 300, with weakly curved ventral margin and well-

developed rostrum; chelicera L 334; palp (Figure 2) total L 288, dorsal length and relative length (% of total length in parentheses) of palp segments: P-1 31 (10.9), P-2 99 (34.4), P-3 56 (19.4), P-4 83 (28.8), P-5 18(6.3); P-2/P-4 ratio 1.19; laterodistal margin of P-2 protruding in form of an obtuse angle, ventral protuberances of P-4 ending in 2 tips separated by a concavity.

Remarks: Due to the reduced size of the genital field and the relatively long median suture line of Cx-2+3 in males and the complete reduction of the median suture of Cx-2+3 in the females, the Turkish specimens show a general conformity with *Torrenticola nana* Di Sabatino & Gerecke, 2003.

Habitat: Rhithral.

Distribution: Israel (Di Sabatino et al., 2003), Iran (Asadi et al., 2003; Pesic et al., 2004).


New record for the Turkish fauna.

Torrenticola jasmineae Bader, 1988

(Figures 3-8)

Material examined: Malatya, Pütürge, Siro Stream, 08.08.2003, leg. Y. Esen (2/0/0).

Description: Male: Idiosoma (Figure 3) L 756, W 531, dorsal shield (Figure 4) L 659, W 200; dorsal plate L 625 (L/W ratio 3.12); shoulder plate L/W 167/62, L/W ratio 2.69; frontal plate L/W 122/63, L/W ratio 1.94; shoulder/frontal plate L ratio 1.37; gnathosomal bay L



Figures 3-8. *Torrenticola jasmineae* Bader, 1988; male: 3 = idiosoma, ventral view; 4 = dorsal shield; 5 = chelicera; 6 = palp; 7 = gnathosoma and palp; 8 = ejaculatory complex. Scale bars = 0.1 mm.

122, Cx-1 L 256, median L 136, Cx-2+3 median L 84; ratio Cx-1 L/Cx-2+3 median L 1.88, Cx-1 median L/Cx-2+3 median L 3.05; genital field L/W 150/113, L/W ratio 1.33; ejaculatory complex (Figure 8) L 194; distance genital field–excretory pore 200, genital field–caudal body margin 244; gnathosoma (Figure 7) ventral L 334; chelicera (Figure 5) L 356, cheliceral basal segment L 323, cheliceral claw L 38, (basal segment/claw L ratio 8.6); palp (Figures 6-7) total L 340, dorsal length and relative length (% of total length in parentheses) of palp segments: P-1 53 (15.5), P-2 133 (39.1), P-3 52 (39.1), P-4 82 (24.1), P-5 20 (58.3); P-2/P-4 ratio 1.62; ventral seta of P-2 inserted on a well-developed ventral projection.

Remarks: After the original description from Elburz Mountain (Bader, 1988) based on a single male, the only further records of *T. jasmineae*, each based on a single female, were published from Israel (Di Sabatino et al.,

2003) and another one from Iran (Pestic et al., 2004). Since in the original description (Bader, 1988), the key characteristic (basal segment/claw ratio of chelicera) has not been reported. Our attribution of *T. jasmineae* is very provisional, based mainly on the approved non-identity with the alternative species (Di Sabatino et al., 2003). Furthermore, the attribution of the populations from Israel and Iran to the species should be verified by studies on male specimens from these populations and from Elburz Mountain.

Distribution: Iran (Bader, 1988; Pestic et al., 2004), Israel (Di Sabatino et al., 2003).

New record for the Turkish fauna.

Acknowledgements

This study was supported by FUBAP-1000 project, Fırat University, Elazığ, TURKEY.

References

- Asadi, M., Pestic, V. and Saboori, A. 2003. Two interesting water mite species (Acari, Hydrachnidia) from Iran, with redescription of the female of *Atractides arcuatus* Thor, 1914. *Zoology in the Middle East*, 30: 95-100.
- Bader, C. 1988. Die Torrenticolidae (Acari, Hydrachnellae). Eine abklärende Studie über eine schwierige Wassermilben-Familie. *Revue Suisse Zool.*, 95 (1): 87-98.
- Gerecke, R. 1991. Taxonomische, faunistische und ökologische Untersuchungen an Wassermilben (Acari, Actinedida) aus Sizilien unter Berücksichtigung anderer aquatischer Invertebraten. *Lauterbornia*, 7: 1-304.
- Di Sabatino, A., Gerecke, R., Smit, H., Pestic, V. and Panesar, A. 2003. Water mites of the family Torrenticolidae (Acari, Actinedida, Hydrachnidia) from the Eastern Mediterranean region. *Archiv für Hydrobiologie Suppl.*, 139 (3): 1-39.
- Özkan, M. 1982. Wassermilben (Acari, Actinedida) aus der Türkei. *Entomol. Basil.*, 7: 29-60.
- Pestic, V., Saboori, A. and Asadi, M. 2004. Water mites of the genus *Torrenticola* Piersig (Acari: Hydrachnidia, Torrenticolidae) from Iran. *Ann. Limnol.-Int. J. Lim.*, 40 (3): 231-236.
- Smit, H. 1995. New records of water mites from Turkey, with 11 species new for the Turkish fauna (Acari: Hydrachnellae). *Storkia*, 4: 10-15.
- Turan, D. and Pestic, V. 2005. Three water mite species of the genus *Torrenticola* Piersig (Acari, Hydrachnidia) new for the Turkish fauna. *Natura Montenegrina* 3: 33-39.
- Wiles, P. R. 1997. Asian and Oriental Torrenticolidae Piersig, 1902 (Acari: Hydrachnidia: Lebertioidea): a revision of the family and description of new species of *Torrenticola* Piersig and *Pseudotorrenticola* Walter, from Southeast Asia. *Journal of Natural History*, 31: 191-236.