

1-1-2009

## Melatonin administration increases antioxidant enzymes activities and reduces lipid peroxidation in the rainbow trout (*Oncorhynchus mykiss*, Walbaum) erythrocytes

İLHAMİ GÜLÇİN

ŞÜKRÜ BEYDEMİR

OLCAY HİSAR

EKREM KÖKSAL

RUSSEL J. REITER

Follow this and additional works at: <https://journals.tubitak.gov.tr/veterinary>



Part of the [Animal Sciences Commons](#), and the [Veterinary Medicine Commons](#)

### Recommended Citation

GÜLÇİN, İLHAMİ; BEYDEMİR, ŞÜKRÜ; HİSAR, OLCAY; KÖKSAL, EKREM; and REITER, RUSSEL J. (2009) "Melatonin administration increases antioxidant enzymes activities and reduces lipid peroxidation in the rainbow trout (*Oncorhynchus mykiss*, Walbaum) erythrocytes," *Turkish Journal of Veterinary & Animal Sciences*: Vol. 33: No. 3, Article 11. <https://doi.org/10.3906/vet-0803-1>  
Available at: <https://journals.tubitak.gov.tr/veterinary/vol33/iss3/11>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Veterinary & Animal Sciences by an authorized editor of TÜBİTAK Academic Journals. For more information, please contact [academic.publications@tubitak.gov.tr](mailto:academic.publications@tubitak.gov.tr).

## Melatonin administration increases antioxidant enzymes activities and reduces lipid peroxidation in the rainbow trout (*Oncorhynchus mykiss*, Walbaum) erythrocytes

İlhami GÜLÇİN<sup>1</sup>, Şükrü BEYDEMİR<sup>1</sup>, Olcay HİSAR<sup>2\*</sup>, Ekrem KÖKSAL<sup>1</sup>, Russel J. REITER<sup>3</sup>

<sup>1</sup>Department of Chemistry, Faculty of Sciences, Atatürk University, 25240 Erzurum - TURKEY

<sup>2</sup>Department of Aquaculture, Faculty of Agriculture, Atatürk University, 25240 Erzurum - TURKEY

<sup>3</sup>Department of Cellular and Structural Biology, Health Science Centre at San Antonio, University of Texas, Mail Code 7762, 7703 Floyd Curl Drive, San Antonio, TX 78229-3900-USA

Received: 03.03.2008

**Abstract:** Effects of melatonin on the levels of antioxidant enzymes including catalase (CAT), peroxidase (POD), glutathione reductase (GRd), and malondialdehyde (MDA), a lipid peroxidation marker, in erythrocytes of rainbow trout (*Oncorhynchus mykiss*) were investigated. The results showed that melatonin statistically activated the CAT, POD, and GRd activity ( $P < 0.05$ ). Moreover, MDA levels were significantly decreased as a result of melatonin treatment ( $P < 0.01$ ).

**Key Words:** Rainbow trout; melatonin; antioxidant enzyme; lipid peroxidation

### Melatonin uygulaması gökkuşuğu alabalık (*Oncorhynchus mykiss*, Walbaum) eritrositlerinde antioksidan enzim aktivitelerini artırırken lipit peroksidasyonunu azaltmaktadır

**Özet:** Çalışmada, melatoninin gökkuşuğu alabalık eritrositlerindeki katalaz (CAT), peroksidaz (POD) ve glutatyon redüktaz (GRd) gibi antioksidan enzimler ile lipit peroksidasyonu göstergesi olan malondialdehit (MDA) üzerine etkileri araştırıldı. Sonuçta, melatonin uygulamasının CAT, POD ve GRd aktivitelerini istatistiksel olarak artırdığı ( $P < 0,05$ ) gösterildi. Ayrıca melatonin uygulamasının bir sonucu olarak MDA seviyesinin önemli derecede azaldığı ( $P < 0,01$ ) belirlenmiştir.

**Anahtar Sözcükler:** Gökkuşuğu alabalığı; melatonin; antioksidan enzimler; lipit peroksidasyonu

#### Introduction

Melatonin is primarily produced by photoreceptor cells, which act as a photoneuroendocrine transducer and secrete melatonin into the blood, in the pineal

gland (1). It is involved in many physiological processes, such as seasonal reproduction, activity rhythms, and sleep/wake cycles (2), in osmoregulation and/or stress adaptation in seawater (3). Moreover, it

\* E-mail: ohisar@atauni.edu.tr

is considered as a powerful free radical scavenger and likely to be a general promoter of anti-oxidative mechanisms and a potential antioxidant in vitro and in vivo (4). Some of its interactions with other hormones such as cortisol and arginine vasotocin (hormones closely associated with electrolyte balance) have been reported in teleosts (5,6).

The antioxidant defense (AD) system of organisms provides a means of dealing with oxidative stress and includes several enzymes and vitamins. A primary role of the AD system is protecting cellular components from damage induced by reactive oxygen species (ROS) (7). The ROS include free radicals, such as superoxide anion radicals ( $O_2^{\cdot-}$ ), hydroxyl radicals ( $OH\cdot$ ) and non-free radicals such as singlet oxygen ( $^1O_2$ ) and hydrogen peroxide ( $H_2O_2$ ). Also, excessive generation of ROS induced by various stimuli and exceeding the antioxidant capacity of the organism leads to a variety of pathophysiological processes. Exposure of organisms to pro-oxidant attack can increase antioxidant defenses, e.g. by increasing synthesis of antioxidant enzymes (8). If antioxidant defenses are effective in detoxifying ROS, then no harmful consequence will result in the tissues. However, if the ROS attack is severe, antioxidant defense systems may be overwhelmed, resulting in inhibition of antioxidant enzymes and oxidative damage to lipids, proteins, DNA, and other key molecules. Such processes may in turn provoke alterations in molecular and membrane structures and functions, leading to cell and tissue damage (9). Increased concentration of lipid peroxidation and DNA damage has been used as indicators of ROS-derived damage in biological systems (10).

Most cellular antioxidants can be regarded as part of the innate nonspecific immune system (11). Like other vertebrates, fish possess an AD system both enzymatic and non-enzymatic. The more relevant AD enzymes consist of glutathione reductase (GRd), superoxide dismutase (SOD), catalase (CAT) and peroxidase (POD), and non-enzymatic defenses include vitamin C,  $\alpha$ -tocopherol, and vitamin A (12).

Although, the influence of melatonin administration on fish reproductive parameters has been extensively investigated (13,14), studies related to the effects of this application on the inhibition or activation of enzymes playing a key role in antioxidant

processes are unfortunately scarce in fish. Therefore, the aim of the present study was to determine the effects of melatonin on the activity of endogenous antioxidant enzymes such as CAT, POD, and GRd, and the levels of MDA as a marker of peroxidized lipid concentrations from rainbow trout (*Oncorhynchus mykiss*) erythrocytes in vivo.

## Materials and Methods

### In vivo experiment

Fish treatments were conducted according to the Applied Research Ethics National Association (15). In the experiment, fish (mean weight:  $200 \pm 20$  g) were divided into 2 groups as control and melatonin. Each included 20 fish. Fish were anaesthetized in water containing tricaine methanesulphonate (MS-222, 1/10,000), and 0.5 ml blood samples from the control group and the melatonin group were collected from the caudal sinus using heparinized syringes and placed into a heparinized vacutainer to determine initial activities of CAT, POD, and GSSG-Px as well as the levels of total MDA. Subsequently, melatonin was dissolved in a small amount of ethanol and then diluted with teleost saline (20 mg  $Na_2CO_3$ /100 ml of 0.6% NaCl) until the desired final concentrations were obtained. Physiological saline was injected to all fish in the control group intraperitoneally (i.p.). For the melatonin group, 10 mg/kg (fish body weight) melatonin was injected i.p. Blood samples were taken from the different fish at 1, 3, and 5 h after injection from the control and the melatonin group ( $n = 5$ ). All blood samples were centrifuged at 2500 rpm for 15 min, then the plasma was removed by drip, the erythrocyte pellet was washed with 0.16 M KCl 3 times and the supernatant was discarded. One volume from the resulting erythrocyte pellet was hemolyzed in 5 volumes of ice-water. Ghost and intact cells were then removed by high-speed centrifugation (20000 rpm for 30 min, Heraeus Sepatech, Suprafuge). As a result, a hemolysate was prepared. Hemoglobin (Hb) concentration in hemolysate was determined by the cyanmethemoglobin method. All of the above-mentioned procedures were carried out at  $+4$  °C.

### Measurement of antioxidant enzyme activities

Catalase activity was assayed by monitoring the decrease in absorbance at 240 nm due to  $H_2O_2$

consumption (16). The reaction mixture contained 0.1 ml hemolysate, 0.5 ml 10 mM H<sub>2</sub>O<sub>2</sub> and 0.9 ml 30 mM potassium phosphate buffer (pH 7.3) and reduction in absorbance was recorded at 240 nm for 60 s. The enzyme activity is expressed as mmoles H<sub>2</sub>O<sub>2</sub> decomposed by using the H<sub>2</sub>O<sub>2</sub> extinction coefficient 36 mM<sup>-1</sup>.cm<sup>-1</sup>.

Peroxidase activity was measured by the method of Shannon et al. (17). For this purpose, 0.1 ml hemolysate were taken to 1.7 ml 0.05 M potassium phosphate buffer (pH 7.3). Then 0.5 ml ABTS were added. The reaction was started by the addition of 0.2 ml 0.2 M H<sub>2</sub>O<sub>2</sub>. Change in absorbance at 470 nm was recorded for 2 min at intervals of 15 s. The enzyme activity is expressed as enzyme unit min<sup>-1</sup>.gHb<sup>-1</sup>.

GRd activity was estimated by the method of Goldberg and Spooner (18). To 0.1 ml hemolysate 2.5 ml 120 mM potassium phosphate buffer (pH 7.3), 0.1 ml 0.015 mM EDTA, and 0.1 ml 0.065 mM oxidized glutathione were added. After 5 min, 0.05 ml 9.6 mM NADH were added and mixed. The absorbance was recorded at 340 nm at 15 s intervals. The enzyme activity is expressed as nmole NADH oxidized min<sup>-1</sup>.gHb<sup>-1</sup> using the molar extinction coefficient 6.23 mM<sup>-1</sup>.cm<sup>-1</sup>.

#### Measurement of MDA level

MDA levels of rainbow trout erythrocytes were estimated according to the modified method of Hunter et al. (19). To 0.5 ml hemolysate, 0.5 ml Tris/HCl buffer (50 mM, pH 7.4) was added followed by further mixing and incubation at room temperature for 10 min; 1.0 ml of 0.75% thiobarbituric acid in 2 M Na<sub>2</sub>SO<sub>4</sub> was added and then the mixture was heated to 100 °C for 45 min. After cooling, 1.0 ml of 70% TCA was added, the mixture was vortexed, and then centrifuged at 1000 × g for 10 min. The absorbance of supernatant was determined at 530 nm. Total thiobarbituric acid-reactive materials are expressed as MDA, using a molar extinction coefficient for MDA of 1.56 × 10<sup>5</sup> cm<sup>-1</sup>.M<sup>-1</sup>.

#### Statistical analyses

The obtained data were analyzed by Student's t-test or one way analysis of variance (ANOVA), followed by Tukey multiple range test to determine significant differences among means at the α: 0.05 level.

## Results

In the present study, it was observed that there were no differences (P > 0.05) between the enzyme activities determined in the control group at 0 (initial), 1, 3, and 5 h for each enzyme. However, as can be seen in the Table, the differences between the mean values of enzyme activities determined at different times were statistically significant (P < 0.05, using the ANOVA). Melatonin increased the CAT activities in rainbow trout at 1 h as compared to the control values. These differences were found to be significant (P < 0.001, using Student's t-test). However, CAT activity returned to the control level at 5 h after treatment (P > 0.05). Like CAT activity, POD and GRd activities were increased after melatonin injection. Maximal increases of POD and GRd activities by the fish dose of melatonin occurred within 1 h after melatonin administration (P < 0.01, using Student's t-test). POD and GRd activities returned to the control level at 3 h and 5 h, respectively after treatment (P > 0.05) (Figure 1).

It was observed that there were no differences (P > 0.05) between the MDA levels determined in the control group at 0 (initial), 1, 3, and 5 h. The decreased concentrations of MDA in the rainbow trout plasma were found after treatment with a 10 mg.kg<sup>-1</sup> dose of melatonin. Melatonin especially reduces lipid peroxidation after 5 h (Figure 2). This difference was found to be statistically significant (P < 0.01, using Student's t-test).

Table. The effect of melatonin on the activities of antioxidant enzymes from rainbow trout (*Oncorhynchus mykiss*) erythrocytes (CAT: catalase, POD: peroxidase GRd: glutathione reductase, EU: enzyme unit, Hb: haemoglobin).

Time (h)	CAT (EU/gHb)	POD (EU/gHb)	GRd (EU/gHb)
0	462.8 ± 30.3 <sup>a</sup>	1339.2 ± 243.3 <sup>a</sup>	11.9 ± 2.3 <sup>a</sup>
1	564.4 ± 28.5 <sup>b</sup>	1809.3 ± 164.6 <sup>b</sup>	26.9 ± 4.4 <sup>b</sup>
3	616.2 ± 48.1 <sup>b</sup>	1532.5 ± 269.4 <sup>a</sup>	16.4 ± 2.4 <sup>a</sup>
5	421.3 ± 69.5 <sup>a</sup>	1140.0 ± 376.3 <sup>a</sup>	12.9 ± 3.2 <sup>a</sup>

Values are means ± SD of 5 fish

Means in a column with different superscripts differ significantly (P < 0.05)




Figure 1. Changes of antioxidant enzymes activities (EU/gHb) in rainbow trout (*Oncorhynchus mykiss*) erythrocytes after the administration pharmacological doses of melatonin ( $10 \text{ mg kg}^{-1}$ ) (differences relative to time 0 h: \* $P \leq 0.05$ ; \*\* $P \leq 0.01$ ; \*\*\* $P \leq 0.001$  versus control using Student's t-test).




Figure 2. Levels of malondialdehyde (MDA) in rainbow trout (*Oncorhynchus mykiss*) plasma after the administration pharmacological doses of melatonin ( $10 \text{ mg kg}^{-1}$ ) (differences relative to time 0 h: \* $P \leq 0.05$ ; \*\* $P \leq 0.01$ ; \*\*\* $P \leq 0.001$  versus control using Student's t-test).

## Discussion

Melatonin and its metabolites are known to prevent oxidative damage to the cell membrane, cytosolic organelles, and nuclear and mitochondrial DNA by electron donation (20,21). These effects can occur by at least 2 mechanisms. In one case, melatonin itself exerts direct antioxidant effects via free-radical scavenging and/or by inhibiting the generation of these toxic reactants. Additionally, melatonin alters the activities of antioxidant enzymes in mammals (22).

A variety of studies have demonstrated an improvement not only in oxidative stress parameters measured in plasma, serum, liver, kidney, and brain, but also in scavenger enzyme activities in other experimental models after melatonin is administered (23). In recent studies, the effect of melatonin on carbonic anhydrase enzyme activity in rainbow trout erythrocytes was determined as in vitro and in vivo. It was observed that melatonin did not inhibit the rainbow trout erythrocyte carbonic anhydrase enzyme activity in vitro. On the contrary, the results did show that rainbow trout erythrocyte carbonic anhydrase was inhibited significantly by melatonin after its injection in vivo ( $P < 0.05$ ) (24). Furthermore,

it has been documented that melatonin has an effect on glucose-6-phosphate dehydrogenase from rainbow trout erythrocytes. In vivo studies showed that, though initial glucose-6-phosphate dehydrogenase activity was  $8.33 \text{ EU/gHb}$ , these values fell after the injection of a pharmacological dose of melatonin (25).

Oxidation of membrane lipids can lead to the loss of cellular or organelle membrane integrity, which can eventually result in membrane rigidity, pathological injury, and cell death (26). Salmonid tissues are characterized by high concentrations of polyunsaturated fatty acids compared with most mammalian tissues, and fish may, therefore, be particularly susceptible to lipid peroxidative cellular damage (27). In previous studies, melatonin levels were found to correlate with the total antioxidant capacity of rat (28) and human serum (29) and in all studies using mammals, melatonin characteristically depressed the products of lipid peroxidation.

In conclusion, our study showed that melatonin reduced levels of lipid peroxides and increased the activity of antioxidant enzymes such as CAT, POD, and GRd in rainbow trout. These findings are consistent with other published reports using other species (22,30).

## References

1. Bayarri, M.J., Rol De Lama, M.A., Madrid, J.A., Sánchez-Vázquez, F.J.: Both pineal and lateral eyes are needed to sustain daily circulating melatonin rhythms in sea bass. *Brain Res.*, 2003; 969: 175-182.
2. Reiter, R.J.: Pineal melatonin: cell biology of its synthesis and of its physiological interactions. *Endocr. Rev.*, 1991, 12: 151-180.
3. Folmar, L.C., Dickhoff, W.W.: Evaluation of some physiological parameters as predictive indices of smoltification. *Aquaculture*, 1981; 23: 309-324.
4. Konar, V., Yilmaz, O., Ozturk, A.I., Kirbag, S., Arslan, M.: Antimicrobial and biological effects of bomphos and phomphos on bacterial and yeast cells. *Bioorg. Chem.*, 2000; 28: 214-225.
5. Delahunty, G., Schreck, C.B., de Vlaming, V.L.: Effect of pinealectomy, reproductive state, feeding regime and photoperiod on plasma cortisol in goldfish. *Am. Zool.*, 1977; 17: 873-376.
6. Kulczykowska, E.: Arginine vasotocin-melatonin interactions in fish: A hypothesis. *Rev. Fish Biol. Fisher.*, 1995; 5: 96-102.
7. Kelly, K.A., Havrilla, C.M., Brady, T.C., Abramo, K.H., Levin, E.D.: Oxidative stress in toxicology: established mammalian and emerging piscine model systems. *Environ. Health Perspect.*, 1998; 106: 375-384.
8. Halliwell, B., Gutteridge, J.: *Free Radicals in Biology and Medicine*. Oxford University Press, Oxford. 1999.
9. Reznick, A.Z., Packer, L., Sen, C.K.: Strategies to assess oxidative stress. In: Reznick, A.Z., Packer, L., Sen, C.K., Holloszy, J.O., Jackson, M.J., Eds., *Oxidative Stress in Skeletal Muscle*. Birkhäuser Verlag, Basel, Switzerland. 1998; 43-58.
10. Halliwell, B., Chirico, S.: Lipid peroxidation: its mechanism, measurement, and significance. *Am. J. Clin. Nutr.*, 1993; 57: 715-725.
11. Kang, C.R., Sweeter, S., Boylan, L.M., Spallholz, J.E.: Oxygen toxicity, biological defense systems and immunity: a historical perspective. *J. Nutr. Immunol.*, 1994; 3: 51-84.
12. Filho, D.W., Giulivi, C., Boveris, A.: Antioxidant defences in marine fish I: Teleosts. *Comp. Biochem. Physiol. C Comp. Pharmacol. Toxicol.*, 1993; 106: 409-413.
13. Amano, M., Iigo, M., Ikuta, K., Kitamura, S., Yamada, H., Yamamori, K.: Roles of melatonin in gonadal maturation of underyearling precocious male Masu salmon. *Gen. Comp. Endocrinol.*, 2000; 120: 190-197.
14. Bornestaf, C., Mayer, I., Borg, B.: Melatonin and maturation pace in female three-spined stickleback *Gasterosteus aculeatus*. *Gen. Comp. Endocrinol.*, 2001; 122: 341-348.
15. ARENA (Applied Research Ethics National Association): *Institutional Animal Care and Use Committee Guidebook*. 2nd edn., National Academy of Sciences, Boston. 2002; 121-125.
16. Aebi, H.: Catalase in vitro. *Methods Enzymol.*, 1984; 105: 121-126.
17. Shannon, L.M., Kay, E., Lew, J.Y.: Peroxidase isoenzymes from horseradish roots. I. Isolation and physical properties. *J. Biol. Chem.*, 1996; 241: 2166-2172.
18. Goldberg, D.M., Spooner R.J.: Glutathione reductase. In: Bergmayer, O.H. Ed., *Methods of Enzymatic Analysis*. 3rd edn., Verlag Chemie Deerfield Beach, FL. 1983; 259-265.
19. Hunter, M.I.S, Nlemadim, B.C., Davidson, D.L.W.: Lipid peroxidation products and antioxidant proteins in plasma and cerebrospinal fluid from multiple sclerosis patients. *Neurochem. Res.*, 1985; 10: 1645-1652.
20. Reiter, R.J., Tan, D.X., Qi, W.B., Manchester, L.C., Karbownik, M., Calvo, J.R.: Pharmacology and physiology of melatonin in the reduction of oxidative stress in vivo. *Biol. Signals Recept.*, 2000; 9: 160-171.
21. Rosen, J., Than, N.N., Koch, D., Poeggeler, B., Laatsch, H. Hardeland, R.: Interactions of melatonin and its metabolites with the ABTS cation radical: extension of the radical scavenger cascade and formation of a novel class of oxidation products, C2-substituted 3-indolinones. *J. Pineal Res.*, 2006; 41: 374-381.
22. Reiter, R., Tang, L., Garcia, J.J., Muñoz-Hoyos, A.: Pharmacological actions of melatonin in oxygen radical pathophysiology. *Life Sci.*, 1997; 60: 2255-2271.
23. Tomás-Zapico, C., Coto-Montes, A.: A proposed mechanism to explain the stimulatory effect of melatonin on antioxidative enzymes. *J. Pineal Res.*, 2005; 39: 99-104.
24. Hisar, O., Beydemir, Ş., Gülçin, İ., Aras Hisar, Ş., Yanık, T., Küfrevioğlu, Ö.İ.: The effects of melatonin hormone on carbonic anhydrase enzyme activity in rainbow trout (*Oncorhynchus mykiss*) erythrocytes in vitro and in vivo. *Turk. J. Vet. Anim. Sci.*, 2005; 29: 841-845.
25. Beydemir, Ş., Gülçin, İ., Hisar, O., Küfrevioğlu, Ö.İ.; Yanık, T.: Effect of melatonin on glucose-6-phosphate dehydrogenase from rainbow trout (*Oncorhynchus mykiss*) erythrocytes in vitro and in vivo. *J. Appl. Anim. Res.*, 2005; 28: 65-68.
26. García, J.J., Reiter, R.J., Pié J., Ortiz G.G., Cabrera J., Sáinz R.M., Acuña-Castroviejo, D.: Role of pinoline and melatonin in stabilizing hepatic microsomal membranes against oxidative stress. *J. Bioenerg. Biomembr.*, 1999; 31: 609-616.
27. Winston, G.W., Di Giulio, R.T.: Prooxidant and antioxidant mechanisms in aquatic organisms. *Aquat. Toxicol.*, 1991; 19: 137-161.
28. Benot, S., Goberna, R., Reiter, R.J., Garcia-Mauriño, S., Osuna, C., Guerrero, J.M.: Physiological levels of melatonin contribute to the antioxidant capacity of human serum. *J. Pineal Res.*, 1999; 27: 59-64.
29. Benot, S., Molinero, P., Soutto, M., Goberna, R., Guerrero, J.M.: Circadian variations in the rat serum total antioxidant status: correlation with melatonin levels. *J. Pineal Res.*, 1998; 25: 1-4.
30. Reiter, R.J., Tan, D.X., Cabrera, J., D'Arpa, D., Sainz, R.M., Mayo, J.C., Ramos, S.: The oxidant/antioxidant network: role of melatonin. *Biol. Signals Recept.*, 1999; 8: 56-63.