

1-1-2012

## Value distribution of meromorphic functions and their differences

RANRAN ZHANG

ZONGXUAN CHEN

Follow this and additional works at: <https://journals.tubitak.gov.tr/math>



Part of the [Mathematics Commons](#)

---

### Recommended Citation

ZHANG, RANRAN and CHEN, ZONGXUAN (2012) "Value distribution of meromorphic functions and their differences," *Turkish Journal of Mathematics*: Vol. 36: No. 3, Article 6. <https://doi.org/10.3906/mat-1101-79>

Available at: <https://journals.tubitak.gov.tr/math/vol36/iss3/6>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Mathematics by an authorized editor of TÜBİTAK Academic Journals. For more information, please contact [academic.publications@tubitak.gov.tr](mailto:academic.publications@tubitak.gov.tr).

## Value distribution of meromorphic functions and their differences\*

Ranran Zhang and Zongxuan Chen

### Abstract

Let  $f(z)$  be a transcendental meromorphic function. Results are proved concerning the value distribution of the  $n$ 'th forward difference  $\Delta^n f(z)$ , in terms of Borel exceptional values of  $f(z)$ . The results may be partly viewed as discrete analogues of a classical theorem of Hayman dealing with the possible relationships between Picard exceptional values of  $f(z)$  and its derivatives.

**Key words and phrases:** Complex difference; value distribution; Borel exceptional value

### 1. Introduction and results

Let  $f(z)$  be a meromorphic function in the plane. We assume that the reader is familiar with the basic notions of Nevanlinna's theory (see [10]). We use  $\sigma(f)$  to denote the order of growth of  $f(z)$ ; and  $\lambda(f)$  and  $\lambda(1/f)$  to denote, respectively, the exponents of convergence of zero and pole sequences of  $f(z)$ . Moreover, we use  $\delta(a, f)$  to denote the Nevanlinna deficiency of  $f(z)$ . For a nonzero constant  $c$ , the forward differences  $\Delta^n f$  are defined (see [1]) by

$$\Delta f(z) = f(z + c) - f(z), \Delta^{n+1} f(z) = \Delta^n f(z + c) - \Delta^n f(z), n = 1, 2, \dots$$

Throughout this paper, we denote by  $S(r, f)$  any function satisfying  $S(r, f) = o(T(r, f))$  as  $r \rightarrow \infty$ , possibly outside a set of  $r$  of finite logarithmic measure. A meromorphic function  $\alpha(z)$  is said to be a small function of  $f(z)$ , if  $T(r, \alpha) = S(r, f)$ .

Recently, there is substantial interest in difference analogues of Nevanlinna's theory, as well as difference equations. The papers [1, 2] investigated the zeros of  $\Delta^n f(z)$  under the assumption that  $f(z)$  is of small growth order, and obtained many profound results. These results may be viewed as discrete analogues of the following existing theorem on the zeros of  $f'(z)$ .

**Theorem A [4, 11]** *Let  $f(z)$  be transcendental and meromorphic in the plane with*

$$\liminf_{r \rightarrow \infty} \frac{T(r, f)}{r} = 0.$$

2000 AMS Mathematics Subject Classification: 30D35, 39A10.

\*This project was supported by the National Natural Science Foundation of China (10871076) and by the Tianyuan Fund for Mathematics (11026096).

Then  $f'(z)$  has infinitely many zeros.

Hayman [9] investigated the possible relationships between Picard exceptional values of  $f(z)$  and its derivatives, and obtained the following classical theorem.

**Theorem B [9]** *If  $f(z)$  is transcendental and meromorphic in the plane, then either  $f(z)$  assumes every finite value infinitely often, or every derivative of  $f(z)$  assumes every finite value except possibly zero infinitely often.*

In this paper, we investigate the value distribution of meromorphic functions and their differences. First we observe that for a general meromorphic function, the difference counterpart of Theorem B doesn't exist, see the following Example 1.1.

**Example 1.1** *Let  $f(z) = ze^z/(2\pi i(e^z + 1))$  and  $c = 2\pi i$ . Then*

$$\Delta f(z) = f(z + 2\pi i) - f(z) = \frac{e^z}{e^z + 1}.$$

We see that  $f(z)$  assumes 0 finitely often and  $\Delta f(z)$  cannot assume 1.

Example 1.1 shows that if  $f(z)$  has only one Borel exceptional value, then  $\Delta f(z)$  may not assume some finite nonzero value. In this paper, we prove that if  $f(z)$  has two Borel exceptional values and if  $f(z)$  is not of period  $c$ , then  $\Delta f(z)$  assumes every finite value except possibly zero infinitely often. Actually, we get the following Theorem, which may be partly viewed as discrete analogues of Theorem B.

**Theorem 1.1** *Let  $f(z)$  be a finite order transcendental meromorphic function with two Borel exceptional values  $a, b$ . Let  $c \in C \setminus \{0\}$  and let  $s(z)$  be a nonzero small function of  $f(z)$ . For every positive integer  $n$ , set*

$$F_n(z) = \Delta^n f(z) - s(z).$$

Suppose that one of the following two conditions holds:

- (i)  $a, b \in C$  and  $c, 2c, \dots, nc$  are not periods of  $f(z)$ ;
- (ii)  $a \in C, b = \infty$  and  $\Delta^n f(z) \not\equiv 0$ .

Then  $F_n(z)$  is transcendentially meromorphic and  $\delta(0, F_n) \leq n/(n + 1)$ .

**Remark** *The following Examples, 1.2–1.4, show that Theorem 1.1 is false, if  $f(z)$  has at most one Borel exceptional value. So the requirement “ $f(z)$  has two Borel exceptional values” in Theorem 1.1 cannot be weakened.*

**Example 1.2** *Let  $f(z)$  and  $c$  be as in Example 1.1, and let  $s(z) = 1$ . Then  $f(z)$  has only one Borel exceptional value 0, and*

$$F_1(z) = \Delta f(z) - s(z) = \frac{e^z}{e^z + 1} - 1 = \frac{-1}{e^z + 1}$$

has no zeros.

**Example 1.3** *Let  $f(z) = e^z + z, c = 2\pi i$  and  $s(z) = \pi i$ . Then  $f(z)$  has only one Borel exceptional value  $\infty$ , and*

$$F_1(z) = \Delta f(z) - s(z) = f(z + 2\pi i) - f(z) - \pi i = \pi i$$

has no zeros.

**Example 1.4** Let  $f(z) = \Gamma'(z)/\Gamma(z)$ ,  $c = 1$  and  $s(z) = 1$ . Then  $f(z)$  has no Borel exceptional values, and

$$F_1(z) = \Delta f(z) - s(z) = f(z+1) - f(z) - 1 = \frac{1}{z} - 1$$

has only one zero.

We give the following two corollaries. Corollary 1.1 is obtained directly from Theorem 1.1. Corollary 1.2 cannot be obtained directly from Theorem 1.1, since the condition “ $N(r, 1/(f - a)) + N(r, f) = S(r, f)$ ” in Corollary 1.2 does not imply that  $a$  and  $\infty$  are Borel exceptional values. However, using the same method as in Part II of proof of Theorem 1.1, we can easily prove Corollary 1.2.

**Corollary 1.1** Let  $f(z)$  be a finite order transcendental meromorphic function with two Borel exceptional values, and let  $c \in C \setminus \{0\}$ . For every positive integer  $n$ , if  $\Delta^n f(z) \not\equiv 0$  and  $c, 2c, \dots, nc$  are not periods of  $f(z)$ , then  $\Delta^n f(z)$  assumes every finite value except possibly zero infinitely often.

**Corollary 1.2** Let  $c \in C \setminus \{0\}$  and  $a \in C$ . Let  $f(z)$  be a transcendental meromorphic function of finite order such that

$$N(r, 1/(f - a)) + N(r, f) = S(r, f).$$

For every positive integer  $n$ , if  $\Delta^n f(z) \not\equiv 0$ , then  $\Delta^n f(z)$  assumes every finite value except possibly zero infinitely often.

Next we give the conditions under which  $\Delta^n f(z)$  assumes every finite value (including zero) infinitely often.

**Theorem 1.2** Let  $f(z)$  be a transcendental meromorphic function with  $1 < \sigma(f) < \infty$ . Let  $c \in C \setminus \{0\}$  and  $a \in C$ . Suppose that

$$\max\{\lambda(f - a), \lambda(1/f)\} < \sigma(f) - 1.$$

Then for every positive integer  $n$ ,  $\Delta^n f(z)$  assumes every finite value infinitely often.

## 2. Lemmas for the proofs of theorems

**Lemma 2.1 [6]** Let  $f(z)$  be a nonconstant meromorphic function of finite order, and let  $\eta_1, \eta_2$  be two arbitrary complex numbers. Then

$$m\left(r, \frac{f(z + \eta_1)}{f(z + \eta_2)}\right) = S(r, f).$$

**Lemma 2.2 [7, 8]** Let  $f(z)$  be a nonconstant finite order meromorphic function and let  $c \neq 0$  be an arbitrary complex number. Then

$$\begin{aligned} T(r + |c|, f) &= T(r, f) + S(r, f), \\ N(r + |c|, f) &= N(r, f) + S(r, f). \end{aligned}$$

It is shown in [5, p. 66], that for an arbitrary  $c \neq 0$ , the following inequalities

$$(1 + o(1))T(r - |c|, f(z)) \leq T(r, f(z + c)) \leq (1 + o(1))T(r + |c|, f(z))$$

hold as  $r \rightarrow \infty$  for a general meromorphic function. From the proof we see that the above relations are also true for  $N(r, f(z+c))$ . So by these relations and Lemma 2.2, we get the following lemma.

**Lemma 2.3** *Let  $f(z)$  be a nonconstant finite order meromorphic function and let  $c \neq 0$  be an arbitrary complex number. Then*

$$T(r, f(z+c)) = T(r, f) + S(r, f),$$

$$N(r, f(z+c)) = N(r, f) + S(r, f).$$

**Remark** *Chiang and Feng [3] have obtained some results similar to the above Lemmas 2.1–2.3, and their work is independent from [6, 7, 8].*

**Lemma 2.4 [13]** *Let  $f(z)$  be a transcendental meromorphic function. Let  $P(f)$  be a polynomial in  $f(z)$  of the form*

$$P(f) = a_n(z)f(z)^n + a_{n-1}(z)f(z)^{n-1} + \cdots + a_0(z),$$

where all coefficients  $a_j(z)$  are small functions of  $f(z)$  and  $a_n(z) \neq 0$ . Then

$$T(r, P(f)) = nT(r, f) + S(r, f).$$

**Lemma 2.5 [12]** *Let  $f(z)$  be a nonconstant meromorphic function, and suppose that*

$$\Psi(z) = a_n(z)f(z)^n + a_{n-1}(z)f(z)^{n-1} + \cdots + a_0(z)$$

has small meromorphic coefficients  $a_j(z)$ ,  $a_n(z) \neq 0$ . Then either

$$T(r, f) \leq \overline{N}(r, 1/\Psi) + \overline{N}(r, f) + S(r, f)$$

or

$$\Psi(z) = a_n \left( f + \frac{a_{n-1}}{na_n} \right)^n.$$

Lemma 2.5 is a version of Tumura-Clunie type theorems. Next we will establish a difference analogue of Lemma 2.5. To this end, we introduce some notations. The difference polynomial  $H(z, f)$  is defined by

$$H(z, f) = \sum_{\lambda \in J} a_\lambda(z) \prod_{j=1}^{\tau_\lambda} f(z + \delta_{\lambda,j})^{\mu_{\lambda,j}}, \tag{2.1}$$

where  $J$  is an index set,  $\delta_{\lambda,j}$  are complex constants,  $\mu_{\lambda,j}$  are nonnegative integers, and  $a_\lambda(z) (\neq 0)$  are small meromorphic functions of  $f(z)$ . The maximal total degree of  $H(z, f)$  in  $f(z)$  and the shifts of  $f(z)$  is defined by

$$\deg_f H = \max_{\lambda \in J} \sum_{j=1}^{\tau_\lambda} \mu_{\lambda,j}.$$

For  $l = 0, 1, \dots, \deg_f H$ , we define

$$J_l = \left\{ \lambda \in J \mid \sum_{j=1}^{\tau_\lambda} \mu_{\lambda,j} = l \right\}. \quad (2.2)$$

**Lemma 2.6** *Let  $f(z)$  be a transcendental meromorphic function of finite order such that*

$$N(r, 1/f) + N(r, f) = S(r, f). \quad (2.3)$$

*Suppose that the difference polynomial (2.1) in  $f(z)$  with small meromorphic coefficients is of maximal total degree  $\deg_f H \geq 1$ . If there exist two different integers  $m, k \in \{0, 1, \dots, \deg_f H\}$  such that*

$$\sum_{\lambda \in J_m} a_\lambda(z) \prod_{j=1}^{\tau_\lambda} f(z + \delta_{\lambda,j})^{\mu_{\lambda,j}} \not\equiv 0, \quad \sum_{\lambda \in J_k} a_\lambda(z) \prod_{j=1}^{\tau_\lambda} f(z + \delta_{\lambda,j})^{\mu_{\lambda,j}} \not\equiv 0, \quad (2.4)$$

*where  $J_m, J_k$  are defined by (2.2), then  $H(z, f)$  is transcendentially meromorphic and*

$$T(r, f) \leq \bar{N}(r, 1/H) + S(r, f).$$

**Proof.** Since there exist two different integers  $m, k \in \{0, \dots, \deg_f H\}$  satisfying (2.4), we may assume, without losing generality, that  $m > k$  and

$$\sum_{\lambda \in J_s} a_\lambda(z) \prod_{j=1}^{\tau_\lambda} f(z + \delta_{\lambda,j})^{\mu_{\lambda,j}} \equiv 0$$

for  $s = m + 1, \dots, \deg_f H$ , where  $J_s$  are defined by (2.2). Thus,  $H(z, f)$  takes the form

$$H(z, f) = \sum_{i=0}^m b_i(z) f(z)^i, \quad (2.5)$$

where for  $i = 0, \dots, m$ ,

$$b_i(z) = \sum_{\lambda \in J_i} a_\lambda(z) \prod_{j=1}^{\tau_\lambda} \left( \frac{f(z + \delta_{\lambda,j})}{f(z)} \right)^{\mu_{\lambda,j}}, \quad J_i = \left\{ \lambda \in J \mid \sum_{j=1}^{\tau_\lambda} \mu_{\lambda,j} = i \right\}.$$

In particular,  $b_m(z) \not\equiv 0$  and  $b_k(z) \not\equiv 0$ .

Since the coefficients  $a_\lambda(z)$  of  $H(z, f)$  are small functions of  $f(z)$ , we have  $T(r, a_\lambda) = S(r, f)$ . So by Lemma 2.1, we get

$$m(r, b_i) = S(r, f)$$

for  $i = 0, 1, \dots, m$ . Moreover, by (2.3) and Lemma 2.3, we have

$$N(r, b_i) \leq \sum_{\lambda \in J_i} \left( N(r, a_\lambda) + \sum_{j=1}^{\tau_\lambda} \mu_{\lambda,j} \left( N(r, f(z + \delta_{\lambda,j})) + N(r, 1/f) \right) \right) + O(1) = S(r, f).$$

So

$$T(r, b_i) = S(r, f) \tag{2.6}$$

for  $i = 0, \dots, m$ . By (2.5), (2.6),  $b_m(z) \not\equiv 0$  and Lemma 2.4, we see that  $H(z, f)$  is transcendently meromorphic.

Applying Lemma 2.5 to (2.5), we get either

$$T(r, f) \leq \overline{N}(r, 1/H) + S(r, f) \tag{2.7}$$

or

$$H(z, f) = b_m \left( f + \frac{b_{m-1}}{mb_m} \right)^m. \tag{2.8}$$

If (2.7) holds, there is nothing to prove. So in the following discussion, we assume that (2.8) holds. First we affirm that  $b_{m-1} \not\equiv 0$ . Otherwise, (2.8) yields

$$H(z, f) = b_m(z)f(z)^m,$$

and so by (2.5), we have

$$\sum_{i=0}^{m-1} b_i(z)f(z)^i \equiv 0.$$

By this equality and Lemma 2.4, we get  $b_i(z) \equiv 0$  for  $i = 0, \dots, m-1$ . This contradicts  $b_k(z) \not\equiv 0$ ,  $k < m$ . Thus,  $b_{m-1} \not\equiv 0$ , and by (2.6) we have

$$T\left(r, \frac{b_{m-1}}{mb_m}\right) = S(r, f).$$

Applying the second main theorem for small target functions and noting (2.3), we get

$$\begin{aligned} T(r, f) &\leq \overline{N}\left(r, \frac{1}{f}\right) + \overline{N}(r, f) + \overline{N}\left(r, \frac{1}{f + \frac{b_{m-1}}{mb_m}}\right) + S(r, f) \\ &= \overline{N}\left(r, \frac{1}{f + \frac{b_{m-1}}{mb_m}}\right) + S(r, f). \end{aligned}$$

Moreover, by (2.8) and  $T(r, b_m) = S(r, f)$ , we have

$$\overline{N}\left(r, \frac{1}{f + \frac{b_{m-1}}{mb_m}}\right) = \overline{N}\left(r, \frac{1}{H}\right) + S(r, f).$$

Therefore

$$T(r, f) \leq \overline{N}(r, 1/H) + S(r, f).$$

□

### 3. Proof of Theorem 1.1

**Part I** We assume that the condition (i) in Theorem 1.1 holds. Set  $g(z) = 1/(f(z) - b)$ . Then  $g(z)$  has two Borel exceptional values  $1/(a - b), \infty$ . Let  $1/(a - b) = d$ . By Hadamard's factorization theory,  $g(z)$  takes the form

$$g(z) = h(z)e^{p(z)} + d, \tag{3.1}$$

where  $p(z)$  is a polynomial and  $h(z)$  is a meromorphic function satisfying  $\sigma(h) < \sigma(g)$ . So  $\sigma(g) = \deg p \geq 1$ , and  $g(z)$  is of regular growth, i.e.,

$$\limsup_{r \rightarrow \infty} \frac{\log T(r, g)}{\log r} = \liminf_{r \rightarrow \infty} \frac{\log T(r, g)}{\log r} = \sigma(g). \tag{3.2}$$

By (3.2) and the fact that  $\infty$  is a Borel exceptional value of  $g(z)$ , we get

$$N(r, g) = S(r, g). \tag{3.3}$$

Observe that

$$\Delta^n f(z) = \sum_{j=0}^n \binom{n}{j} (-1)^{n-j} f(z + jc), \tag{3.4}$$

$$\sum_{j=0}^n \binom{n}{j} (-1)^{n-j} = (1 - 1)^n = 0, \tag{3.5}$$

where  $\binom{n}{j}$  are the binomial coefficients. Substituting  $f(z) = 1/g(z) + b$  into (3.4) and noting (3.5), we get

$$\Delta^n f(z) = \sum_{j=0}^n \binom{n}{j} (-1)^{n-j} \left( \frac{1}{g(z + jc)} + b \right) = \sum_{j=0}^n \binom{n}{j} (-1)^{n-j} \frac{1}{g(z + jc)}.$$

So

$$F_n(z) = \Delta^n f(z) - s(z) = \frac{E_1(z) - s(z) \prod_{j=0}^n g(z + jc)}{\prod_{j=0}^n g(z + jc)}, \tag{3.6}$$

where

$$E_1(z) = \sum_{j=0}^n \binom{n}{j} (-1)^{n-j} \prod_{\substack{i=0 \\ i \neq j}}^n g(z + ic). \tag{3.7}$$

Let  $g_1(z) = g(z) - d$ . Then  $g_1(z)$  has two Borel exceptional values  $0, \infty$ , and  $g_1(z)$  is of regular growth.

So

$$N(r, g_1) = S(r, g_1), \quad N(r, 1/g_1) = S(r, g_1).$$

Set

$$E_2(z) = E_1(z) - s(z) \prod_{j=0}^n g(z + jc).$$



Substituting (3.7) into  $E_2(z)$  and then replacing  $g(z)$  by  $g(z) = g_1(z) + d$ , we get

$$E_2(z) = \sum_{j=0}^n \binom{n}{j} (-1)^{n-j} \prod_{\substack{i=0 \\ i \neq j}}^n (g_1(z + ic) + d) - s(z) \prod_{j=0}^n (g_1(z + jc) + d). \quad (3.8)$$

By calculation, we obtain

$$-s(z) \prod_{j=0}^n (g_1(z + jc) + d) = -s(z) \prod_{j=0}^n g_1(z + jc) + P(z, g_1) - s(z)d^{n+1}, \quad (3.9)$$

and for  $j = 0, \dots, n$ ,

$$\prod_{\substack{i=0 \\ i \neq j}}^n (g_1(z + ic) + d) = P_j(z, g_1) + d^n, \quad (3.10)$$

where  $P(z, g_1)$  and  $P_j(z, g_1)$  are difference polynomials in  $g_1(z)$  and its shifts such that the degree of every term in  $P(z, g_1)$  and  $P_j(z, g_1)$  is at most  $n$  and at least 1. By (3.8)–(3.10) and noting (3.5), we get

$$E_2(z) = -s(z) \prod_{j=0}^n g_1(z + jc) + P(z, g_1) + \sum_{j=0}^n \binom{n}{j} (-1)^{n-j} P_j(z, g_1) - s(z)d^{n+1}.$$

Since  $s(z) \neq 0$  and  $d = 1/(a - b) \neq 0$ , we have  $-s(z) \prod_{j=0}^n g_1(z + jc) \neq 0$  and  $-s(z)d^{n+1} \neq 0$ . So by Lemma 2.6, we get  $E_2(z) \neq 0$  and

$$T(r, g_1) \leq \overline{N}(r, 1/E_2) + S(r, g_1).$$

By the above results and noting that  $F_n(z) = E_2(z) / \prod_{j=0}^n g(z + jc)$  and  $g_1(z) = g(z) - d$ , we obtain

$$F_n(z) \neq 0$$

and

$$T(r, g) \leq \overline{N}(r, 1/E_2) + S(r, g). \quad (3.11)$$

In order to estimate the zeros of  $F_n(z)$ , we proceed to discuss the common zeros of  $E_2(z)$  and  $\prod_{j=0}^n g(z + jc)$ . Let  $z_0$  be such a common zero. Then  $z_0$  is a zero of  $E_1(z)$  or a pole of  $s(z)$ . Assume that  $z_0$  is a zero of  $E_1(z)$  and that

$$g(z_0 + jc) \neq \infty \quad (3.12)$$

for  $j = 0, \dots, n$ . Since  $\prod_{j=0}^n g(z_0 + jc) = 0$ , there exists an integer  $l \in \{0, \dots, n\}$  such that  $g(z_0 + lc) = 0$ . By (3.7), (3.12) and the fact that  $g(z_0 + lc) = 0$ ,  $E_1(z_0) = 0$ , we get

$$\binom{n}{l} (-1)^{n-l} \prod_{\substack{i=0 \\ i \neq l}}^n g(z_0 + ic) = 0.$$

This equality shows that there exists an integer  $s \in \{0, \dots, n\} \setminus \{l\}$  such that  $g(z_0 + sc) = 0$ . So we have

$$g(z_0 + lc) - g(z_0 + sc) = 0.$$

Since  $g(z) = 1/(f(z) - b)$  and  $c, 2c, \dots, nc$  are not periods of  $f(z)$ , we have

$$g(z + lc) - g(z + sc) \neq 0.$$

Thus, the integrated counting function of the common zeros of  $E_2(z)$  and  $\prod_{j=0}^n g(z + jc)$ , denoted by  $N_1(r)$ , satisfies

$$N_1(r) \leq N(r, s) + \sum_{j=0}^n N(r, g(z + jc)) + \sum_{\substack{l \neq s \\ l, s \in \{0, \dots, n\}}} N\left(r, \frac{1}{g(z + lc) - g(z + sc)}\right).$$

By (3.3), Lemma 2.3 and  $T(r, s) = S(r, g)$ , the above inequality becomes

$$N_1(r) \leq S(r, g) + \sum_{\substack{l \neq s \\ l, s \in \{0, \dots, n\}}} N\left(r, \frac{1}{g(z + lc) - g(z + sc)}\right). \tag{3.13}$$

Since  $p(z)$  in (3.1) is a polynomial of degree  $\deg p = \sigma(g) \geq 1$ , we have

$$p(z) = a_m z^m + p_1(z),$$

where  $a_m (\neq 0)$  is a constant,  $m = \sigma(g) \geq 1$ , and  $p_1(z)$  is a polynomial of degree at most  $m - 1$ . For  $l \neq s$ ,

$$p(z + lc) - p(z + sc) = cm(l - s)a_m z^{m-1} + \dots = p_{l,s}(z), \tag{3.14}$$

where  $p_{l,s}(z)$  are polynomials of degree  $m - 1$ . By (3.1) and (3.14), we have

$$g(z + lc) - g(z + sc) = (h(z + lc)e^{p_{l,s}(z)} - h(z + sc))e^{p(z+sc)}.$$

Since  $\sigma(h) < \sigma(g)$  and  $\sigma(e^{p_{l,s}(z)}) < \sigma(g)$ , it follows by (3.2) and Lemma 2.3 that

$$T(r, h(z + lc)e^{p_{l,s}(z)} - h(z + sc)) = S(r, g).$$

So for  $l, s \in \{0, \dots, n\}$ ,  $l \neq s$ , we have

$$N\left(r, \frac{1}{g(z + lc) - g(z + sc)}\right) = S(r, g). \tag{3.15}$$

By (3.13) and (3.15), we get

$$N_1(r) \leq S(r, g). \tag{3.16}$$

Since  $N_1(r)$  denotes the common zeros of  $E_2(z)$  and  $\prod_{j=0}^n g(z + jc)$ , it follows from (3.6) that

$$N(r, 1/F_n) \geq N(r, 1/E_2) - N_1(r).$$

Combining this inequality with (3.11) and (3.16), we get

$$T(r, g) \leq N(r, 1/F_n) + S(r, g).$$

Moreover,  $g(z) = 1/(f(z) - b)$ . So

$$T(r, f) \leq N(r, 1/F_n) + S(r, f). \tag{3.17}$$

By (3.17),  $F_n(z) \not\equiv 0$  and noting that  $f(z)$  is a transcendental meromorphic function, we see that  $F_n(z)$  is transcendentially meromorphic.

By  $F_n(z) = \Delta^n f(z) - s(z)$  and (3.4), we get

$$T(r, F_n) \leq \sum_{j=0}^n T(r, f(z + jc)) + S(r, f).$$

So by Lemma 2.3,  $T(r, F_n)$  satisfies

$$T(r, F_n) \leq (n + 1)T(r, f) + S(r, f). \tag{3.18}$$

Combining (3.17) and (3.18), we get

$$\delta(0, F_n) = 1 - \limsup_{r \rightarrow \infty} \frac{N(r, 1/F_n)}{T(r, F_n)} \leq n/(n + 1).$$

**Part II** We assume that the condition (ii) in Theorem 1.1 holds. Set  $g_2(z) = f(z) - a$ . Then  $0, \infty$  are Borel exceptional values of  $g_2(z)$ , and

$$N(r, g_2) = S(r, g_2), \quad N(r, 1/g_2) = S(r, g_2).$$

Substituting  $f(z) = g_2(z) + a$  into  $F_n(z)$ , we get

$$F_n(z) = \Delta^n (g_2(z) + a) - s(z) = \Delta^n g_2(z) - s(z).$$

Since  $\Delta^n g_2(z) = \Delta^n f(z) \not\equiv 0$  and  $s(z) \not\equiv 0$ , by Lemma 2.6, it follows that  $F_n(z)$  is transcendentially meromorphic and

$$T(r, g_2) \leq \overline{N}(r, 1/F_n) + S(r, g_2),$$

and so

$$T(r, f) \leq \overline{N}(r, 1/F_n) + S(r, f). \tag{3.19}$$

Moreover, we still have (3.18). By (3.18) and (3.19), we get

$$\delta(0, F_n) \leq n/(n + 1).$$

#### 4. Proof of Theorem 1.2

Since  $\max\{\lambda(f - a), \lambda(1/f)\} < \sigma(f) - 1$  and  $1 < \sigma(f) < \infty$ , we have

$$f(z) = a + h(z)e^{q(z)}, \tag{4.1}$$

where  $q(z)$  is a polynomial of degree  $\deg q = \sigma(f) > 1$ , and  $h(z)$  is a nonzero meromorphic function satisfying  $\sigma(h) < \sigma(f) - 1$ . Let

$$q(z) = d_k z^k + \tilde{q}(z),$$

where  $d_k (\neq 0)$  is a constant,  $k = \sigma(f) > 1$ , and  $\tilde{q}(z)$  is a polynomial of degree at most  $k - 1$ . For  $j = 1, \dots, n$ ,

$$q(z + jc) - q(z) = jkd_k cz^{k-1} + q_j(z), \tag{4.2}$$

where  $q_j(z)$  are polynomials of degree at most  $k - 2$ . Let  $q_0(z) \equiv 0$ . By (3.4), (3.5), (4.1) and (4.2), we have

$$\begin{aligned} \Delta^n f(z) &= \sum_{j=0}^n \binom{n}{j} (-1)^{n-j} f(z + jc) \\ &= \sum_{j=0}^n \binom{n}{j} (-1)^{n-j} \left( a + h(z + jc)e^{q(z+jc)} \right) \\ &= e^{q(z)} \sum_{j=0}^n \binom{n}{j} (-1)^{n-j} h(z + jc)e^{q_j(z)} e^{jkd_k cz^{k-1}}. \end{aligned}$$

Set

$$T(z) = \sum_{j=0}^n \binom{n}{j} (-1)^{n-j} h(z + jc)e^{q_j(z)} e^{jkd_k cz^{k-1}}, \quad t(z) = e^{kd_k cz^{k-1}}.$$

Then we have

$$T(z) = \sum_{j=0}^n \alpha_j(z) t^j(z), \tag{4.3}$$

where for  $j = 0, \dots, n$ ,

$$\alpha_j(z) = \binom{n}{j} (-1)^{n-j} h(z + jc)e^{q_j(z)} \neq 0. \tag{4.4}$$

Since  $t(z)$  is of regular growth  $\sigma(t) = k - 1 > 0$  and noting that  $\sigma(h) < k - 1$  and  $\sigma(e^{q_j(z)}) \leq k - 2$ , we get

$$T(r, \alpha_j) = S(r, t) \tag{4.5}$$

for  $j = 0, \dots, n$ . By Lemma 2.4 and (4.3)–(4.5), we get  $T(z) \neq 0$ . So  $\Delta^n f(z) = e^{q(z)} T(z) \neq 0$  and the condition (ii) in Theorem 1.1 holds. Thus,  $\Delta^n f(z)$  assumes every nonzero finite value infinitely often. Moreover, applying Lemma 2.6 to (4.3), we get

$$T(r, t) \leq \overline{N}(r, 1/T) + S(r, t) = \overline{N}(r, 1/\Delta^n f) + S(r, t).$$

Therefore,  $\Delta^n f(z)$  assumes every finite value infinitely often.

### Acknowledgement

The authors are very grateful to the referee for valuable comments and suggestions.

### References

- [1] Bergweiler, W. and Langley, J.K.: Zeros of differences of meromorphic functions. *Math. Proc. Cambridge Philos. Soc.* 142, 133–147 (2007).

- [2] Chen, Z.X. and Shon, K.H.: On zeros and fixed points of differences of meromorphic functions. *J. Math. Anal. Appl.* 344, 373–383 (2008).
- [3] Chiang, Y.M. and Feng, S.J.: On the Nevanlinna characteristic of  $f(z + \eta)$  and difference equations in the complex plane. *Ramanujan J.* 16, 105–129 (2008).
- [4] Eremenko, A., Langley, J. and Rossi, J.: On the zeros of meromorphic functions of the form  $f(z) = \sum_{k=1}^{\infty} \frac{a_k}{z-z_k}$ . *J. Anal. Math.* 62, 271–286 (1994).
- [5] Goldberg, A.A. and Ostrovskii, I.V.: *Distribution of Values of Meromorphic Functions*. Nauka, Moscow, 1970.
- [6] Halburd, R.G. and Korhonen, R.J.: Difference analogue of the lemma on the logarithmic derivative with applications to difference equations. *J. Math. Anal. Appl.* 314, 477–487 (2006).
- [7] Halburd, R.G. and Korhonen, R.J.: Meromorphic solutions of difference equations, integrability and the discrete Painlevé equations. *J. Phys. A* 40, 1–38 (2007).
- [8] Halburd, R.G. and Korhonen, R.J.: Finite-order meromorphic solutions and the discrete Painlevé equations. *Proc. Lond. Math. Soc.* (3) 94, 443–474 (2007).
- [9] Hayman, W.K.: Picard values of meromorphic functions and their derivatives. *Ann. of Math.* (2) 70, 9–42 (1959).
- [10] Hayman, W.K.: *Meromorphic Functions*. Clarendon Press, Oxford, 1964.
- [11] Hinchliffe, J.D.: The Bergweiler-Eremenko theorem for finite lower order. *Results Math.* 43, 121–128 (2003).
- [12] Weissenborn, G.: On the theorem of Tumura and Clunie. *Bull. Lond. Math. Soc.* 18, 371–373 (1986).
- [13] Yang, C.C.: On deficiencies of differential polynomials, II. *Math. Z.* 125, 107–112 (1972).

Ranran ZHANG, Zongxuan CHEN  
 School of Mathematical Sciences,  
 South China Normal University,  
 Guangzhou 510631, People's Republic of CHINA  
 e-mails: zhrr19820315@163.com, chzx@vip.sina.com

Received: 21.01.2011