

1-1-2016

Removing the specimen with traction during robotic radical prostatectomy does not cause a positive surgical margin

SERKAN ALTINOVA

ABİDİN EGEMEN İŞGÖREN

ZİYA AKBULUT

MUHAMMET FUAT ÖZCAN

ABDULLAH ERDEM CANDA

See next page for additional authors

Follow this and additional works at: <https://journals.tubitak.gov.tr/medical>



Part of the [Medical Sciences Commons](#)

Recommended Citation

ALTINOVA, SERKAN; İŞGÖREN, ABİDİN EGEMEN; AKBULUT, ZİYA; ÖZCAN, MUHAMMET FUAT; CANDA, ABDULLAH ERDEM; ATMACA, ALİ FUAT; and BALBAY, MEVLANA DERYA (2016) "Removing the specimen with traction during robotic radical prostatectomy does not cause a positive surgical margin," *Turkish Journal of Medical Sciences*: Vol. 46: No. 6, Article 7. <https://doi.org/10.3906/sag-1503-21>
Available at: <https://journals.tubitak.gov.tr/medical/vol46/iss6/7>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Medical Sciences by an authorized editor of TÜBİTAK Academic Journals. For more information, please contact academic.publications@tubitak.gov.tr.

Removing the specimen with traction during robotic radical prostatectomy does not cause a positive surgical margin

Authors

SERKAN ALTINOVA, ABİDİN EGEMEN İŞGÖREN, ZİYA AKBULUT, MUHAMMET FUAT ÖZCAN, ABDULLAH ERDEM CANDA, ALİ FUAT ATMACA, and MEVLANA DERYA BALBAY

Removing the specimen with traction during robotic radical prostatectomy does not cause a positive surgical margin

Serkan ALTINOVA^{1*}, Abidin Egemen İŞGÖREN¹, Ziya AKBULUT², Muhammet Fuat ÖZCAN¹,
Abdullah Erdem CANDA³, Ali Fuat ATMACA³, Mevlana Derya BALBAY⁴

¹Department of Urology, Atatürk Training and Research Hospital, Ankara, Turkey

²Department of Urology, Faculty of Medicine, Medipol University, İstanbul, Turkey

³Department of Urology, Faculty of Medicine, Yıldırım Beyazıt University, Ankara, Turkey

⁴Department of Urology, Memorial Şişli Hospital, İstanbul, Turkey

Received: 03.03.2015 • Accepted/Published Online: 14.02.2016 • Final Version: 20.12.2016

Background/aim: The aim of this study was to gauge whether removal of a specimen with traction during robot-assisted laparoscopic radical prostatectomy causes a positive surgical margin or not.

Materials and methods: One hundred and sixty-nine patients with localized prostate cancer who underwent robot-assisted laparoscopic radical prostatectomy from 2009 to 2011 were included in the study. After dividing the patients into two groups, we recorded their characteristics and pre-op/post-op evaluations.

Results: There were 111 and 58 patients in groups 1 (with traction) and 2 (without traction), respectively. We evaluated the patients' ages, follow-up time, body mass index (BMI), prostate-specific antigen (PSA) values, pre-op and post-op Gleason score values, pathological stage, positive surgical margin rates, and biochemical PSA recurrence rates. There was no statistically significant difference between the groups for age, pre-op PSA values, BMI, pre-op and post-op Gleason scores, positive surgical margin rates and biochemical recurrence rates. There was a significant difference between prostate weight, tumor volume, and clinical stage.

Conclusion: Removing the specimen with traction during robot-assisted laparoscopic radical prostatectomy does not cause a positive surgical margin. The incision should be as small as possible for cosmetic appearance.

Key words: Prostate cancer, radical prostatectomy, specimen, traction

1. Introduction

Robot-assisted laparoscopic radical prostatectomy (RALP) has become the preferred surgical technique for localized prostate cancer. One of the most important factors in its oncological success is the surgical margin status (1). Positive surgical margin (PSM) status has a possible relation to the surgeon, surgical technique, and disease burden (1,2). Our aim was to evaluate the effect of traction, likely the cause of the PSM, during specimen removal. There are many studies comparing PSMs according to techniques, pathological findings, and clinical stage, but we found none regarding the technique of specimen removal (3).

2. Materials and methods

A total of 169 patients who were treated by RALP for localized prostate cancer between 2009 and 2011 were included in this study. All the patients were evaluated and ethics committee permission was given for each.

* Correspondence: saltinova@yahoo.com

We planned this study to investigate patients with postoperative PSM but no prostate-specific antigen (PSA) recurrence. Patients were randomized into two groups, A and B, according to whether traction was used or not while removing the specimen. A traction procedure consists of removing the specimen from a small incision. Nontraction procedures remove the specimen from an incision larger than the prostate, which eases the procedure.

Student's t-test was used for follow-up, age, body mass index (BMI), PSA, prostate weight, and tumor volume. The chi-square test was used for Gleason grade, stage, surgical margin invasion (SMI), and biochemical recurrence rate (BCR). All values were calculated as mean and SD. SPSS 16 was used.

3. Results

Group A (traction group) had 111 patients, while group B (nontraction group) had 58. There was a

statistically significant difference between the groups for prostate weight, tumor volume, and clinical stage. Age, BMI, preoperative PSA levels, biopsy Gleason score, prostatectomy Gleason score, pathological stage, SMI status, and BCR were similar for both groups. Patients' preoperative and postoperative characteristics are summarized in Tables 1 and 2. Although there were pT0 patients in both groups, there was no additional therapy, such as androgen deprivation therapy, given preoperatively.

4. Discussion

RALP is currently the main surgical technique for localized prostate cancer. In the United States, 85% of radical prostatectomies are performed robotically (4). Generally, PSM rates after different techniques for radical prostatectomy seem to be equal, but the surgical technique performed may occasionally affect these rates (5,6). Oncologic outcomes of robotic surgery are generally similar with laparoscopic and open surgery (7–10).

Table 1. Preoperative characteristics of patients. *: Statistically significant.

| | Total (n: 169) | Traction group (n: 111, 65.7%) | Nontraction Group (n: 58, 34.3%) | P-value |
|-------------------------------|----------------|--------------------------------|----------------------------------|----------|
| Follow-up, months, mean ± SD | 33.85 ± 8.45 | 38.62 ± 6.30 | 24.72 ± 2.26 | < 0.001* |
| Age, years, mean ± SD | 61.11 ± 6.65 | 61.22 ± 6.81 | 60.91 ± 6.40 | 0.822 |
| BMI, mean ± SD | 26.90 ± 2.97 | 27.07 ± 2.99 | 26.50 ± 2.92 | 0.522 |
| Preoperative PSA, mean ± SD | 8.5 ± 5.73 | 8.88 ± 6.25 | 7.76 ± 4.56 | 0.084 |
| Prostate weight, g, mean ± SD | 53.20 ± 19.13 | 50.22 ± 17.39 | 58.91 ± 21.10 | 0.037* |
| Tumor volume, mL, mean ± SD | 7.85 ± 1.62 | 8.80 ± 1.90 | 6.05 ± 8.54 | 0.029* |
| Biopsy Gleason score, n (%) | | | | 0.336 |
| ≤ 6 | 121 (71.6) | 84 (49.7) (75.7) | 37 (21.9) (63.8) | |
| 3+4 | 26 (15.4) | 15 (8.9) (13.5) | 11 (6.5) (19) | |
| 4+3 | 9 (5.3) | 4 (2.3) (3.6) | 5 (3) (8.6) | |
| >7 | 13 (7.7) | 8 (4.7) (7.2) | 5 (3) (8.6) | |
| Clinical stage, n (%) | | | | < 0.001* |
| cT1 | 78 (46.2) | 78 (46.2) (70.3) | - | |
| cT2 | 91 (53.8) | 33 (19.5) (29.7) | 58 (34.3) (100) | |

Table 2. Postoperative findings of patients.

| | Total (n: 169) | Traction group (n: 111, 65.7%) | Nontraction group (n: 58, 34.3%) | P-value |
|------------------------------------|----------------|--------------------------------|----------------------------------|---------|
| Prostatectomy Gleason score, n (%) | | | | 0.462 |
| pT0 | 9 (5.3) | 6 (3.6) (5.4) | 3 (1.7) (5.2) | |
| ≤ 6 | 92 (54.4) | 66 (39.2) (59.5) | 26 (15.4) (44.8) | |
| 3+4 | 40 (23.7) | 23 (13.6) (20.7) | 17 (10.1) (29.3) | |
| 4+3 | 16 (9.5) | 9 (5.3) (8.1) | 7 (4.2) (12.1) | |
| >7 | 12 (7.1) | 7 (4.1) (6.3) | 5 (3) (8.6) | |
| Pathological stage, n (%) | | | | 0.064 |
| pT0 | 8 (4.7) | 5 (3) (4.5) | 3 (1.7) (5.1) | |
| pT2 | 123 (72.8) | 75 (44.4) (67.6) | 48 (28.4) (82.8) | |
| pT3a | 38 (22.5) | 31 (18.4) (27.9) | 7 (4.1) (12.1) | |
| SMI | | | | 0.746 |
| Negative | 142 (84) | 94 (55.6) (84.7) | 48 (28.4) (82.8) | |
| Positive | 27 (16) | 17 (10.1) (15.3) | 10 (5.9) (17.2) | |
| BCR | 11 (6.5) | 8 (4.7) (7.2) | 3 (1.8) (5.2) | 0.611 |

However, some results suggest that the rates are different for different techniques (11–13). It is well known that the PSM may be related to the disease burden, surgeon, and technique. Robotic surgery has some advantages versus laparoscopic surgery, and these are related to both the patient and the surgeon. To determine if traction may cause a PSM, we randomized the patients into two groups as traction and nontraction. We hypothesized that traction may cause damage to the prostate capsule and show a pseudopositive surgical margin. In our study, PSM rates were similar in both groups. Higher tumor volume and stage can affect PSM rates (2). Although the traction group had higher tumor volume rates and lower clinical stage, the PSM rates were similar. In addition, prostatectomy Gleason scores were similar for both groups. Higher preoperative PSA levels (>10 ng/mL) may have an effect on PSM formation, but our groups had no difference in PSA levels (14). We included operations performed by the same person because PSM rates can differ among

surgeons. Some authors have described a “capsular incision index” to show the damage to the capsule that may cause pseudopositive surgical margins (2). We think that, because the traction made by the fourth arm of the robot may cause a pseudopositive surgical margin, the pathologist must reveal a possible positive margin via colored ink; they must also see the capsule of the prostate. If they do not, this may not be a positive margin. This is very important because it can affect extra therapy options. To avoid unnecessary treatment, both the surgeon and the pathologist must be very careful, as mistakes may not only increase morbidity, but also cost.

In conclusion, surgical margin status after radical prostatectomy is an important topic. Surgical technique is vital for prevention of PSMs, but pathological findings are equally important for determining additional treatment. Removing the specimen with traction during RALP does not cause PSM. The incision should be as small as possible for cosmetic appearance.

References

1. Wiezer AZ, Strope S, Wood DP. Margin control in robotic and laparoscopic prostatectomy: what are the REAL outcomes? *Urol Oncol* 2010; 28: 210-214.
2. Hong H, Mel L, Taylor J, Wu Q, Reeves H. Effects of robotic-assisted laparoscopic prostatectomy on surgical pathology specimens. *Diagn Pathol* 2012; 7: 24-30.
3. Tewari A, Sooriakumaran P, Bloch DA, Seshadri-Kreaden U, Hebert AE, Wiklund P. Positive surgical margin and perioperative complication rates of primary surgical treatments for prostate cancer: a systematic review and meta-analysis comparing retropubic, laparoscopic and robotic prostatectomy. *Eur Urol* 2012; 62: 1-15.
4. Lowrance WT, Parekh DJ. The rapid uptake of robotic prostatectomy and its collateral effects. *Cancer* 2012; 118: 4-7.
5. Philippou P, Waine E, Rowe E. Robot-assisted laparoscopic prostatectomy versus open: comparison of the learning curve of a single surgeon. *J Endourol* 2012; 26: 1002-1008.
6. Coelho RF, Rocco B, Patel MB, Orvieto MA, Chauhan S, Ficarra V, Melegari S, Palmer KJ, Patel VR. Retropubic, laparoscopic and robot-assisted radical prostatectomy: a critical review of outcomes reported by high volume centers. *J Endourol* 2010; 24: 2003-2015.
7. Parsons JK, Bennett JL. Outcomes of retropubic, laparoscopic, and robotic-assisted prostatectomy. *Urology* 2008; 72: 412-416.
8. Ficarra V, Novara G, Fracalanza S, D'Elia C, Secco S, Iafrate M, Cavalleri S, Artibani W. A prospective, non-randomized trial comparing robot-assisted laparoscopic and retropubic radical prostatectomy in one European institution. *BJU Int* 2009; 104: 534-539.
9. Schroeck FR, Sun L, Freedland SJ, Albala DM, Mouraviev V, Polascik TJ, Moul JW. Comparison of prostate-specific antigen recurrence-free survival in a contemporary cohort of patients undergoing either radical retropubic or robot-assisted laparoscopic radical prostatectomy. *BJU Int* 2008; 102: 28-32.
10. Laurila TA, Huang W, Jarrard DF. Robotic-assisted laparoscopic and radical retropubic prostatectomy generate similar positive margin rates in low and intermediate risk patients. *Urol Oncol* 2009; 27: 529-533.
11. Williams SB, Chen MH, D'Amico AV, Weinberg AC, Kacker R, Hirsch MS, Richie JP, Hu JC. Radical retropubic prostatectomy and robotic-assisted laparoscopic prostatectomy: likelihood of positive surgical margin(s). *Urology* 2010; 76: 1097-1101.
12. Cathcart P, Murphy DG, Moon D, Costello AJ, Frydenberg M. Perioperative, functional and oncological outcomes after open and minimally invasive prostate cancer surgery: experience from Australia. *BJU Int* 2011; 107: 11-19.
13. Magheli A, Gonzalgo ML, Su LM, Guzzo TJ, Netto G, Humphreys EB, Han M, Partin AW, Pavlovich CP. Impact of surgical technique (open vs laparoscopic vs robotic-assisted) on pathological and biochemical outcomes following radical prostatectomy: an analysis using propensity score matching. *BJU Int* 2011; 107: 1956-1962.
14. Özkanlı SŞ, Zemheri IE, Yıldırım A, Gür HD, Balbay MD, Şenol S, Özkanlı AO, Alkan E, Zenginkinet T, Aydın A et al. Gleason score at the margin can predict biochemical recurrence after radical prostatectomy in addition to preoperative PSA and surgical margin status. *Turk J Med Sci* 2013; 44: 397-403.