

1-1-2000

## Seasonal Changes in the Phytoplankton of the Northeastern Mediterranean(Bay of Iskenderun)

SEVİM POLAT

ERCAN SARIHAN

TUFAN KORAY

Follow this and additional works at: <https://journals.tubitak.gov.tr/botany>



Part of the [Botany Commons](#)

---

### Recommended Citation

POLAT, SEVİM; SARIHAN, ERCAN; and KORAY, TUFAN (2000) "Seasonal Changes in the Phytoplankton of the Northeastern Mediterranean(Bay of Iskenderun)," *Turkish Journal of Botany*. Vol. 24: No. 1, Article 1. Available at: <https://journals.tubitak.gov.tr/botany/vol24/iss1/1>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Botany by an authorized editor of TÜBİTAK Academic Journals. For more information, please contact [academic.publications@tubitak.gov.tr](mailto:academic.publications@tubitak.gov.tr).

## Seasonal Changes in the Phytoplankton of the Northeastern Mediterranean (Bay of İskenderun)\*

Sevim POLAT, Ercan SARIHAN

Çukurova University, Faculty of Fisheries, 01330, Balcalı, Adana-TURKEY

Tufan KORAY

Ege University, Faculty of Fisheries, 35100, İzmir-TURKEY

Received: 02.10.1998

Accepted: 21.06.1999

**Abstract:** This study was carried out in the northern part of the Bay of İskenderun in the northeastern Mediterranean during the period 1994-1995. The quantitative and qualitative distribution of phytoplankton, the environmental factors that affect its distribution and the changes in succession over time were investigated. At each of the seven stations, monthly sampling was done from the surface and 10.0 m depth. In spring, the sampling was performed once every two weeks. In addition, the measurements of some physical and chemical parameters i.e. temperature, salinity,  $(NO_3^- + NO_2^-)$ -N and  $PO_4^{3-}$ -P were carried out.

At the end of the study, a total of 170 taxa from 5 algae classes, *Cyanophyceae*, *Dinophyceae*, *Prymnesiophyceae*, *Dictyochophyceae* and *Bacillariophyceae* were determined. *Bacillariophyceae* was found to be richer than the other groups in terms of abundance and species number and followed by *Dinophyceae*. *Cyanophyceae*, *Prymnesiophyceae* and *Dictyochophyceae* were represented by only one species each.

**Key Words:** Northeastern Mediterranean, Bay of İskenderun, phytoplankton, diversity, succession, physico-chemical parameters.

### Kuzeydoğu Akdeniz'de (İskenderun Körfezi) Fitoplanktonun Mevsimsel Değişimi

**Özet:** Bu çalışma, İskenderun Körfezi'nin (Doğu Akdeniz) kuzey kesimindeki 1994-1995 yılları arasında sürdürülmüştür. Çalışmada fitoplanktonun nitel ve nicel dağılımı, çevresel faktörlerin bu dağılıma etkileri ve zaman içindeki değişimleri araştırılmıştır. Herbir istasyonda aylık ve bahar aylarında iki haftada bir olmak üzere yüzeyden ve 10.0 m derinlikten örneklemeler yapılmıştır. Ayrıca, sıcaklık, tuzluluk,  $(NO_3^- + NO_2^-)$ -N ve  $PO_4^{3-}$ -P gibi bazı fizikokimyasal parametrelerin de ölçümü yapılmıştır.

Çalışma sonunda *Cyanophyceae*, *Dinophyceae*, *Prymnesiophyceae*, *Dictyochophyceae* ve *Bacillariophyceae* olmak üzere 5 alg sınıfından 170 adet taksa saptanmıştır. *Bacillariophyceae* sınıfı tür sayısı ve yoğunluk olarak diğer sınıflardan daha zengin bulunmuş ve bunu *Dinophyceae* takip etmiştir. *Cyanophyceae*, *Prymnesiophyceae* ve *Dictyochophyceae* birer tür ile temsil edilmiştir.

**Anahtar Sözcükler:** Kuzeydoğu Akdeniz, İskenderun Körfezi, fitoplankton, çeşitlilik, süksesyon, fiziko-kimyasal özellikler.

### Introduction

Many studies have been carried out on phytoplankton ecology in Turkish seas. Some important studies on this subject are those of Gökalp (1) in the Gulf of Edremit, Bodrum and İskenderun and Koray in the Bay of İzmir (2, 3). However, few studies have been carried out, on the phytoplankton composition and seasonal succession at the Northeastern Mediterranean coast of Turkey

especially in the Bay of İskenderun which is a very important area for fisheries.

In this study, the temporal distribution of phytoplankton species were studied in relation to some physico-chemical parameters i.e. temperature, salinity and N/P ratios in the selected area of the Bay of İskenderun.

\* This research was supported by the research fund of Çukurova University (P.no: 94.116) and partly supported by TÜBİTAK, YDABÇAG-296 project.

**Materials and Methods**

This study was carried out at the Bay of Iskenderun, in the northeastern Mediterranean. The coast between Yumurtalik and the Toros Fertilizer Factory in the northern part of the bay was selected and a total of seven sampling stations in the shallow and deeper parts of the area were defined (Fig. 1). For species identification, from each sampling station horizontal samples were taken. For this aim, a standard plankton net with a 55µ mesh size and 40 cm diameter was used. After the haulings, plankton samples were placed in plastic jars, fixed with standard lugol solution and taken to the laboratory for microscopic examination. Samples were preserved in a refrigerator until species identifications. Before microscopic identifications, samples were treated against organic substances. Identifications of the species were done with phase-contrast and binocular research microscopes. The studies used for species identifications were Massuti and Margalef (4), Tregouboff and Rose (5), Brunel (6), Patrick and Reimer (7), Drebes (8), Vinyard (9), Vinyard (10), Cupp (11), Rampi and Bernhard (12), Rampi and Bernhard (13), Koray and Gökpinar (14), Gökpinar and Koray (15), Sournia (16), Ricard (17), Delgado and Fortuna (18).

Quantitative plankton sampling was done vertically with a nansen bottle. The samples were placed in two liter jars and fixed with standard lugol solution. Then formaldehyde was added until a concentration of 4%. Samples were precipitated and the upper layer was removed by siphoning and standard single drop methods were used for counting and also a Sedgwick-Rafter counting chamber for bigger phytoplankters. The results of countings were summarized as cells/l (19). Shannon-Wiener (H') diversity and Pielou evenness (J) indices were used for estimation of diversity and Pielou evenness (20). These formulas are:

$$H' = - \sum_{i=1}^s pi(\log_2 pi)$$

$$J = \frac{H'}{\log_2 S}$$

where pi , cell number of species; i, total cell number; S, species number.

The salinity and temperature measurements of each stations were made with a YSI model salinometer. The




Figure 1. The sampling stations of the Bay of Iskenderun.

analysis of inorganic nitrate and phosphate phosphorus was made according to the standard methods of Parsons et al. (21). N/P ratios were calculated from  $(\text{NO}_3^- + \text{NO}_2^-)$ -N and  $\text{PO}_4^{3-}$ -P values.

## Results

### Physico-chemical parameters

#### Temperature

During the research period, the mean temperature increased to the highest level in August ( $28.9 \pm 0.16^\circ\text{C}$ ) in surface water and decreased to the lowest in January ( $15.8 \pm 0.57^\circ\text{C}$ ) at 10 m depth (Fig. 2). There were no important differences between stations due to temperature. The number of phytoplankton species was poorer in August than in January. The species number of the class *Bacillariophyceae* was more than the class *Dinophyceae* in both August and January.

#### Salinity

Changes in salinity at the surface and 10.0 m depth during the research period showed that salinity reaches the highest level during the summer months. The highest mean salinity was measured in October as  $39.02 \pm 0.04$  ‰ at 10.0 m depth (Fig. 3). The lowest mean salinity was  $37.9 \pm 0.43$  ‰ in September at 10.0 m depth. During winter the salinity was lower and the




Figure 2. The annual cycle of seawater temperature in the Bay of Iskenderun ( $^\circ\text{C}$ ).




Figure 3. The annual cycle of salinity in the Bay of Iskenderun (‰).

number of species was higher in the same period. Total species number was found to be lower during July and August in which the salinity increased to high levels.

#### N/P Ratio

The highest and lowest mean N/P ratios were found to be  $8.09 \pm 2.96$  at the surface in April and  $1.38 \pm 1.05$  at 10 m in July respectively (Table 1). Conspicuous differences were found among stations. Among the stations, the highest N/P ratio was calculated to be 21.2 at the surface at the 1st station in October, 1995. The lowest N/P ratio was calculated to be 0.09 at 10 m depth at the 5th station in October, 1995 and phytoplankton abundance increased at the same station.

#### Phytoplankton composition and its distribution

Members of five algae classes, *Cyanophyceae*, *Dinophyceae*, *Prymnesiophyceae*, *Dictyochophyceae* and *Bacillariophyceae* were determined during the research period (from September 1994 to October, 1995). Classes were represented as: 38 genera, 83 species, three varieties and two formas from *Bacillariophyceae*; 16 genera, 73 species, 25 varieties and two formas from *Dinophyceae*; one species from *Cyanophyceae*; one species from *Prymnesiophyceae*, one species from *Dictyochophyceae* (Table 2). A total of 170 taxa were determined from these algae classes. *Bacillariophyceae* was superior to the other classes in terms of diversity and abundance. The most common genera from *Bacillariophyceae* were *Cerataulina* H. Peragallo, *Chaetoceros* Ehrenberg, *Leptocylindricus* Cleve, *Rhizosolenia* (Ehrenberg) Brightwell, *Thalassiothrix* Cleve & Grunow and *Nitzschia* Hassall. *Chaetoceros* and

Table 1. The monthly distribution of N/P means and  $\pm$  standard deviations (n=7 for 0.0 m, n=6 for 10.0 m.).

	0 m	10 m
N/P ratios		
October 94	$4.87 \pm 2.42$	$4.40 \pm 2.57$
November	$3.07 \pm 2.81$	$1.99 \pm 0.66$
8 January	$3.31 \pm 1.60$	$2.83 \pm 0.93$
31 January	$4.31 \pm 3.23$	$3.72 \pm 2.87$
February	$3.34 \pm 1.32$	$3.83 \pm 0.83$
March	$4.90 \pm 2.00$	$6.46 \pm 5.24$
4 April	$3.75 \pm 0.68$	$3.75 \pm 2.81$
16 April	$8.09 \pm 2.96$	$7.52 \pm 3.22$
30 April	$2.52 \pm 0.29$	$2.52 \pm 0.52$
May	$2.09 \pm 0.66$	$2.25 \pm 0.78$
June	$4.41 \pm 0.82$	$3.05 \pm 2.64$
July	$1.42 \pm 0.78$	$1.38 \pm 1.05$
August	$4.68 \pm 2.17$	$4.60 \pm 2.04$
September	$5.01 \pm 1.35$	$4.90 \pm 1.76$
October 95	$7.09 \pm 7.31$	$2.13 \pm 1.49$

Table 2. The monthly distribution of phytoplankton species.

	S	O	N	J	M	O	N	T	H	S						
	94				J	F	M	A	A	A	M	J	J	A	S	O
																95
<b>CYANOPHYCEAE</b>																
<i>Oscillatoria</i> sp.	+	+	+	+	+	+		+						+	+	+
<b>DINOPHYCEAE</b>																
<i>Amphisolenia bidentata</i> Schröder		+	+	+	+	+	+	+			+					+
<i>Amphisolenia truncata</i> Kofoid end Michener															+	
<i>Ceratium arietinum</i> var. <i>arietinum</i>		+	+	+	+	+	+	+		+	+		+	+		+
<i>Ceratium biceps</i> Claparede et Lachm.			+	+	+	+	+	+	+	+	+	+	+			+
<i>Ceratium breve</i> var. <i>schmidtii</i>															+	
<i>Ceratium breve</i> var. <i>parallellum</i> (Schmidt) Jörgensen		+	+				+									
<i>Ceratium candelabrum</i> var. <i>candlabrum</i>	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
<i>Ceratium candelabrum</i> var. <i>depressum</i> (Pouc) Jörgensen								+	+	+	+	+	+			
<i>Ceratium carriense</i> var. <i>carriense</i>				+	+			+								
<i>Ceratium carriense</i> var. <i>volans</i>		+	+	+	+	+	+		+		+		+			+
<i>Ceratium concilians</i> Jörgensen			+	+	+	+		+		+	+					
<i>Ceratium contortum</i> var. <i>contortum</i> (Gourret) Cleve			+	+	+	+	+									
<i>Ceratium contortum</i> var. <i>karstenii</i> (Pavillard) Sournia			+	+	+	+	+	+		+						+
<i>Ceratium contrarium</i> (Gourret) Pavillard					+						+	+	+			
<i>Ceratium declinatum</i> var. <i>majus</i> Jörgensen						+	+	+	+	+	+					+
<i>Ceratium eucarvatum</i> Jörgensen		+	+	+	+	+	+	+	+	+	+		+			+
<i>Ceratium furca</i> var. <i>furca</i>			+	+	+	+	+	+	+	+	+	+				+
<i>Ceratium fusus</i> var. <i>fuscus</i>		+	+	+	+	+	+	+		+	+	+	+			+
<i>Ceratium gibberum</i> var. <i>dispar</i> (Pouch.) Sournia						+										
<i>Ceratium hexacanthum</i> var. <i>contortum</i> Lemmermann						+	+	+			+	+				
<i>Ceratium hexacanthum</i> var. <i>hexacanthum</i>					+									+		
<i>Ceratium horridum</i> var. <i>denticulatum</i> Jörgensen			+				+	+	+	+						+
<i>Ceratium horridum</i> var. <i>horridum</i>				+		+					+	+				
<i>Ceratium inflatum</i> (Kofoid) Jörgensen	+	+	+	+	+	+	+	+	+	+	+		+		+	+
<i>Ceratium kofoidi</i> Jörgensen	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
<i>Ceratium limulus</i> Gourret				+	+											
<i>Ceratium longirostrum</i> Gourret		+		+	+	+	+				+					
<i>Ceratium macroceros</i> var. <i>gallicum</i> Kofoid	+	+	+	+	+	+	+	+	+	+	+	+	+		+	+
<i>Ceratium macroceros</i> var. <i>macroceros</i>					+		+									
<i>Ceratium massiliense</i> var. <i>massiliense</i>			+			+	+	+	+	+	+	+	+			+
<i>Ceratium pentagonum</i> Gourret			+	+	+	+	+									+
<i>Ceratium ranipes</i> var. <i>palmatum</i> (Schröder) Cleve			+	+	+	+										
<i>Ceratium symmetricum</i> var. <i>orthoceros</i> Jörg. Grah. et Bronikovsky			+	+		+		+			+					
<i>Ceratium symmetricum</i> var. <i>symmetricum</i>		+	+	+	+	+	+		+	+						+
<i>Ceratium teres</i> Kofoid					+	+										+
<i>Ceratium trichoceros</i> (Ehrenberg) Kofoid		+	+	+	+	+	+	+	+	+	+	+	+	+		+
<i>Ceratium tripos</i> var. <i>atlanticum</i> (Ostenfeld) Paulsen	+	+		+		+	+	+	+	+	+	+	+	+	+	+
<i>Ceratium tripos</i> var. <i>pulchellum</i> (B. Schröder) Lopez				+			+	+	+	+	+	+				+
f. <i>semipulchellum</i> Jörgensen							+	+	+	+	+	+				+
f. <i>pulchellum</i>		+	+	+	+	+	+		+	+	+	+	+	+	+	+
<i>Ceratocorys armata</i> (Schütt) Kofoid															+	
<i>Ceratocorys gourreti</i> Paulsen						+										+
<i>Ceratocorys horrida</i> Stein	+	+	+	+	+	+	+	+		+	+	+	+	+	+	+
<i>Dinophysis caudata</i> Saville-Kent	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
<i>Dinophysis doryphorum</i> (Stein) Abé											+		+			

Table 2. Continued.

	S	O	N	J	J	F	M	A	A	A	M	J	J	A	S	O
	94															95
<i>Dinophysis hastata</i> Stein								+			+			+		
<i>Dinophysis mitra</i> (Schütt) Abé				+												
<i>Diplopsalis lenticula</i> Bergh			+				+						+			
<i>Gonyaulax birostris</i> Stein					+											
<i>Gonyaulax diegensis</i> Kofoid													+	+		
<i>Gonyaulax monocantha</i> Pavillard											+					
<i>Gonyaulax polyedra</i> Stein				+												
<i>Gonyaulax spinifera</i> (Claparede et Lachmann)Diesing	+							+	+	+			+	+	+	
<i>Gonyaulax polygramma</i> Stein	+	+		+	+	+					+			+	+	+
<i>Gonyaulax turbynei</i> Murray et Whitting																+
<i>Heteraulacus polyedricus</i> (Pouchet)Drugg etLoeblich				+	+	+	+	+			+	+	+	+		
<i>Heteraulacus sphaericus</i> (Murrat et Whitting)LoeblichIII				+												
<i>Kofoidinium velelloides</i> Pavillard				+							+			+		
<i>Ornithocercus quadratus</i> Schütt		+	+		+	+					+	+		+		+
<i>Oxytoxum scolopax</i> Stein	+															+
<i>Oxytoxum</i> sp.				+				+			+			+		+
<i>Podolampas bipes</i> Stein	+			+		+					+	+	+	+	+	+
<i>Podolampas spinifera</i> Okamura											+		+			+
<i>Prorocentrum compressum</i> (Bailey) AbÉ ex Dodge				+												
<i>Prorocentrum micans</i> Ehrenberg		+	+		+	+		+	+	+	+	+		+	+	+
<i>Protoperidinium brochi</i> (Kofoid et Swezy) Balech				+	+	+		+					+	+		+
<i>Protoperidinium claudicans</i> (Paulsen) Balech						+	+	+					+			
<i>Protoperidinium conicum</i> (Gran)Balech	+						+	+	+	+	+	+	+	+	+	+
<i>Protoperidinium depressum</i> (Bailey)Balech							+			+	+	+	+	+		
<i>Protoperidinium divergens</i> (Ehrenberg)Balech	+		+	+		+	+	+	+	+	+	+	+	+	+	
<i>Protoperidinium grande</i> (Kofoid)Balech	+										+	+	+			+
<i>Protoperidinium globolus</i> (Stein)Balech	+	+				+										
<i>Protoperidinium leonis</i> (Pavillard)Balech					+											
<i>Protoperidinium oceanicum</i> (Vanhöffen) Balech	+			+	+			+	+	+	+				+	+
<i>Protoperidinium pallidum</i> (Ostenfeld)Balech						+										+
<i>Protoperidinium mediterraneum</i> (Kofoid)Balech	+			+	+	+	+	+	+	+	+	+	+	+		+
<i>Protoperidinium pedunculatum</i> (Schütt)Balech					+											
<i>Protoperidinium pyriforme</i> (Paulsen) Balech					+						+					
<i>Protoperidinium quarnerense</i> (Schröder)Balech												+	+			
<i>Protoperidinium steinii</i> (Jörgensen)Balech																+
<i>Pyrocystis fusiformis</i> W.Thomson	+															
<i>Pyrophacus horologium</i> Stein	+										+	+	+	+	+	
<i>Pyrophacus steinii</i> (J.Schiller) Wall et Dale	+			+						+			+	+	+	
<i>Spiraulax jollifei</i> (Murray et Whitting) Kofoid				+								+	+			
<b>PRYMNESIOPHYCEAE</b>																
<i>Halosphaeria viridis</i> Schmitz				+	+											
<b>DICTYOCOPHYCEAE</b>																
<i>Dictyocha fibula</i> Ehrenberg	+	+	+	+	+	+		+						+	+	+
<b>BACILLARIOPHYCEAE</b>				+												
<i>Amphiprora gigantea</i> Grunow			+	+	+	+	+								+	+
<i>Asterionella japonica</i> Cleve et Möller				+	+	+	+	+	+	+	+	+	+			
<i>Asterolampra grevillei</i> (Wallich) Greville		+	+	+		+	+	+								
<i>Asterolampra marylandica</i> Ehrenberg	+	+	+	+	+	+									+	+
<i>Asteromphalus flabellatus</i> (Brebisson)Greville															+	

Seasonal Changes in the Phytoplankton of the Northeastern Mediterranean (Bay of Iskenderun)

Table 2. Continued.

	S	O	N	J	J	F	M	A	A	A	M	J	J	A	S	O
	94															95
<i>Bacteriastrium delicatulum</i> Cleve	+					+	+	+	+	+	+		+		+	+
<i>Bacteriastrium hyalinum</i> Lauder		+	+	+	+		+					+	+	+		+
<i>Campylodiscus</i> sp.		+	+										+			
<i>Cerataulina pelagica</i> (Cleve) Hendey	+	+		+	+	+	+	+	+	+		+	+	+	+	+
<i>Chaetoceros affine</i> var. <i>willei</i> (Gran)Hustedt.	+			+	+	+	+	+	+	+		+	+	+	+	+
<i>Chaetoceros anastomosans</i> Grunow in VanHeurck											+					
<i>Chaetoceros atlanticum</i> Cleve				+	+	+	+	+	+		+			+		+
<i>Chaetoceros coarctatum</i> Lauder						+										
<i>Chaetoceros compressum</i> Lauder							+					+				
<i>Chaetoceros constrictum</i> Gran						+	+	+	+							
<i>Chaetoceros costatum</i> Pavillard								+	+	+		+				
<i>Chaetoceros curvisetum</i> Cleve			+		+	+	+					+		+	+	
<i>Chaetoceros danicum</i> Cleve								+	+	+						
<i>Chaetoceros debile</i> Cleve							+									
<i>Chaetoceros decipiens</i> Cleve	+	+	+	+	+	+	+	+	+	+	+	+	+	+		+
<i>Chaetoceros didymum</i> Ehrenberg																
var. <i>protuberans</i> (Lauder)Gran.et Yendo.								+	+	+			+	+		+
<i>Chaetoceros diversum</i> Cleve	+					+		+	+		+					
<i>Chaetoceros gracile</i> Schütt							+	+	+	+						
<i>Chaetoceros lacinosum</i> Schütt												+	+			+
<i>Chaetoceros lorenzianum</i> Grunow							+	+	+			+	+			+
<i>Chaetoceros messanense</i> Castracane					+											
<i>Chaetoceros peruvianum</i> Brightwell												+				+
<i>Chaetoceros pseudocurvisetum</i> Mangin				+	+	+	+	+	+	+	+	+	+	+	+	+
<i>Chaetoceros rostratum</i> Lauder							+	+	+	+	+					
<i>Chaetoceros tetrastichon</i> Cleve							+	+	+	+	+	+	+		+	+
<i>Chaetoceros tortissimum</i> Gran							+									
<i>Climacosphaenia moniligera</i> Ehrenberg							+	+		+	+		+	+		+
<i>Coscinodiscus perforatus</i> Ehrenberg		+	+													
<i>Coscinodiscus radiatus</i> Ehrenberg				+	+											+
<i>Coscinodiscus</i> sp.	+	+	+	+	+	+		+							+	+
<i>Cylindrotheca closterium</i> (Ehren.)Lewin&Reimann			+	+	+		+				+		+	+	+	+
<i>Dactyliosolen mediterraneus</i> H. Peragallo							+	+								+
<i>Ditylum brightwellii</i> (T.West)Grunow in Van Heurck				+												
<i>Eucampia zoodiacus</i> Ehrenberg				+	+	+		+	+				+			
<i>Grammatophora</i> sp.				+												
<i>Guinardia flaccida</i> (Castracane) H.Peragallo	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
<i>Gyrosigma balticum</i> (Ehrenberg) Rabenhorst													+			
<i>Gyrosigma spenceri</i> (Quekett.) Griffith et Henfery			+	+	+											
<i>Gyrosigma</i> sp.		+	+	+	+					+					+	
<i>Gyrosigma tenuissimum</i> (W.Smith)Griff. et Henfrey	+											+	+	+	+	+
<i>Helicotheca thamesis</i> (Shrubsole)Ricard	+		+	+	+									+	+	
<i>Hemiaulus membranaceus</i> Cleve			+			+								+	+	
<i>Hemiaulus hauckii</i> Grunow in Van Heurck	+	+		+	+	+	+	+	+	+	+	+	+	+	+	+
<i>Lauderia borealis</i> Gran		+														
<i>Leptocylindricus danicus</i> Cleve	+	+	+	+	+	+	+	+	+	+	+		+	+	+	+
<i>Leptocylindricus minimus</i> Gran									+	+	+	+	+	+	+	+
<i>Licmophora abbreviata</i> Agardh				+	+	+	+	+					+	+	+	+
<i>Lithodesmium undulatum</i> Ehrenberg				+	+	+	+			+			+			
<i>Melosira sulcata</i> (Ehrenberg)Kützing			+													

Table 2. Continued.

	S	O	N	J	J	F	M	A	A	A	M	J	J	A	S	O
	94															95
<i>Navicula</i> sp.	+	+	+		+	+										
<i>Nitzschia longissima</i> (Brebis.in Kützing.) Ralf in Prit.			+	+	+	+	+	+	+		+	+	+		+	
<i>Nitzschia paradoxa</i> Grunow in Cleve et Grunow	+	+	+	+	+	+		+		+			+		+	
<i>Nitzschia sigma</i> (Kützing)W. Smith		+	+	+			+	+	+				+		+	+
<i>Nitzschia</i> sp.	+		+							+						
<i>Odontella mobiliensis</i> (Bailey) Grunow	+	+	+	+	+						+		+	+	+	+
<i>Pleurosigma elongatum</i> W. Smith																
<i>Pleurosigma normanii</i> Ralfs in Pritchard.	+	+	+	+	+	+	+						+	+	+	+
<i>Pleurosigma</i> sp1.						+										
<i>Pleurosigma</i> sp 2.															+	
<i>Podocystis</i> sp.					+				+				+			+
<i>Pseudonitzschia pungens</i> (Grunow ex Cleve) Hasle	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
<i>Rhabdonema adriaticum</i> Kützing				+	+	+	+	+	+	+		+				+
<i>Rhizosolenia alata</i> . f. <i>gracillima</i> (Cleve)Gran	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
<i>Rhizosolenia alata</i> .f. <i>indica</i> (H.Peragallo.)Gran	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
<i>Rhizosolenia calcar-avis</i> Schultze	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
<i>Rhizosolenia castracanei</i> H.Peragallo				+												
<i>Rhizosolenia fragilissima</i> Bergon														+	+	
<i>Rhizosolenia imbricata</i> var. <i>shrubsolaei</i> (Cleve)Schütt.	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
<i>Rhizosolenia robusta</i> Norman in Pritchard	+	+		+	+	+	+	+	+	+	+	+	+	+	+	
<i>Rhizosolenia setigera</i> Brightwell	+				+		+	+	+	+	+		+			
<i>Rhizosolenia stolterfothii</i> H. Peragallo	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
<i>Rhizosolenia sytliiformis</i> Brightwell	+	+	+												+	+
<i>Skeletonema costatum</i> (Greville) Cleve			+	+	+		+	+	+	+	+			+		
<i>Striatella unipunctata</i> (Lyngbye)Agardh					+											+
<i>Surirella fastuosa</i> Shrubsole	+		+		+											
<i>Synedra ulna</i> (Nitzsch.) Ehrenberg	+	+														
<i>Thalassionema nitzschioides</i> Hustedt	+		+	+	+		+	+			+					+
<i>Thalassiothrix fraunfeldii</i> Grunow	+	+	+	+	+	+	+	+		+	+	+	+	+	+	+
<i>Thalassiothrix mediterranea</i> Pavillard		+	+									+	+	+	+	+

*Rhizosolenia* were represented as 22 and nine species respectively.

*Cerataulina pelagica* (Cleve) Hendey, *Guinardia flaccida* (Castracane) H .Peragallo and *Pseudonitzschia pungens* (Grunow ex Cleve) Hasle from *Bacillariophyceae* were very important in terms of abundance and they were responsible for phytoplankton blooms during the research period. *Thalassiothrix fraunfeldii* Grunow, *Hemiaulus hauckii* Grunow, *Rhizosolenia calcar-avis* Schultze and *R. stolterfothii* H. Peragallo were not very abundant but existed in all sampling periods.

Dinophyceae were the second important class after *Bacillariophyceae* in terms of species distribution and abundance. *Ceratium* Schrank, *Protoperdinium* Bergh, *Prorocentrum* Ehrenberg and *Dinophysis* Ehrenberg were the most abundant genera from *Dinophyceae*. A total of 27 *Ceratium* species were determined and the

genus *Ceratium* was represented with the highest number of species. *Cyanophyceae*, *Dictyochophyceae* and *Prymnesiophyceae* were represented with only one genus and species.

In all stations, an increase in phytoplankton abundance was determined at the end of January and April in general. But local increases in cell numbers were determined in August and October (Figs. 4, 5). An increase in phytoplankton abundance in January was because of *Cerataulina pelagica* bloomings from the class *Bacillariophyceae*. The increase in phytoplankton abundance in April was mostly because of *Cerataulina pelagica* and *Chaetoceros atlanticum* that reached 66.300 cells/ l and 6040 cells/l in abundance respectively. Apart from these increases, local increases were also determined especially at stations near the coast line. Especially at the 5th station, *Pseudonitzschia*



*pungens* exhibited frequent bloomings in August and October. The abundance of this species reached 320.480 cells/l in August, 380.000 cells/l in October. The taxa from *Dinophyceae* were determined at all stations while the abundance of them was low compared with *Bacillariophyceae*.

The increase in abundance of *Dinophyceae* was determined only in May due to *Prorocentrum micans* Ehrenberg (Fig. 4). In this period, the abundance of *P. micans* reached 6460 cells/l at the 5th station.

The monthly changes in diversity indices are shown in Figs. 6 and 7. Shannon-Wiener diversity (H') in surface waters were found to be between 0.10 and 2.96. Evenness indices (J) were evaluated as 0.04 and 0.84 in

surface waters. The highest values were observed in February because the species number did not show important changes but cell numbers were decreased.

During the research period, the lowest H' and J values in surface waters were found in August. The abundance of *Pseudonitzschia pungens* increased especially at the 5th station in August. In October, cell numbers of *Pseudonitzschia pungens* increased and thus diversity indices decreased again (Fig. 6). Diversity indices calculated for 10 m depth were found lower than for surface waters. The highest values at 10 m depth were found in February. The lowest values were found in August and October for Shannon-Wiener diversity and Pielou evenness indices (Fig. 7).




Figure 4. The monthly changes in the mean cell numbers of diatoms and dinoflagellates at surface water (mean cell concentrations and standard deviations).




Figure 5. The monthly changes in the mean cell number of diatoms and dinoflagellates at 10 m depth (mean cell concentrations and standard deviations)




Figure 6. The monthly changes in Shannon-Wiener diversity ( $H'$ ) and Pielou evenness ( $J$ ) indices of phytoplankton species at surface.




Figure 7. The monthly changes in Shannon-Wiener diversity ( $H'$ ) and Pielou evenness ( $J$ ) indices of phytoplankton species at 10 m.

## Discussion

At the end of the study, the phytoplanktonic members of algae from five classes, *Cyanophyceae*, *Dinophyceae*, *Prymnesiophyceae*, *Dictyochophyceae* and *Bacillariophyceae* were determined. Among them the members of *Bacillariophyceae* were generally dominant during the research period. When species number and abundance are taken into consideration, the members of this class were abundant especially in nutrient-rich shallow stations. It is well known that because of coastal inputs, Iskenderun Bay is more productive than the other regions of the northeastern Mediterranean which has oligotrophic waters in terms of nutrients and primary production. Yilmaz et al. (22) reported that, primary productivity in Iskenderun Bay is two to four times higher than that of the open sea. However, the mean N : P ratios observed were below the theoretical assimilation ratio (15:1) during the whole sampling period.

Phytoplankton composition and abundance were found to be different due to the location of stations. At the coastal stations, diversity and abundance of diatoms were higher than those of dinoflagellates. It was stated by Azov (23) that diatoms were found to be higher, when compared with the dinoflagellates at neritic stations on the Israeli coast.

In terms of qualitative distribution of phytoplankton, it can be said that most of the species identified in the present study consist of phytoplanktonic algae reported in different regions of the Mediterranean before. Most of the identified taxa were neritic, temperate and subtropical species. Also oceanic species were determined. Lakkis and Lakkis (24) studied the coast of Lebanon and stated that the phytoplankton of this region is cosmopolitan, temperate and tropical, distributed in all temperate and subtropical seas in general.

In the present study, an increase of phytoplanktonic organisms was detected at the end of January and in the middle of April. It is considered that, especially *Cerataulina pelagica* was responsible for the phytoplankton increase in January. Koray (3) reported that this species was one of the widespread species in Izmir Bay, Aegean Sea. The phytoplankton increase in April was caused by *C. pelagica* and *Chaetoceros atlanticum*. In addition some stations near the coast have nutrient-rich waters and some local increases were determined at these stations (especially at the 5th stations) in August and October as a result of bloomings of *Pseudonitzschia pungens*. The abundance of this species in August and October reached 320.480 cells/l and 380.000 cells/l respectively. A low diversity index value was found during the same period. On the other hand, the lowest phytoplankton abundance was determined in February and March.

Species of the genera *Chaetoceros*, *Rhizosolenia*, *Brightwell* and *Hemiaulus hauckii*, *Thalassiothrix fraunfeldii*, *Ceratulina pelagica*, *Leptocylindricus danicus* Cleve were common during the whole research period. Species from the genus *Chaetoceros* were observed almost in all stations and their abundance was high in coastal stations. Therefore, it was considered that the members of the genus *Chaetoceros* were important in terms of diversity and abundance. *C. decipiens* Cleve, *C. atlanticum*, *C. affine* were widespread species. Among them, *C. atlanticum* was the most abundant species. *C. decipiens* and *C. atlanticum* are known as oceanic species (2). It was thought that these species were transferred with water circulation from the open sea and they can adapt to coastal waters. Indeed, the species belonging to the genus *Chaetoceros* were the most abundant diatom species in Gökova Bay (Koray et al., 25), at Levantine Basin and the coast of Israeli (23).

Species from the genus *Rhizosolenia* were the second most important species after the genus *Chaetoceros* in terms of the species richness. In present study, a total of nine species from the genus *Rhizosolenia* were identified. This number is lower than that reported by Kideyş et al. (28) and Koray (3). These studies were carried out at the Erdemli coast, the Northeastern Mediterranean and Izmir Bay in which 15 and 14 species and varieties belonging to the genus *Rhizosolenia* were reported respectively. *R. robusta* Norman, *R. alata*, *R. calcar-avis*, *R. stolterfothii* were the most common species. Among them, *R. alata* is known as a cosmopolitan, oceanic and neritic species (26). *R. calcar-avis* is considered a characteristic species of nutrient poor oligotrophic waters (27). In the present study, species from *Rhizosolenia*

were represented with at least a few species in each sampling period, but they never increased very much in any period.

*Hemiaulus hauckii* was considered one of the common species in this study. This species was identified in almost all stations during the research period but it never reached high abundance. This species is considered as characteristic species of nutrient-poor oligotrophic waters. In addition, it was stated that *H. hauckii* continued to exist in waters after spring blooms (27). Therefore, it is clearly observed that there is a similarity between these opinions and the results obtained in the present research.

*Ceratium* was the most common and abundant genus in *Dinophyceae*. In this study, a total of 27 species were identified from the genus *Ceratium*, which is smaller than that given by Koray and Gökpinar (14). He studied Izmir Bay and a total of 28 species were identified from the same genus. In the present study it is observed that, *Ceratium kofoidi* Jörgensen, *C. trichoceros* (Ehrenberg) Kofoid, *C. candelabrum*, *C. furca* var. *furca* from the genus *Ceratium* were the most common species. The members from the genus *Proto-peridinium* were the second after the genus *Ceratium* in terms of species number. A total of 15 species were identified belonging to the genus *Proto-peridinium*. Five species were recorded by Koray et al. (25) from this genus in Gökova Bay, a number smaller than that obtained from the present study. In this study, *Proto-peridinium divergens* (Ehrenberg) Balech was the most abundant species of the genus. *Prorocentrum micans* from *Dinophyceae* is usually described as cosmopolitan, eurythermic and euryhaline because of its very wide distribution (27). In the present study, this species was also common. In general, the numbers of the dinoflagellat species were found to be low at coastal stations, but high in deep stations. Azov (23) stated that dinoflagellat abundance was higher in pelagic stations. On the other hand, the increase in dinoflagellat abundance in phytoplankton blooms did not reach the level of the increase in diatome abundance during the same period. In phytoplankton blooms after nutrient rich periods in coastal waters, diatoms were found to be dominant.

The *Cyanophyceae* class was represented by only one species. *Oscillatoria* sp. from *Cyanophyceae* was identified rarely and its abundance were always at a quite low level during the research period. It was observed during all the periods except the end of spring and the beginning of summer.

*Dictyocha fibula* Ehrenberg was the only species from *Dictyochophyceae*. Two species belonging to *Dictyocha* Ehrenberg were also reported by Kideyş et al. (28) at the Erdemli coast, Northeastern Mediterranean. However, the abundance of this species was generally low during the research period in present study.

If the physical and chemical data are taken into consideration, the lowest temperature was recorded in January, which is the time when the phytoplanktonic organisms increased. Cell numbers of phytoplankton were found to be low except at the 5<sup>th</sup> station in August, which is the time that highest temperature was

measured and the high salinity levels was recorded in summer conditions. According to Smayda (29), Braarud and McLachlan stated that wide salinity ranges are good conditions for growth of phytoplankton species. However, because of the euryhaline nature of the phytoplankton and the modest seasonal fluctuations in the salinity, salinity does not appear to be a significant regulatory factor in phytoplankton succession (29). In the present study N/P ratios were also found to be low. The highest and lowest mean N/P ratios were estimated in April and July respectively. April is the time of the phytoplankton abundance increased. In general, N/P ratios were relatively high during the period when phytoplankton abundance increased.

## References

- Gökalp, N., Edremit, Bodrum ve Iskenderun Körfezlerinin Plankton Durumunun Karşılaştırmalı İncelenmesi, İst. Üniv. Fen Fak. Hid. Ar. Enst. Yay., sayı 3, 71, (1972).
- Koray, T., One Celled Microplankton Species of Izmir Bay (Aegean Sea): A Species List. A Comparison With Records of Adjacent Regions, Doğa, TÜJ Biology, 129-146, (1987).
- Koray, T., Phytoplankton Species Succession, Diversity and Nutrients in Neritic Waters of The Aegean Sea (Bay of Izmir). TÜBITAK, Journal of Botany, 531-544, (1995).
- Massuti, M., Margalef, R., Introducción Al Estudio Del Plancton Marino, Patronato Juan De La Cierva De Inves Tigation Tecnica, Seccion De Biologica Marina, Barcelona, (1930).
- Tregouboff, G., Rose, M., Manuel de Planctonologie Mediterranee, Centre National De la Resherche Scientifique, I, II., 587 p., 207 pl. Paris, (1957).
- Brunel, J., Le Phytoplankton de la Baie des Chaleurs, Contributions Du Ministere De la Chasse Et Des Pecheries, Province De Quebec, No:91 ,(1962).
- Patrick, R., Reimer, C.W., The Diatoms of the United States, vol 1, The Academy of Natural Sciences of Philadelphia, No 13, (1966).
- Drebes G., Marines Phytoplankton, Eine auswahl der Helgolander Planktonalgen (Diatomeen, Peridineen), George Thieme Verlag, Stuttgart, (1974).
- Vinyard, W.C., Key to the Genera of Diatoms of the Inland Waters of Temperate North America, Mad River Press, California, (1974).
- Vinyard, W. C., A Key to The Genera of Marine Planktonic Diatoms of The Pacific Coast of North America, Mad River Press, California, (1975).
- Cupp, E. E., Marine Plankton Diatoms of the West Coast of North America, Otto Koeltz Science Publishers, Koenigstein, (1977).
- Rampi, L., Bernhard, M., Chrave per la Determinazione Delle Peridinee Pelagiche Mediterranee, C.N.E.N., RT/BIO, (1980).
- Rampi, L., Bernhard, M., Key For The Determination of Mediterranean Pelagic Diatoms, C.N.E.N., RT/BIO, (1978).
- Koray, T., Gökpinar, Ş., Ceratium Schrank Genusu Türleri'nin Izmir Körfezi'ndeki Kalitatif ve Kantitatif Özellikleri, E. Ü. Fac. of Sci. Jour. Ser. B, Suppl. Izmir, 178-200, (1983).
- Gökpinar, Ş., Koray, T., Izmir Körfezi Planktonunda Rastlanan Rhizosolenia (Ehrenberg) Brightwell Genusu Üzerine Gözlemler, E. Ü. Fac. of Sci. Jour. Ser. B, Suppl., Izmir, 201-219, (1983).
- Sournia, A., Atlas Du Phytoplankton Marin, Vol., Cyanophycees, Dictyochophycees, Dinophycees, Raphidophycees, Editions Du National De la Resherche Scientifique, Paris, (1986).
- Ricard, M., Atlas Du Phytoplankton Marin, Vol. II, Diatomophycees Editions Du Centre National De la Resherche Scientifique, Paris, (1987).
- Delgado, M., Fortuna, J. M., Atlas De Fitoplancton Del Mar Mediterraneo, Scientia Marina, Barcelona, Vol. 55, 133, (1991).
- Semina, H. J., Treatment of an Aliquot Sample, Phytoplankton Manual, Editor; A. Sournia, UNESCO, (1978).
- Omori, M., Ikeda, T., Methods in Marine Zooplankton Ecology, A Wiley Interscience Publication 269-275, (1984).
- Parsons, T.R., Maita, Y., Lalli, C.M., A Manual of Chemical and Biological Methods for Seawater Analysis, Pergamon Press, Oxford, (1984).
- Yılmaz, A., Baştürk, Ö., Saydam, C., Ediger, D., Yılmaz, K., Hatipoglu, E., Eutrophication in Iskenderun Bay, Northeastern Mediterranean. Science of the Total Environment, Supplement, Elsevier Science Publishers B.V., Amsterdam, 705-717, (1992).

23. Azov, Y., Seasonal Patterns of Phytoplankton Productivity and Abundance in Nearshore Oligothropic Waters of The Levantin Basin (Mediterranean), *Journal of Plankton Research*, 8:41-53, (1986).
24. Lakkis, S., Novel, Lakkis, V., Composition, Annual Cycle and Species Diversity of Phytoplankton in Lebanese Coastal Water, *Jour. of Plank. Res.* Vol 3, No 1, 123-135, (1980).
25. Koray, T., Benli, H.A., Sever, T.M., Gökova Körfezi'nin (Ege Denizi) Fitoplankton ve Protozooplankton Türleri Üzerine Bir Araştırma, *E. Ü. Su Ürünleri Dergisi*, Sayı: 42-43, 11:9-15, (1994).
26. Marshall, G. H., Distribution and Concentration Patterns of Ubiquitous Diatoms for the Northeastern Continental Shelf of the United States, 9th DIATOM-SYMPOSIUM, 75-85, (1986).
27. Kırmor, B., Round Table on Indicator Species in Marine Phytoplankton, II. Background Presentation, *Rapp. Comm. int. Mer. Medit.*, 29, 9, 65-79, (1985).
28. Kıdeyş, A. E., Ünsal, M., Bingel, F., Seasonal Changes Net Phytoplankton of Erdemli Northeastern Mediterranean, *TÜBITAK Journal of Botany*, vol. 45-54, (1989).
29. Smayda, T. J., Phytoplankton Species Succession, *Physiological Ecology of Phytoplankton*, Morris, I., Blackwell Scientific Publications, Oxford, p.493-569, (1980).