

1-1-2000

## A New Liverwort for the Flora of Turkey

İSA GÖKLER

HATİCE ÖZENOĞLU KİREMİT

FERHAT KİREMİT

Follow this and additional works at: <https://journals.tubitak.gov.tr/botany>



Part of the [Botany Commons](#)

---

### Recommended Citation

GÖKLER, İSA; KİREMİT, HATİCE ÖZENOĞLU; and KİREMİT, FERHAT (2000) "A New Liverwort for the Flora of Turkey," *Turkish Journal of Botany*: Vol. 24: No. 1, Article 9. Available at: <https://journals.tubitak.gov.tr/botany/vol24/iss1/9>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Botany by an authorized editor of TÜBİTAK Academic Journals. For more information, please contact [academic.publications@tubitak.gov.tr](mailto:academic.publications@tubitak.gov.tr).

## A New Liverwort for the Flora of Turkey

İsa GÖKLER, Hatice ÖZENOĞLU, Ferhat KİREMIT

Dokuz Eylül Univ., Buca Education Fac., Biology Department, 35150 Buca, İzmir-TURKEY

Received: 10.03.1999

Accepted: 20.08.1999

**Abstract:** The thaloid liverwort *Riccia crozalsii* Levier belonging to the class *Marchantiopsida* (*Hepaticae*) of the division *Bryophyta* is reported for the first time from Turkey, bringing the total number of *Riccia* species known from Turkey to seventeen. The species was collected from Dilek Peninsula National Park near Kuşadası in Aydın

**Key Words:** Liverwort, *Riccia crozalsii*, *Marchantiopsida*, *Hepaticae*.

### Türkiye Florası İçin Yeni Bir Ciğerotu

**Özet:** *Bryophyta* bölümünün *Marchantiopsida* (*Hepaticae*) sınıfında bulunan, talluslu ciğerotlarından *Riccia crozalsii* Levier türü Türkiye'den ilk defa rapor edilmektedir. Böylece ülkemizden bilinen *Riccia* türü sayısı onyediyeye yükselmektedir. Bu tür Aydın'ın Kuşadası İlçesi'ndeki Dilek Yarımadası Milli Parkı'ndan toplanmıştır.

**Anahtar Sözcükler:** Ciğerotu, *Riccia crozalsii*, *Marchantiopsida*, *Hepaticae*.

### Introduction

The genus *Riccia* L. (*Ricciaceae*) is one of the richest genera in the Turkish Liverwort Flora, with 16 species reported up to this time (1, 2). *Ricciocarpos* Corda is the only other Turkish genus of the family, with only one aquatic species collected from Sakarya in 1989 (3).

The first *Riccia* species (*R. bifurca* Hoffm.) was recorded from Turkey in 1905 by Penther and Zederbauer (4). This was followed by *R. macrocarpa* Levier in 1908, reported by Schiffner (5). Two further additions were made by Bornmüller (6) in 1931 and Jovet-Ast (7) in 1957. Thus, prior to 1960, the total number of *Riccia* species recorded was only four. Since 1960, investigations by Jovet-Ast (8), Walther (9), Crundwell and Nyholm (10), Çetin (11) and the authors of the present study (1-3) have increased the number of species to 16 for this genus.

### Materials and Methods

Specimens of bryophytes were collected from Samsun Mountain in the Dilek Peninsula National Park near Kuşadası, Aydın (37° 38' N, 27° 09' E) in 1998 (Fig.1). The material was cleaned with water so as to remove the soil and obtain a clear view of the colour, which is an important feature in the identification of some

bryophytes. The specimens of hepatics werebrought to the laboratory in small polyethylene bags for sorting, then put into special envelopes without pressing, and were then air-dried. Parts of some specimens were kept fresh for the study of their morphological and anatomical characteristics.

Identification was carried out using different published lists as well as floras (1, 8, 11, 12-14). The specimens were deposited in the herbarium of the Department of Biology, Buca Faculty of Education, Dokuz Eylül University.

### Description

*Riccia crozalsii* Levier, Rev. Bryol. 29: 73 (1902)

Plants green or glaucous green to purplish beneath, usually forming partial rosettes (Fig. 2). Thalli 2-3 times dichotomously branched, to 5 mm. long, furrowed lobes 0.4-1 mm. wide, not narrowed to apex, grooved deeply at lobe tips (Fig. 3. 1-6). Lobes thick in t.s., 1-2 times as broad as thick (Fig. 3. 2-6). Epidermal cells of thallus thin-walled, pear-shaped (Fig. 3.7). Thallus finely papillose, margin rounded, cilia present on younger and sometimes older marginal part of plant; cilia 100-300 µm. long, erect to spreading when moist (Fig. 3.8), inflexed over thallus in older parts, surface of cilia




Figure 1. The study site, Dilek Peninsula National Park.



Figure 2. General view of *Riccia crozalsii* specimens (X24).

covered with fine tubercles. Ventral scales colourless to purplish, soon vanishing. Plant monoecious, capsules very common in winter to spring. Spores dark brown to black, 65-105  $\mu\text{m}$ ., convex (distal) face with 6-8 areolae, 12-18  $\mu\text{m}$ . wide, truncate tubercles present at corners of areolae, triradiate (proximal) face with smaller areolae,

wing entire. Chromosome number  $n=7+m$ , 8. It is found on banks, paths, arable fields, mud-capped walls and soil usually near the sea (pH: 4.5-7.5). The species occurs in England, Spain, Portugal, France, Switzerland, Cyprus, Lebanon, Israel, Tunisia, Algeria, Morocco, Australia, New Zealand (8).




Figure 3. *Riccia crozalsii*. 1. General view; 2-6: Transverse sections from different parts of thallus; 7: pear-shaped epidermal cells; 8: cilia. Adapted and drawn by F. Kiremit from Jovet-Ast (8).

## References

- Gökler, I. and Öztürk, M., Liverworts of Turkey and their position in south-west Asia, *Candollea* 46: 359-366 (1991).
- Gökler, I. and Aysel, V., A New Aquatic Liverwort for the Flora of Turkey, *Tr. J. of Bot.* 22: 355-357 (1988).
- Seçmen, Ö., Leblebici, E. and Gökler, I., Türkiye İçin Yeni Bir Ciğerotu: *Ricciocarpus natans* (L.) Corda, *Doğa TU. Bot. D.* 13, 2: 311-313 (1989).
- Penther, A., Zederbauer, E., *Ergebnisse einer naturwissenschaftlichen Reise zum Erdschias-Dagh*, *Ann. Nathist. Hofmus Wien* 20, 385-388 (1905).
- Schiffner, V., *Beitrage zur Kenntnis der Bryophyten von Persien and Lydien*, *Öst. Bot. Zeitschr.* 58, 341-349 (1908).
- Bornmüller, J., *Zur Bryophyten-Flora Kleinasiens*, *Magyar Bot. Lapok* 30, 1-21 (1931).
- Jovet-Ast, S., *Riccia frostii* Aust. au Sahara et en Turquie, *Rev. Bryol. Lichen.* 26, 67-68 (1957).
- Jovet-Ast, S., *Les Riccia de la region Mediterraneenne*, *Crypt. Bryol. Lichen.* 7, 3, 283-431 (1986).
- Walther, K., *Beitrage zur Moosflora Westanatoliens I.*, *Mitt. Staatsinst. Allg. Bot. Hamburg*, 12: 129-188, (1967).
- Crundwell, C. C., Nyholm, E., *Some additions to the bryophyte flora of Turkey I. Hepaticae*, *J. Bryol.*, 10: 479-789, (1979).
- Çetin, B., *Checklist of the Liverworts and Hornworts of Turkey*, *Lindbergia*, 14, 12-14 (1988).
- Smith, A. J. E., *The Liverworts of Britain and Ireland*, Cambridge University Press, (1991).
- Augier, J., *Flore des Bryophytes*, *Cent. Nat. de la Rech. Sci., Paris* (1966).
- Frey, W., Kürschner, H., *Conspectus Bryophytorum Orientalum et Arabicorum*, J. Cramer Verlags. Berlin (1991).