

1-1-2011

A note on weighted $A_p(G)$ -modules

SERAP ÖZTOP

Follow this and additional works at: <https://journals.tubitak.gov.tr/math>



Part of the [Mathematics Commons](#)

Recommended Citation

ÖZTOP, SERAP (2011) "A note on weighted $A_p(G)$ -modules," *Turkish Journal of Mathematics*: Vol. 35: No. 2, Article 7. <https://doi.org/10.3906/mat-0904-26>

Available at: <https://journals.tubitak.gov.tr/math/vol35/iss2/7>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Mathematics by an authorized editor of TÜBİTAK Academic Journals. For more information, please contact academic.publications@tubitak.gov.tr.

A note on weighted $A_p(G)$ -modules

Serap Öztop

Abstract

Let G be a locally compact abelian group and w be a weight function on G . In this paper, we show that the space $A_{p,w}(G)$ is a Banach module over the Figà-Talamanca Herz algebra $A_p(G)$ and study the multiplier space from $A_p(G)$ to $A_{p,w}(G)$.

Key Words: Multiplier, Order free, Essential, Banach module.

1. Introduction

Let G be a locally compact abelian group with Haar measure, w be a weight function on G and $1 < p < \infty$. The work of R. Spector [15] on $A_p(G)$ has motivated us to be interested in the structure theory of weighted $A_p(G)$ denoted by $A_{p,w}(G)$. We show that $A_{p,w}(G)$ is a Banach $A_p(G)$ -module under pointwise multiplication and, as such, a fixed $v \in A_{p,w}(G)$ induces by multiplication an operator T_v from $A_p(G)$ to $A_{p,w}(G)$ defined by $T_v(u) = uv$. Following the work of Friedberg [6] we show that the compact multiplier T_v is trivial if G is a nondiscrete. We also study some multiplier problems from $A_p(G)$ to $A_{p,w}(G)$ spaces.

2. Preliminaries

Let $(A, \|\cdot\|_A)$ be a Banach algebra. A Banach space $(B, \|\cdot\|_B)$ is called a Banach A -module if there exists a continuous algebra representation T of A to $BL(B)$, the algebra of all continuous linear operators from B to B , with $\|T_a\| \leq \|a\|_A$ and $T_{a_1 \bullet a_2} = T_{a_1} \circ T_{a_2}$, where \bullet is the multiplication on A . For $b \in B$, $T_a(b)$ is denoted by $a \bullet b$. Such a module is order-free if 0 is the only $b \in B$ for which $a \bullet b = 0$ for all $a \in A$. A Banach A -module B is called essential if the closed linear span of $A \bullet B$, called the essential part of B and denoted by B_e , coincides with B . If the Banach algebra $(A, \|\cdot\|_A)$ contains a bounded approximate identity, i.e., a bounded net $(u_\alpha)_{\alpha \in I}$ such that $\lim_{\alpha} \|u_\alpha a - a\|_A = 0$ for all $a \in A$, then a Banach A -module B is essential if and only if $\lim_{\alpha} \|u_\alpha b - b\|_B = 0$ for all $b \in B$ [3]. If B is a Banach A -module, then

$$\text{Hom}_A(B) = \{ T \in BL(B) \mid \forall a \in A, \forall b \in B, T(a \bullet b) = a \bullet T(b) \}$$

is the space of all A -module homomorphisms. The elements of $\text{Hom}_A(B)$ are traditionally called the multipliers from B to B .

Let G be a locally compact abelian group with Haar measure dx . A continuous function on G satisfying $w(x) \geq 1$, $w(x+y) \leq w(x)w(y)$ for $x, y \in G$ is called a weight function. Then the space $L_w^p(G) = \{f \mid fw \in L^p(G)\}$, $1 \leq p < \infty$, is a Banach space on G with the norm $\|f\|_{p,w} = \|fw\|_p$, and its dual space is $L_{w^{-1}}^{p'}(G)$, where $\frac{1}{p} + \frac{1}{p'} = 1$. Moreover, if $1 < p < \infty$, then $L_w^p(G)$ is a reflexive Banach space. If $p = 1$, $L_w^1(G)$ is a Banach algebra on G with respect to convolution and contains a bounded approximate identity. It is called a Beurling algebra on G [12]. We know that $L_w^p(G)$ is an essential, order-free Banach $L_w^1(G)$ -module with respect to convolution [11]. We denote by $C_0(G)$ the space of all continuous functions on G vanishing at infinity, and by $C_{0,w}(G)$ the space of functions f on G such that $fw \in C_0(G)$. Throughout we assume that w is even function, i.e., w satisfies $w(x) = w(-x)$ for all $x \in G$.

The following is a classical technique of harmonic analysis.

Proposition 2.1 *Let $f \in L_w^p(G)$ and $g \in L_{w^{-1}}^{p'}(G)$, where $1 < p < \infty$ and $\frac{1}{p} + \frac{1}{p'} = 1$. Then $f * g \in C_{0,w^{-1}}(G)$ and*

$$\|f * g\|_{\infty, w^{-1}} \leq \|f\|_{p,w} \|g\|_{p', w^{-1}}.$$

By Proposition 2.1, a bilinear map b can be defined from $L_w^p(G) \times L_{w^{-1}}^{p'}(G)$ into $C_{0,w^{-1}}(G)$ by

$$b(f, g) = f^\sim * g, \quad f^\sim(x) = f(-x) \quad f \in L_w^p(G), \quad g \in L_{w^{-1}}^{p'}(G)$$

such that $\|b\| \leq 1$. Then b lifts to a linear map B from $L_w^p(G) \otimes_\gamma L_{w^{-1}}^{p'}(G)$, the projective tensor product of $L_w^p(G)$ and $L_{w^{-1}}^{p'}(G)$ considered as a Banach space, into $C_{0,w^{-1}}(G)$ such that $\|B\| \leq 1$ (see [1], [7] and [14]).

Definition 2.2 The range of B consisting of all functions $v = \sum_{i=1}^\infty f_i^\sim * g_i$ on G with $f_i \in L_w^p(G)$, $g_i \in L_{w^{-1}}^{p'}(G)$ and $\sum_{i=1}^\infty \|f_i\|_{p,w} \|g_i\|_{p', w^{-1}} < \infty$ equipped with the norm

$$\|v\|_{A_{p,w}} = \inf \left\{ \sum_{i=1}^\infty \|f_i\|_{p,w} \|g_i\|_{p', w^{-1}} \mid v = \sum_{i=1}^\infty f_i^\sim * g_i \right\}$$

will be denoted by $A_{p,w}(G)$.

The range $A_{p,w}(G)$ is a Banach space of functions on G and can be viewed as a subspace of $C_{0,w^{-1}}(G)$. It can also be identified with the quotient Banach space of $L_w^p(G) \otimes_\gamma L_{w^{-1}}^{p'}(G)$ with K , i.e.,

$$A_{p,w}(G) \cong L_w^p(G) \otimes_\gamma L_{w^{-1}}^{p'}(G) / K,$$

where K is the kernel of the linear form B .

Since $L_w^p(G)$ is a reflexive Banach space and a Banach $L_w^1(G)$ -module with respect to convolution, by [11] we have

$$\text{Hom}_{L_w^1(G)}(L_w^p(G)) \cong \left(L_w^p(G) \otimes_\gamma L_{w^{-1}}^{p'}(G) / K \right)^* \cong K^\perp \cong (A_{p,w}(G))^*.$$

Letting

$$\text{Hom}_G(L_w^p(G)) = \{T \in BL(L_w^p(G)) \mid \forall x \in G, TL_x = L_xT\},$$

we obtain

$$\text{Hom}_G(L_w^p(G)) = \text{Hom}_{L_w^1(G)}(L_w^p(G)).$$

Indeed, $T \in \text{Hom}_{L_w^1(G)}(L_w^p(G))$, and let $(u_\alpha)_{\alpha \in I}$ be an approximate identity of $L_w^1(G)$. Since $L_w^p(G)$ is an essential Banach $L_w^1(G)$ -module, for $f \in L_w^p(G)$, we have

$$\begin{aligned} T(L_x f) &= \lim_\alpha T(u_\alpha * L_x f) = \lim_\alpha ((L_x u_\alpha) * T f), \\ L_x(T f) &= \lim_\alpha L_x(T(u_\alpha * f)) = \lim_\alpha ((L_x u_\alpha) * T f). \end{aligned}$$

Hence $T \in \text{Hom}_G(L_w^p(G))$. Since $L_w^p(G)$ is an order-free Banach $L_w^1(G)$ -module, it is easy to prove the converse. So we obtain

$$\text{Hom}_G(L_w^p(G)) = \text{Hom}_{L_w^1(G)}(L_w^p(G)) \cong (A_{p,w}(G))^*.$$

Let $C_c(G)$ denote the space of continuous functions on G with compact support. Since $C_c(G)$ is dense in $L_w^p(G)$ and in $L_{w-1}^p(G)$, and the convolution of two functions with compact support has compact support, it follows that $C_c(G) \cap A_{p,w}(G)$ is dense in $A_{p,w}(G)$. Also $L_w^p(G)$ and $A_{p,w}(G)$ are essential Banach $L_w^1(G)$ -modules under convolution [11]. Hence, for every $v \in A_{p,w}(G)$ and $x \in G$, $x \rightarrow L_x v$ is a continuous function, where $L_x v(y) = v(y - x)$ for all $y \in G$.

3. $A_{p,w}(G)$ as a Banach $A_p(G)$ -module

It is well known that the Banach algebra $A_p(G)$, called the Figà-Talamanca Herz algebra, has a bounded approximate identity, i.e., there exists $(e_\alpha)_{\alpha \in I} \subset A_p(G)$ such that $\|e_\alpha\|_{A_p} \leq 1$ and $\lim_\alpha \|e_\alpha u - u\|_{A_p} = 0$ for all $u \in A_p(G)$ (see [4], [5], [8] and [15]).

Theorem 3.1 $A_{p,w}(G)$ is an essential, order-free Banach $A_p(G)$ -module under pointwise multiplication.

Proof. Let us consider two functions $u = h^\sim * k$ and $v = f^\sim * g$, where $f, g, h, k \in C_c(G)$. Then

$$\begin{aligned} \|u\|_{A_p} &\leq \|h\|_p \|k\|_{p'} \\ \|v\|_{A_{p,w}} &\leq \|f\|_{p,w} \|g\|_{p',w^{-1}} \\ u(x) &= (h^\sim * k)(x) \\ &= \int_G h(-x - z) k(-z) dz = \int_G h(-x - y - z) k(-y - z) dz \\ v(x) &= (f^\sim * g)(x) \\ &= \int_G f^\sim(x + y) g(-y) dy = \int_G f(-x - y) g(-y) dy \end{aligned}$$

$$\begin{aligned} (uv)(x) &= \int_G \int_G f(-x-y) g(-y) h(-x-y-z) k(-y-z) dy dz \\ &= \int_G A_z(x) dz, \end{aligned}$$

where $A_z = a_z \tilde{*} b_z$ and $a_z, b_z \in C_c(G)$ are given by $a_z(x) = f(x) h(x-z)$, $b_z(x) = g(x) k(x-z)$. Since the mapping $z \rightarrow A_z$ is continuous from G into $A_{p,w}(G)$, the integral $\int_G A_z(x) dz$ is in $A_{p,w}(G)$. By [12], it suffices to show that

$$\int_G \|A_z\|_{A_{p,w}} dz \leq \|f\|_{p,w} \|g\|_{p',w^{-1}} \|h\|_p \|k\|_{p'}.$$

It follows from the Hölder inequality that

$$\begin{aligned} \int_G \|A_z\|_{A_{p,w}} dz &\leq \int_G \|a_z\|_{p,w} \|b_z\|_{p',w^{-1}} dz \leq \\ &\leq \left(\int_G \|a_z\|_{p,w}^p dz \right)^{\frac{1}{p}} \left(\int_G \|b_z\|_{p',w^{-1}} dz \right)^{\frac{1}{p'}} \\ &= \left(\int_G |f(x) w(x)|^p dx \int_G |h(x-z)|^p dz \right)^{\frac{1}{p}} \cdot \\ &\quad \left(\int_G |g(x) w^{-1}(x)|^{p'} dx \int_G |k(x-z)|^{p'} dz \right)^{\frac{1}{p'}} \\ &= \|f\|_{p,w} \|g\|_{p',w^{-1}} \|h\|_p \|k\|_{p'}. \end{aligned}$$

So we obtain that

$$\|uv\|_{A_{p,w}} \leq \|u\|_{A_p} \|v\|_{A_{p,w}},$$

where $u \in A_p(G)$, $v \in A_{p,w}(G)$.

Since $C_c(G) \cap A_{p,w}(G)$ is dense in $A_{p,w}(G)$ and $A_p(G)$ has a bounded approximate identity, it follows that $A_{p,w}(G)$ is an essential Banach module and is order-free, i.e., if $v \in A_{p,w}(G)$ and $uv = 0$ for all $u \in A_p(G)$, then $v = 0$. □

Remark 3.2 Since $A_2(G) \subset A_r(G) \subset A_p(G)$ for $1 < p < r < 2$ or for $2 \leq r < p < \infty$, $A_{p,w}(G)$ is also $A_r(G)$ -module for $1 < p < r \leq 2$ or for $2 \leq r < p < \infty$, that is, $A_r(G) A_{p,w}(G) \subset A_{p,w}(G)$.

In particular, for $r = 2$, since $A_2(G)$ can be identified with $L^1(\hat{G})^\wedge = \mathcal{F}(L^1(\hat{G}))$, where \mathcal{F} denotes the Fourier transform, $A_{p,w}(G)$ has the structure of an $L^1(\hat{G})$ -module, and the action of φ is given by $\varphi^\wedge h \in A_{p,w}(G)$ for $\varphi \in L^1(\hat{G})$, $h \in A_{p,w}(G)$.

4. Multipliers from $A_p(G)$ to $A_{p,w}(G)$

Proposition 4.1 $[Hom_{A_p(G)}(A_p(G), A_{p,\omega}(G))]_e \cong A_{p,\omega}(G)$.

Proof. Since $A_{p,\omega}(G)$ is an essential module, it follows from Theorem 4.5 in [13]. □

Proposition 4.2 If G is compact, then $Hom_{A_p(G)}(A_p(G), A_{p,\omega}(G))$ is an essential $A_p(G)$ -module.

Proof. Since G is compact $A_{p,\omega}(G) = A_p(G)$ and $Hom_{A_p(G)}(A_p(G), A_p(G)) \cong A_p(G)$ by Theorem 5.2 in [10]. Again, using Theorem 4.5 in [13], we get the result.

Since $A_{p,w}(G)$ is a Banach $A_p(G)$ -module, in particular, a fixed $v \in A_{p,w}(G)$ induces a linear operator T_v from $A_p(G)$ into $A_{p,w}(G)$ by means of

$$T_v(u) = uv, \quad u \in A_p(G).$$

It is easy see that T_v is a multiplier from $A_p(G)$ into $A_{p,w}(G)$. □

Remark 4.3 Using the fact of the inclusion $A_{p,\omega}(G) \subset Hom_{A_p(G)}(A_p(G), A_{p,\omega}(G))$, we have the following result.

Proposition 4.4 $A_{p,\omega}(G)$ is dense in $Hom_{A_p(G)}(A_p(G), A_{p,\omega}(G))$ for the strong operator topology on $A_{p,\omega}(G)$.

Proof. Let $(e_\alpha)_{\alpha \in I}$ be a bounded approximate identity for $A_p(G)$ and let $T \in Hom_{A_p(G)}(A_p(G), A_{p,\omega}(G))$. Put $u_\alpha = T(e_\alpha)$. Since $A_{p,\omega}(G)$ is an essential Banach $A_p(G)$ module for every $u \in A_p(G)$, we can write

$$\|T_{u_\alpha}(u) - T(u)\| = \|u_\alpha u - T(u)\| = \|(Te_\alpha)u - T(u)\| = \|(Tu)e_\alpha - T(u)\| \rightarrow 0.$$

for all $u \in A_p(G)$. □

Theorem 4.5 *If G is a nondiscrete locally compact abelian group and $T_v : A_p(G) \rightarrow A_{p,w}(G)$ is a compact multiplier, then T_v is trivial.*

Proof. Let V be an open set in G whose closure is compact. As it is well known [9], for every $x \in V$, there exists $u \in A_p(G)$ such that 1) $u(x) = 1$; 2) $\text{supp } u \subset V$; 3) $\|u\|_{A_p} = 1$. Now assume that $v \neq 0$. Then there exists an open set V with compact closure and $\delta > 0$ such that $|v(x)| \geq \delta$ for all $x \in V$. Let (V_n) be a sequence of disjoint open sets in V such that the closure of any $V_n (n = 1, 2, \dots)$ is compact. Choose a sequence (x_n) such that $x_n \in V_n (n = 1, 2, \dots)$. As we already noted above, there exists a sequence (u_n) in $A_p(G)$ such that $u_n(x_m) = \delta_{nm}$ (here, δ_{nm} is the Kronecker symbol) and $\|u_n\|_{A_p} = 1$.

For $n \neq m$ we have

$$\begin{aligned} \|T_v u_n - T_v u_m\|_{A_{p,w}} &\geq \|v u_n - v u_m\|_{\infty, w^{-1}} \\ &\geq \sup_k \left(|v(x_k) u_n(x_k) - v(x_k) u_m(x_k)| \frac{1}{w(x_k)} \right) \\ &\geq \delta \inf \left\{ \frac{1}{w(x)} \mid x \in \overline{V} \right\} \sup_k |u_n(x_k) - u_m(x_k)| \\ &= \inf \left\{ \frac{1}{w(x)} \mid x \in \overline{V} \right\}. \end{aligned}$$

This contradicts compactness of T_v . □

Let us note that we have the inclusion

$$\text{Hom}_{A_p(G)}(A_p(G), E) \subset \text{Hom}_{A_p(G)}(A_p(G), A_{p,w}(G))$$

whenever E is a Banach $A_p(G)$ -module contained in $A_{p,w}(G)$. Using the method in [2], we get the following theorem.

Theorem 4.6 *Let E be a Banach $A_p(G)$ -module contained in $A_{p,w}(G)$. If $T \in \text{Hom}_{A_p(G)}(A_p(G), A_{p,w}(G))$ such that the net $T(e_\alpha)$ converges in E , where $(e_\alpha)_{\alpha \in I}$ is a bounded approximate identity of $A_p(G)$, then T is a multiplier from $A_p(G)$ to E .*

Proof. For $u \in A_p(G)$, consider (x_α) defined by $x_\alpha = e_\alpha u T(e_\alpha)$. Since $T \in \text{Hom}_{A_p(G)}(A_p(G), A_{p,w}(G))$, we have

$$\begin{aligned} \|x_\alpha - x_\beta\|_E &= \|e_\alpha u T(e_\alpha) - e_\beta u T(e_\beta)\|_E \\ &\leq \|e_\alpha u T(e_\beta) - e_\beta u T(e_\beta)\|_E + \|e_\alpha u T(e_\alpha) - e_\alpha u T(e_\beta)\|_E \\ &\leq \|e_\alpha u - e_\beta u\|_{A_p} \|T(e_\beta)\|_E + \|e_\alpha u\|_{A_p} \|T(e_\alpha) - T(e_\beta)\|_E \rightarrow 0, \end{aligned}$$

which shows that (x_α) is a Cauchy net in E . Hence for $u \in A_p(G)$, the net $x_\alpha = e_\alpha u T(e_\alpha)$ converges to an element x in E . Now we want to show that x is equal to Tu as an element in $A_{p,w}(G)$.

Let $h \in A_p(G)$. Then $h x_\alpha = (h e_\alpha)(u T(e_\alpha)) = (h e_\alpha)(T(u e_\alpha)) \rightarrow h T u$ in $A_{p,w}(G)$. On the other hand, since $x_\alpha \rightarrow x$ in E , we have $h x_\alpha \rightarrow h x$ in $A_{p,w}(G)$. Hence we have that $x = T u$. □

It is easy to obtain the following corollary.

Corollary 4.7 *Let E be a Banach $A_p(G)$ -module contained in $A_{p,w}(G)$ and let*

$A = \{ T \in \text{Hom}_{A_p(G)}(A_p(G), A_{p,w}(G)) \mid (T(e_\alpha)) \text{ converges in } E \}$, where $(e_\alpha)_{\alpha \in I}$ is a bounded approximate identity of $A_p(G)$. Then the multiplier space $\text{Hom}_{A_p(G)}(A_p(G), E)$ is identified with the subspace A of $\text{Hom}_{A_p(G)}(A_p(G), A_{p,w}(G))$.

Similarly, if we consider that $A_{p,w}(G)$ is a Banach $L_w^1(G)$ -module under convolution, we get the following theorem.

Theorem 4.8 *Let $E \subseteq A_{p,w}(G)$ two Banach $L_w^1(G)$ -modules then $T \in \text{Hom}_{L_w^1(G)}(L_w^1(G), A_{p,w}(G))$ is a multiplier from $L_w^1(G)$ to E if and only if $(Te_\alpha) \in E$ and $\sup_\alpha \|(Te_\alpha)\|_E < \infty$ where $(e_\alpha)_{\alpha \in I}$ is a bounded approximate identity of $L_w^1(G)$.*

Acknowledgement

The author would like to thank the referees for carefully reading the manuscript.

References

- [1] Bonsall, F.F., Duncan, J. : *Complete Normed Algebras*, Springer-Verlag, Berlin and New York, 1973.
- [2] Datry, C., Muraz, G., Quek, T.S. and Yap, L.Y.H. : *Homomorphismes de $L^1(G)$ -modules*, prépublication de l'Institut Fourier **76** (1987).
- [3] Doran, R. S., Wichmann, J. : *Approximate Identities and Factorization in Banach Modules*, Lecture Notes in Mathematics **768**, Springer-Verlag, Berlin and New York , 1979.
- [4] Eymard, P. : *L'algèbre de Fourier d'un groupe localement compact*, Bull. Soc. Math.France **92**, 181-236 (1964).
- [5] Figà-Talamanca, A. : *Multipliers of p -integrable functions*, Bull. Amer. Math. Soc. **70**, 666-669 (1964).
- [6] Friedberg, S.H. : *Compact multipliers on Banach algebras*, Proc. Amer. Math. Soc. **77** , 210 (1979).
- [7] Grothendieck, A. : *Produits tensoriels topologiques et espaces nucléaires*, Mem. Amer. Math. Soc.**16**, Providence, (1955).
- [8] Herz, C. : *The theory of p -spaces with an application to convolution operators*, Trans. Amer. Math. Soc. **154**, 69-82 (1971).
- [9] Herz, C. : *Harmonic synthesis for subgroups*, Ann. Inst. Fourier (Grenoble) **23**, 91-123 (1973).
- [10] Lai, H. and Chen, I. *Harmonic analysis on the Fourier algebra $A_{1,p}(G)$* , J. Austral. Math. Soc. (Series A)**30**, 438-452 (1981).
- [11] Öztop, S. and Gürkanlı A. T. : *Multipliers and tensor products of weighted L^p -spaces*, Acta. Math. Sci. Ser. B **21**, 41-49 (2001).
- [12] Reiter, H. : *Classical Harmonic Analysis and Locally Compact Groups*, Oxford Univ. Press, 1968.
- [13] Rieffel, M.A. : *Induced Banach representations of Banach algebras and locally compact groups*, J. Funct. Anal. **1**, 443-491 (1967).
- [14] Rieffel, M.A. : *Multipliers and tensor products of L^p -spaces of locally compact groups*, Studia Math. **33**, 71-82 (1969).
- [15] Spector, R. : *Sur la structure locale des groupes abéliens localement compacts*, Bull. Soc. Math. France, Mémoire **24** (1970).

Serap ÖZTOP
 İstanbul University, Faculty of Science,
 Department of Mathematics,
 34134 Vezneciler, İstanbul-TURKEY
 e-mail: oztops@istanbul.edu.tr

Received: 16.04.2009