

1-1-2011

## Invariant subspace problem for positive L-weakly and M-weakly compact operators

CEVRİYE TONYALI

ERDAL BAYRAM

Follow this and additional works at: <https://dctubitak.researchcommons.org/math>



Part of the [Mathematics Commons](#)

---

### Recommended Citation

TONYALI, CEVRİYE and BAYRAM, ERDAL (2011) "Invariant subspace problem for positive L-weakly and M-weakly compact operators," *Turkish Journal of Mathematics*: Vol. 35: No. 2, Article 8. <https://doi.org/10.3906/mat-0903-37>

Available at: <https://dctubitak.researchcommons.org/math/vol35/iss2/8>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Mathematics by an authorized editor of TÜBİTAK Academic Journals.

# Invariant subspace problem for positive $L$ -weakly and $M$ -weakly compact operators

*Cevriye Tonyalı and Erdal Bayram*

## Abstract

In this paper, we show that positive  $L$ -weakly and  $M$ -weakly compact operators on some real Banach lattices have a non-trivial closed invariant subspace. Also, we prove that any positive  $L$ -weakly (or  $M$ -weakly) compact operator  $T : E \rightarrow E$  has a non-trivial closed invariant subspace if there exists a Dunford-Pettis operator  $S : E \rightarrow E$  satisfying  $0 \leq T \leq S$ , where  $E$  is Banach lattice.

**Key word and phrases:** Invariant subspace,  $L$ - and  $M$ -weakly compact operator, Polynomially  $L$ -weakly ( $M$ -weakly) compact operator, Dunford-Pettis operator.

## 1. Introduction

Bulk of the papers in Banach lattice theory concern the open problem every positive operator on a Banach lattice of dimension at least two has a non-trivial closed invariant subspace. The problem was solved for positive compact operators on Banach lattices [1], [7], [12].

Our objective in this work is to investigate whether or not every positive  $L$ -weakly and  $M$ -weakly operator on a real Banach lattice does possess a non-trivial closed invariant subspace. First, we will prove that every  $L$ -weakly compact operator on a Banach lattice without order continuous norm has a non-trivial closed invariant subspace. Also, we shall show that if  $E$  is a Banach lattice such that either  $E$  and  $E'$  has order continuous norm, then every bounded operator that commutes with a positive  $L$ -weakly ( $M$ -weakly) compact operator that on  $E$  has a non-trivial closed invariant subspace. Furthermore, we will investigate invariant subspaces of polynomially  $L$ -weakly ( $M$ -weakly) compact operators for Banach lattices without order continuous norm. Next, we will prove that every bounded operator that commutes with a positive  $M$ -weakly (or positive  $L$ -weakly) compact operator on a Banach lattice  $E$  has a non-trivial closed invariant subspace if it is dominated by a Dunford-Pettis operator. Also, we will see that any positive operator on a Banach lattice of which order dual has order continuous norm has a non-trivial closed invariant subspace if it is dominated by a Dunford-Pettis operator.

**2. Main results**

Throughout this paper, unless otherwise state,  $E$  will denote an infinite dimensional separable real Banach lattice with norm dual  $E'$  and all operators on Banach lattices will be assumed to be non-scalar and non-zero. In the rest of this article, by the term “an operator” between two Banach lattices, we shall mean “a linear norm bounded operator”. We refer the reader to [5], [14] and [16] for any unexplained terms from Banach lattice theory and for further details on the theory of invariant subspaces see [1], [2], [15].

Recall that a non-empty bounded subset  $A$  of Banach lattice  $E$  is said to be  $L$ -weakly compact if  $\|x_n\| \rightarrow 0$  for every disjoint sequence  $(x_n)$  in the solid hull of  $A$ . A bounded linear operator  $T$  from a Banach space  $X$  into  $E$  is called  $L$ -weakly compact if  $T(U_X)$  is  $L$ -weakly compact in  $E$ , where  $U_X$  denotes the closed unit ball of  $X$ . A bounded linear operator from  $E$  into  $X$  is  $M$ -weakly compact if  $\|Tx_n\| \rightarrow 0$  as  $n \rightarrow \infty$  for every disjoint sequence  $(x_n)$  in  $U_E$ . That  $L$ -weakly compact and  $M$ -weakly compact operators are weakly compact operators was shown by P. Meyer-Nieberg [13]. It is known that

$$E^a = \{x \in E : \text{every monoton sequence in } [0, |x|] \text{ is convergent}\}$$

is the maximal closed order ideal in  $E$  on which the induced norm is order continuous and it is known that any  $L$ -weakly compact subset is contained in  $E^a$  ([14], p. 92 and p. 212). Recall that a Banach lattice  $E$  is said to have an *order continuous norm* if  $x_\alpha \downarrow 0$  in  $E$  implies  $\|x_\alpha\| \downarrow 0$ .

**Proposition 1** *If  $E$  is a Banach lattice without order continuous norm, then every  $L$ -weakly compact operator  $T : E \rightarrow E$  has a non-trivial closed invariant ideal.*

**Proof.** If  $E$  does not have order continuous norm,  $E^a \neq E$  since  $E^a$  has order continuous norm. Take any  $x \in E^a$  (we can suppose that  $\|x\| \leq 1$  without loss of generality).  $T(U_E) \subset E^a$  since  $T$  is an  $L$ -weakly compact operator and every  $L$ -weakly compact subset is contained in  $E^a$ . Therefore  $T(x) \in E^a$  i.e. ,  $T(E^a) \subseteq E^a$ . If  $E^a = \{0\}$  then  $T(U_E) = \{0\}$  which implies that  $T = 0$ . Since we take  $T$  different from zero operator,  $E^a \neq \{0\}$ . Hence  $E^a$  is the invariant ideal for  $T$  which we are looking for. □

**Corollary 1** *Let  $E$  be a Banach lattice without order continuous norm, let  $E^a \neq \{0\}$  and let  $T : E \rightarrow E$  be a non-scalar regular operator. If there exists some element  $0 < x_0 \in E^a$  and  $n \in \mathbb{N}$  such that  $T^n x_0 \neq 0$  and  $T^n : E \rightarrow E$  is an  $L$ -weakly compact operator, then every regular operator on  $E$  has a non-trivial invariant closed order ideal.*

**Proof.** Assume that  $T^n : E \rightarrow E$  is an  $L$ -weakly compact operator for some  $n \in \mathbb{N}^+$  and  $S : E \rightarrow E$  is any regular operator. Since in  $\mathcal{L}^r(E)$ , the space of the regular operators on  $E$ , the  $L$ -weakly compact regular operators form closed two-sided ideal,  $S^m T^n : E \rightarrow E$  is an  $L$ -weakly compact operator for each  $m \in \mathbb{N}^+$ . Thus,  $S^m T^n (x_0) \in E^a$  for all  $m \in \mathbb{N}^+$ . Next, choose the closed order ideal  $W$  generated by

$$\{T^n x_0, ST^n x_0, S^2 T^n x_0, \dots, S^m T^n x_0, \dots\}.$$

Hence,  $S(W) \subseteq W$ , it follows that  $W$  is a non-trivial invariant closed order ideal for  $S$ . □

The order ideal generated by an element  $0 < x \in E$  is precisely

$$I_x = \{y \in E : \exists \lambda > 0 \text{ with } |y| \leq \lambda x\}.$$

For every  $z \in I_x$ ,

$$\|z\|_\infty = \inf \{\lambda > 0; |z| \leq \lambda x\}$$

defines a norm on  $I_x$ . Thus,  $(I_x, \|\cdot\|_\infty)$  is an  $AM$ -space and, moreover, its closed unit ball is the interval  $[-x, x]$ , see [4],[5],[14].

**Theorem 1** *If  $T : E \rightarrow E$  is a positive  $L$ -weakly compact operator on a Banach lattice  $E$  with order continuous norm, then the operator  $T^2$  is compact.*

**Proof.** If  $T : E \rightarrow E$  is a positive  $L$ -weakly compact operator, then for each  $n \in \mathbb{N}^+$  there exists some  $0 < u_n \in E$  lying in the order ideal generated by  $T(E)$  satisfying

$$\left\| (|Tx| - u_n)^+ \right\| < n^{-1}$$

for all  $x \in U_E$  ([5], Th. 18.9, p. 313). From the identity  $|Tx| = |Tx| \wedge u_n + (|Tx| - u_n)^+$  we see that

$$T(U_E^+) \subseteq [0, u_n] + n^{-1}U_E \tag{2.1}$$

for all  $n \in \mathbb{N}^+$ . Let  $0 < y = \sum_{n=1}^\infty \frac{1}{2^n \|u_n\|} u_n \in E^+$  and let  $I_y$  be the order ideal generated by  $y$ . The restriction  $T|_{I_y} : (I_y, \|\cdot\|_\infty) \rightarrow E$  is a positive  $L$ -weakly compact operator, and  $(I_y, \|\cdot\|_\infty)'$  is an  $AL$ -space ([5], Th. 18.11, p. 315). Thus, since  $I_y$  is an  $AM$ -space, it satisfies Dunford-Pettis property and so  $T|_{I_y}$  is a Dunford-Pettis operator by ([5], Theorems 19.4 and 19.6). Moreover we know that  $I_y'$  is an  $AL$ -space with order continuous norm and  $E$  has order continuous norm by hypothesis, therefore,  $T|_{I_y} : I_y \rightarrow E$  is compact ([6], Theorem 2.12(2)ii). Let  $\alpha_n = 2^n \|u_n\|$  for every  $n \in \mathbb{N}^+$ . From the inequality  $u_n \leq \alpha_n y$  and (2.1) we obtain that

$$T^2(U_E^+) \subseteq T[0, u_n] + n^{-1} \|T\| U_E \subseteq \alpha_n T[0, y] + n^{-1} \|T\| U_E$$

and

$$T^2(U_E^+) \subseteq \alpha_n T[0, y] + n^{-1} \|T\| U_E = \alpha_n T|_{I_y}[0, y] + n^{-1} \|T\| U_E.$$

Since  $\{n^{-1} \|T\| U_E\}$  is a norm-neighborhood system at zero,  $T^2(U_E^+)$  is a norm-totally bounded set, from which it follows that the operator  $T^2$  is compact. □

**Corollary 2** *Every  $L$ -weakly compact operator on a Banach lattice has a non-trivial closed invariant subspace.*

As an immediate consequence of Theorem 1, we obtain the following result:

**Corollary 3** *If  $T$  is a positive  $M$ -weakly compact operator on a Banach lattice  $E$  such that  $E'$  has order continuous norm, then  $T^2$  is compact.*

**Proof.** It is enough to consider that  $T'$  is a positive  $L$ -weakly compact operator. □

**Theorem 2** *Let  $E$  and  $F$  be a Banach lattice and  $T : E \rightarrow F$  be a positive  $M$ -weakly compact operator. If  $F$  has order continuous norm, then the operator  $T$  is compact.*

**Proof.** If  $T : E \rightarrow F$  is a positive  $M$ -weakly compact operator, then for each  $n \in \mathbb{N}^+$  there exists some  $0 < u_n \in E$  such that

$$\left\| T(|x| - u_n)^+ \right\| < n^{-1}$$

holds for all  $x \in U_E$  ([5], Th. 18.9, p.313).

Let  $y = \sum_{n=1}^{\infty} \frac{1}{2^n \|u_n\|} u_n \in E^+$  and let  $I_y$  be the order ideal generated by  $y$ . It is clear that  $y$  is well defined. Since the operator  $T : E \rightarrow F$  is  $M$ -weakly compact, the operator  $T|_{I_y} : (I_y, \|\cdot\|_{\infty}) \rightarrow F$  is  $M$ -weakly compact, and so  $T|_{I_y}$  is compact by Theorem 2.12. in [6]. Let  $\alpha_n$  be as mentioned in the proof of above theorem. From the inequality  $u_n \leq \alpha_n y$  and the identity  $|x| = |x| \wedge u_n + (|x| - u_n)^+$  we see that

$$T(U_E^+) \subseteq T[0, u_n] + n^{-1}U_F \subseteq \alpha_n T[0, y] + n^{-1}U_F.$$

Thus, we obtain that

$$T(U_E^+) \subseteq \alpha_n T[0, y] + n^{-1}U_F = \alpha_n T|_{I_y}[0, y] + n^{-1}U_F.$$

Moreover,  $T|_{I_y}[0, y] \subset F$  is a norm-totally bounded set since  $T|_{I_y}$  is a compact operator. Hence,  $T(U_E^+)$  is also a norm-totally bounded set because  $\{n^{-1}U_F\}$ ,  $n \in \mathbb{N}^+$  is a norm-neighborhood system at zero. Therefore,  $T$  is a compact operator, as desired. □

Since the dual of an  $L$ -weakly compact operator is  $M$ -weakly compact, we can state the following result:

**Corollary 4** *Let  $T : E \rightarrow F$  be a positive  $L$ -weakly compact operator between Banach lattices. If  $E'$  has order continuous norm, then  $T$  is compact.*

We obtain another consequence of Theorem 1 and 2:

**Corollary 5** *If  $E$  is a Banach lattice such that either  $E$  or  $E'$  has order continuous norm, then every bounded operator that commutes with a positive  $L$ -weakly (positive  $M$ -weakly) compact operator on  $E$  has a non-trivial closed invariant subspace.*


Let  $T$  be a bounded operator on a Banach lattice  $E$ .  $T$  is said to be a *polynomially  $L$ -weakly (polynomially  $M$ -weakly) compact operator* whenever there exists a non-zero polynomial  $p$  such that  $p(T)$  is  $L$ -weakly ( $M$ -weakly) compact. It is clear that every  $L$ -weakly compact operator on a Banach lattice is polynomially  $L$ -weakly compact, but sometimes the converse of this statement is false even for  $AL$ -spaces. The following example illustrates this point.

**Example 1** Define an operator  $T : L^1 [0, 1] \rightarrow L^1 [0, 1]$  by

$$Tf(x) := \begin{cases} 0 & \text{if } 0 \leq x \leq 1/2 \\ f(x - \frac{1}{2}) & \text{if } 1/2 < x \leq 1, \end{cases}$$

then  $T$  is not weakly compact, but  $T$  is polynomially  $L$ -weakly compact since  $T^2 = 0$ .

The following diagram is held for all operators defined on  $AL$ -spaces.



Again, we will consider Banach lattices in different properties while we seek invariant subspaces for polynomially  $L$ -weakly compact operators.

**Theorem 3** Every polynomially  $L$ -weakly compact operator on a Banach lattice without order continuous norm has a non-trivial closed invariant subspace.

**Proof.** Let  $E$  be a Banach lattice without order continuous norm and  $T : E \rightarrow E$  be a polynomially  $L$ -weakly compact operator. Choose a non-zero polynomial  $p(t) = a_0 + a_1t + a_2t^2 + \dots + a_{n-1}t^{n-1} + a_nt^n$  such that  $p(t)$  is  $L$ -weakly compact.

Assume that  $p(T) = 0$  (In this case, there exists a non-zero element  $x$  in  $E$  such that  $p(T)(x) = 0$ ). Let  $V$  denote the non-zero closed subspace generated by the set  $\{x, Tx, \dots, T^{n-1}x\}$ . Since  $E$  is infinite dimensional, we have  $V \neq E$  and we can see easily that  $V$  is  $T$ -invariant.

Now suppose that  $p(T) \neq 0$ . Fix any non-zero vector  $x_0 \in E^a$  such that  $p(T)(x_0) \neq 0$ . Since  $p(T)$  is  $L$ -weakly compact, we have  $p(T)(E^a) \subset E^a$  by Prop.1. Moreover, for each  $k = 0, 1, 2, \dots$ ,  $T^k p(T)(E^a) = p(T)T^k(E^a) \subset E^a$  holds because of  $L$ -weakly compactness of  $p(T)T^k$ . Let  $V$  be the non-zero closed subspace generated by the set  $\{p(T)(x_0), Tp(T)(x_0), T^2p(T)(x_0), \dots, T^k p(T)(x_0), \dots\} \subset E^a$ . Since  $E$  does not have order continuous norm, again we have  $V \neq E$ , and it can be seen that  $V$  is  $T$ -invariant.  $\square$

**Remark 1** Not every operator  $p(T)$ , where  $p$  is a polynomial, is positive when  $T$  is positive. Thus we cannot say that every polynomially  $L$ -weakly compact operator on a Banach lattice with order continuous norm has a non-trivial invariant subspace.

Let  $X, Y$  be Banach spaces and let  $E$  be a Banach lattice. A linear operator  $T : E \rightarrow X$  is called  $AM$ -compact if  $T[-x, x]$  is relatively compact for every  $x \in E^+$ . And, we say that  $T : X \rightarrow Y$  is a *Dunford-Pettis operator* whenever  $x_n \xrightarrow{w} 0$  in  $X$  implies  $\lim \|Tx_n\| = 0$  [5], [10], [11], [14]. The  $o$ -weakly compact operators have been characterized by P.G. Dodds [8]. Recall that a continuous operator  $T : E \rightarrow X$  is  $o$ -weakly compact whenever  $T[0, x]$  is a relatively weakly compact subset of  $X$  for each  $0 < x \in E$ . It is clear that every weakly compact operator is  $o$ -weakly compact.

**Theorem 4** *Let  $E$  be a Banach lattice and  $T : E \rightarrow E$  be a positive  $M$ -weakly compact operator. If there exists a Dunford-Pettis operator  $S : E \rightarrow E$  such that  $0 \leq T \leq S$ , then the operator  $T^3$  is compact.*

**Proof.** Assume that  $T : E \rightarrow E$  is a positive  $M$ -weakly compact operator and  $S : E \rightarrow E$  is a Dunford-Pettis operator satisfying  $0 \leq T \leq S$ . Thus,  $T^2 : E \rightarrow E$  is a Dunford-Pettis operator by ([5], Cor. 19.15, p. 340). Moreover, the operator  $T$  is  $o$ -weakly compact because it is an  $M$ -weakly compact operator ([5], p. 311). Thus,  $T[0, x]$  is a relatively weakly compact set for every  $x \in E^+$ . Therefore,  $T^2(T[0, x]) = T^3[0, x]$  is a norm-totally bounded set for every  $x \in E^+$  since the Dunford-Pettis operator  $T^2$  carries relatively weakly compact subsets of  $E$  onto norm-totally bounded subsets of  $E$  ([5], Th. 19.3, p. 334). It follows that the  $M$ -weakly compact operator  $T^3$  is  $AM$ -compact. This implies that  $T^3$  is a compact operator ([14], Prop. 3.7.4, p.219).  $\square$

**Corollary 6** *Let  $E$  be a Banach lattice and  $T : E \rightarrow E$  be a positive  $M$ -weakly compact operator. If there exists a Dunford-Pettis operator  $S : E \rightarrow E$  such that  $0 \leq T \leq S$ , then every bounded operator that commutes with  $T$  has a non-trivial closed invariant subspace.*

We can state a similar theorem to the previous theorem for  $L$ -weakly compact operators.

**Theorem 5** *Let  $E$  be a Banach lattice and  $T : E \rightarrow E$  be a positive  $L$ -weakly compact operator. If there exists a Dunford-Pettis operator  $S : E \rightarrow E$  satisfying  $0 \leq T \leq S$ , then  $T^4$  is a compact operator.*

**Proof.** We already known that  $T^2$  is a Dunford-Pettis operator and the operator  $T$  is  $o$ -weakly compact as mentioned in Theorem 4. Moreover, the Dunford-Pettis operator  $T^2$  carries the relatively weakly compact sets  $T[0, x]$  for each  $x \in E^+$  onto norm-totally bounded sets  $T^3[0, x]$ . Since the operator  $T$  is a  $L$ -weakly compact, for each  $\varepsilon > 0$  there exists some  $x_\varepsilon \in E^+$  such that

$$T(U_E^+) \subseteq [0, x_\varepsilon] + \varepsilon U_E.$$

Thus, we obtain that

$$T^4(U_E^+) \subseteq T^3[0, x_\varepsilon] + \varepsilon \|T\|^3 U_E,$$

and so,  $T^4(U_E^+)$  is a norm-totally bounded set because  $\{\varepsilon \|T\|^3 U_E\}$  is a base at zero.  $\square$

**Corollary 7** *Let  $E$  be a Banach lattice and  $T : E \rightarrow E$  be a positive  $L$ -weakly compact operator. If there exists a Dunford-Pettis operator  $S : E \rightarrow E$  satisfying  $0 \leq T \leq S$ , then every bounded operator that commutes with  $T$  has a non-trivial closed invariant subspace.*

## References

- [1] Abramovich, Y.A.; Aliprantis, C.D., An Invitation to Operator Theory, Graduate Texts in Math, American Math. Soc., Providence, R.I., (2002).
- [2] Abramovich, Y.A.; Aliprantis, C.D., Problems in Operator Theory, Graduate Texts in Math, Vol.51, American Math. Soc., Providence, R.I., (2002).

- [3] Aliprantis, C.D.; Burkinshaw, Dunford-Pettis operators on Banach lattices, *Tras. Amer. Math. Soc.*, 274(1), 227-238, (1982).
- [4] Aliprantis, C.D.; Burkinshaw, O., *Locally Solid Riesz Spaces*, Academic Pres, New-York-London, (1978).
- [5] Aliprantis, C.D.; Burkinshaw, O., *Positive Operators*, Academic Press, New york-London, (1985).
- [6] Aqzzouz, B., Nouira, R.; Zraoula, L., About positive Dunford-Pettis operators on Banach lattices, *Journal Of Math. Analysis And Appl.*, 324: 49-59, (2006).
- [7] Aronszajn, N.; Smith, K.T., Invariant subspaces of completely continuous operators, *Ann. Of Math.*, 60, 345-350, (1954).
- [8] Dodds, P.G.,  $\sigma$ -weakly compact mappings of Riesz spaces, *Trans. Amer. Math. Soc.*, 214, 389-402, (1975).
- [9] Dodds, P.G.; Fremlin, D.H., Compact operators in Banach lattices, *Israel J. Math.*, 34, 287-320, (1979).
- [10] Dunford, N.; Pettis, P.J., Linear operations on summable functions, *Trans. Amer. Math. Soc.*, 47, 323-392, (1940).
- [11] Grothendieck, A., Sur les applications lineaires faiblement compactes d'espaces du type  $C(K)$ , *Canad. J. Math.*, 5, 129-173, MR 15, 438, (1953).
- [12] Lomonosov, V.I., Invariant subspaces of the family of operators that commute with a completely continuous operator, *Funktsional. Anal. i Prilozhen*, 7, No:3, 55-56, (1973).
- [13] Meyer-Nieberg, P., Über klassen schwach kompakter operatoren in Banachverbänden, *Math. Z.*, 138, 145-159, (1974).
- [14] Meyer-Nieberg, P., *Banach Lattices*, Springer-Verlag, Berlin-Heidelberg, (1991).
- [15] Radjavi, H.; Rosenthal, P., *Invariant Subspaces*, Edition, Dover, Mineola, New York, (2003).
- [16] Schaefer, H.H., *Banach Lattices and Positive Operators*, Springer-Verlag, Berlin, New York, (1974).

Cevriye TONYALI, Erdal BAYRAM  
 Department of Mathematics  
 Faculty of Science and Arts, Gazi Universty  
 Ankara-TURKEY  
 e-mail: bayramer@gmail.com

Received: 25.03.2009