

1-1-2010

On Abelian Rings

NAZIM AGAYEV

ABDULLAH HARMANCI

SAİT HALICIOĞLU

Follow this and additional works at: <https://dctubitak.researchcommons.org/math>



Part of the [Mathematics Commons](#)

Recommended Citation

AGAYEV, NAZIM; HARMANCI, ABDULLAH; and HALICIOĞLU, SAİT (2010) "On Abelian Rings," *Turkish Journal of Mathematics*: Vol. 34: No. 4, Article 4. <https://doi.org/10.3906/mat-0711-1>
Available at: <https://dctubitak.researchcommons.org/math/vol34/iss4/4>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Mathematics by an authorized editor of TÜBİTAK Academic Journals.

On Abelian Rings

Nazim Agayev, Abdullah Harmancı, Sait Halıcıoğlu

Abstract

Let α be an endomorphism of an arbitrary ring R with identity. In this note, we introduce the notion of α -abelian rings which generalizes abelian rings. We prove that α -reduced rings, α -symmetric rings, α -semicommutative rings and α -Armendariz rings are α -abelian. For a right principally projective ring R , we also prove that R is α -reduced if and only if R is α -symmetric if and only if R is α -semicommutative if and only if R is α -Armendariz if and only if R is α -Armendariz of power series type if and only if R is α -abelian.

Key word and phrases: α -reduced rings, α -symmetric rings, α -semicommutative rings, α -Armendariz rings, α -abelian rings.

1. Introduction

Throughout this paper R denotes an associative ring with identity 1 and α denotes a non-zero and non-identity endomorphism of a given ring with $\alpha(1) = 1$, and $\mathbf{1}$ denotes identity endomorphism, unless specified otherwise.

We write $R[x]$, $R[[x]]$, $R[x, x^{-1}]$ and $R[[x, x^{-1}]]$ for the polynomial ring, the power series ring, the Laurent polynomial ring and the Laurent power series ring over R , respectively. Consider

$$R[x, \alpha] = \left\{ \sum_{i=0}^s a_i x^i : s \geq 0, a_i \in R \right\},$$

$$R[[x, \alpha]] = \left\{ \sum_{i=0}^{\infty} a_i x^i : a_i \in R \right\},$$

$$R[x, x^{-1}, \alpha] = \left\{ \sum_{i=-s}^t a_i x^i : s \geq 0, t \geq 0, a_i \in R \right\},$$

$$R[[x, x^{-1}, \alpha]] = \left\{ \sum_{i=-s}^{\infty} a_i x^i : s \geq 0, a_i \in R \right\}.$$

Each of these is an abelian group under an obvious addition operation. Moreover, $R[x, \alpha]$ becomes a ring under the following product operation:

$$\text{For } f(x) = \sum_{i=0}^s a_i x^i, g(x) = \sum_{i=0}^t b_i x^i \in R[x, \alpha]$$

$$f(x)g(x) = \sum_{k=0}^{s+t} \left(\sum_{i+j=k} a_i \alpha^i(b_j) \right) x^k.$$

Similarly, $R[[x, \alpha]]$ is a ring. The rings $R[x, \alpha]$ and $R[[x, \alpha]]$ are called the *skew polynomial extension* and the *skew power series extension of R* , respectively. If $\alpha \in \text{Aut}(R)$, then with a similar scalar product, $R[[x, x^{-1}, \alpha]]$ (resp. $R[x, x^{-1}, \alpha]$) becomes a ring. The rings $R[x, x^{-1}, \alpha]$ and $R[[x, x^{-1}, \alpha]]$ are called the *skew Laurent polynomial extension* and the *skew Laurent power series extension of R* , respectively.

In [8], *Baer rings* are introduced as rings in which the right(left) annihilator of every nonempty subset is generated by an idempotent. According to Clark [4], a ring R is said to be *quasi-Baer ring* if the right annihilator of each right ideal of R is generated(as a right ideal) by an idempotent. These definitions are left-right symmetric. A ring R is called *right principally quasi-Baer ring (or simply, right p.q.-Baer ring)* if the right annihilator of a principally right ideal of R is generated by an idempotent. Finally, a ring R is called *right principally projective ring (or simply, right p.p.-ring)* if the right annihilator of an element of R is generated by an idempotent [2].

2. Abelian Rings

In this section the notion of an α -abelian ring is introduced as a generalization of an abelian ring. We show that many results of abelian rings can be extended to α -abelian rings for this general settings.

The ring R is called *abelian* if every idempotent is central, that is, $ae = ea$ for any $e^2 = e$, $a \in R$.

Definition 2.1 A ring R is called α -abelian if, for any $a, b \in R$ and any idempotent $e \in R$,

- (i) $ea = ae$,
- (ii) $ab = 0$ if and only if $a\alpha(b) = 0$.

So a ring R is *abelian* if and only if it is **1**-abelian.

Example 2.2 Let \mathbb{Z}_4 be the ring of integers modulo 4. Consider the ring $R = \left\{ \begin{pmatrix} a & b \\ 0 & a \end{pmatrix} \mid a, b \in \mathbb{Z}_4 \right\}$ with the usual matrix operations. Let $\alpha : R \rightarrow R$ be defined by $\alpha\left(\begin{pmatrix} a & b \\ 0 & a \end{pmatrix}\right) = \begin{pmatrix} a & -b \\ 0 & a \end{pmatrix}$. It is easy to check that α is a homomorphism of R . We show that R is an α -abelian ring. Since R is commutative, R is abelian. To complete the proof we check that for any $r, s \in R$, $rs = 0$ if and only if $r\alpha(s) = 0$. We prove one way implication. The other way is similar. So let $r = \begin{pmatrix} a & b \\ 0 & a \end{pmatrix}$, $s = \begin{pmatrix} x & y \\ 0 & x \end{pmatrix} \in R$. Assume that $rs = 0$ and

r and s are nonzero. Then we have $ax = 0$ and $ay + bx = 0$. If $a = 0$, then easy calculation shows that $r\alpha(s) = 0$. So we suppose $a \neq 0$. If $x = 0$ then $r\alpha(s) = 0$. Assume $x \neq 0$. Then $a = 2$ and $x = 2$. It implies $r\alpha(s) = 0$. Therefore R is α -abelian.

Lemma 2.3 *Let R be a ring such that for any $a, b \in R$, $ab = 0$ implies $a\alpha(b) = 0$, then $\alpha(e) = e$ for every idempotent $e \in R$.*

Proof. Since $e(1-e) = 0$ and $\alpha(1) = 1$, then $0 = e\alpha(1-e) = e - e\alpha(e)$. So $e = e\alpha(e)$. Further, $(1-e)e = 0$. Then $(1-e)\alpha(e) = 0$. Therefore, $\alpha(e) = e\alpha(e)$. So, we have $e = e\alpha(e)$ and $\alpha(e) = e\alpha(e)$. Hence, $e = \alpha(e)$. \square

Example 2.4 shows that there exists an abelian ring, but it is not α -abelian.

Example 2.4 Let R be the ring $\mathbb{Z} \oplus \mathbb{Z}$ with the usual componentwise operations. It is clear that R is an abelian ring. Let $\alpha : R \rightarrow R$ be defined by $\alpha(a, b) = (b, a)$. Then $(1, 0)(0, 1) = 0$, but $(1, 0)\alpha(0, 1) \neq 0$. Hence R is not α -abelian.

The ring R is called *semicommutative* if $ab = 0$ implies $aRb = 0$, for any $a, b \in R$. A ring R is called α -*semicommutative* if $ab = 0$ implies $aR\alpha(b) = 0$, for any $a, b \in R$. Agayev and Harmanci studied basic properties of α -semicommutative rings and focused on the semicommutativity of subrings of matrix rings (see [1]). In this note, the ring R is said to be α -*semicommutative* if, for any $a, b \in R$,

- (i) $ab = 0$ implies $aRb = 0$,
- (ii) $ab = 0$ if and only if $a\alpha(b) = 0$.

It is clear that a ring R is semicommutative if and only if it is **1**-semicommutative. The first part of Lemma 2.5 is proved in [7]. We give the proof for the sake of completeness.

Lemma 2.5 *If the ring R is α -semicommutative, then R is α -abelian. The converse holds if R is a right p.p.-ring.*

Proof. If e is an idempotent in R , then $e(1-e) = 0$. Since R is α -semicommutative, we have $ea(1-e) = 0$ for any $a \in R$ and so $ea = eae$. On the other hand, $(1-e)e = 0$ implies that $(1-e)ae = 0$, so we have $ae = eae$. Therefore, $ae = ea$. Suppose now R is an α -abelian and right p.p.-ring. Let $a, b \in R$ with $ab = 0$. Then $a \in r(b) = eR$ for some $e^2 = e \in R$ and so $be = 0$ and $a = ea$. Since R is α -abelian, we have $arb = earb = arbe = 0$ for any $r \in R$, that is, $aRb = 0$. Therefore R is α -semicommutative. \square

Corollary 2.6 *If the ring R is semicommutative, then R is abelian. The converse holds if R is a right p.p.-ring.*

Corollary 2.7 *Let R be an α -abelian and right p.p.-ring. Then $r(a) = r(aR)$, for any $a \in R$.*

Corollary 2.8 *Let R be an α -abelian and right p.p.-ring. Then R is a right p.q.-Baer ring.*

Proof. It follows from Corollary 2.7. \square

For a right R -module M , consider $M[x, \alpha] = \{\sum_{i=0}^s m_i x^i : s \geq 0, m_i \in M\}$. $M[x, \alpha]$ is an abelian group under an obvious addition operation and becomes a right module over $R[x; \alpha]$ under the following scalar product operation:

$$\text{For } m(x) = \sum_{i=0}^s m_i x^i \in M[x, \alpha] \text{ and } f(x) = \sum_{i=0}^t a_i x^i \in R[x, \alpha]$$

$$m(x)f(x) = \sum_{k=0}^{s+t} \left(\sum_{i+j=k} m_i \alpha^i(a_j) \right) x^k.$$

In [12], the ring R is called *Armendariz* if for any $f(x) = \sum_{i=0}^n a_i x^i, g(x) = \sum_{j=0}^s b_j x^j \in R[x]$, $f(x)g(x) = 0$ implies $a_i b_j = 0$ for all i and j . This definition of Armendariz ring is extended to modules in [11]. A module M is called α -*Armendariz* if the following conditions (1) and (2) are satisfied, and the module M is called α -*Armendariz of power series type* if the following conditions (1) and (3) are satisfied:

- (1) For $m \in M$ and $a \in R$, $ma = 0$ if and only if $m\alpha(a) = 0$.
- (2) For any $m(x) = \sum_{i=0}^n m_i x^i \in M[x, \alpha]$, $f(x) = \sum_{j=0}^s a_j x^j \in R[x, \alpha]$, $m(x)f(x) = 0$ implies $m_i \alpha^i(a_j) = 0$ for all i and j .
- (3) For any $m(x) = \sum_{i=0}^\infty m_i x^i \in M[[x, \alpha]]$, $f(x) = \sum_{j=0}^\infty a_j x^j \in R[[x, \alpha]]$, $m(x)f(x) = 0$ implies $m_i \alpha^i(a_j) = 0$ for all i and j .

In this note, the ring R is called α -*Armendariz* (α -*Armendariz of power series type*) if R_R is α -*Armendariz* (α -*Armendariz of power series type*) module. Hence R is an Armendariz (Armendariz of power series type) ring if and only if R_R is an **1**-Armendariz (**1**-Armendariz of power series type) module.

Theorem 2.9 *If the ring R is α -Armendariz, then R is α -abelian. The converse holds if R is a right p.p.-ring.*

Proof. Let $f_1(x) = e - ea(1 - e)x$, $f_2(x) = (1 - e) - (1 - e)aux$, $g_1(x) = 1 - e + ea(1 - e)x$, $g_2(x) = e + (1 - e)aux \in R[x, \alpha]$, where e is an idempotent in R and $a \in R$. Then $f_1(x)g_1(x) = 0$ and $f_2(x)g_2(x) = 0$. Since R is α -Armendariz, we have $ea(1 - e)\alpha(1 - e) = 0$. By Lemma 2.3, $\alpha(1 - e) = 1 - e$ and so $ea(1 - e) = 0$. Similarly, $f_2(x)g_2(x) = 0$ implies that $(1 - e)ae = 0$. Then $ae = eae = ea$, so R is α -abelian.

Suppose now R is an α -abelian and right p.p.-ring. Then R is abelian, and so every idempotent is central. By Lemma 2.3, $\alpha(e) = e$ for every idempotent $e \in R$. From Lemma 2.5, R is α -semicommutative, i.e., $ab = 0$ implies $aRb = 0$ for any $a, b \in R$. Let $f(x) = \sum_{i=0}^s a_i x^i$, $g(x) = \sum_{j=0}^t b_j x^j \in R[x, \alpha]$. Assume $f(x)g(x) = 0$. Then we have:

$$a_0 b_0 = 0 \tag{1}$$

$$a_0 b_1 + a_1 \alpha(b_0) = 0 \tag{2}$$

$$a_0 b_2 + a_1 \alpha(b_1) + a_2 \alpha^2(b_0) = 0 \tag{3}$$

...

By hypothesis there exist idempotents $e_i \in R$ such that $r(a_i) = e_i R$ for all i . So $b_0 = e_0 b_0$ and $a_0 e_0 = 0$. Multiply (2) from the right by e_0 , we have $0 = a_0 b_1 e_0 + a_1 \alpha(b_0) e_0 = a_0 e_0 b_1 + a_1 \alpha(b_0) \alpha(e_0) = a_1 \alpha(b_0)$. By (2) $a_0 b_1 = 0$ and so $b_1 = e_0 b_1$. Again, multiply (3) from the right by e_0 , we have $0 = a_0 b_2 e_0 + a_1 \alpha(b_1) e_0 + a_2 \alpha^2(b_0) e_0 = a_1 \alpha(b_1) + a_2 \alpha^2(b_0)$. Multiply this equation from right by e_1 , we have $0 = a_1 \alpha(b_1) e_1 + a_2 \alpha^2(b_0) e_1 = a_2 \alpha^2(b_0)$. Continuing in this way, we may conclude that $a_i \alpha^i(b_j) = 0$ for all $1 \leq i \leq s$ and $1 \leq j \leq t$. Hence R is α -Armendariz. This completes the proof. \square

Corollary 2.10 *If the ring R is Armendariz, then R is abelian. The converse holds if R is a right p.p.-ring.*

Proposition 2.11 *If the ring R is α -Armendariz of power series type, then R is α -abelian. The converse holds if R is a right p.p.-ring.*

Proof. Similar to the proof of Theorem 2.9. \square

Recall that a ring is *reduced* if it has no nonzero nilpotent elements. In [11], Lee and Zhou introduced α -reduced module. A module M is called α -reduced if, for any $m \in M$ and any $a \in R$,

- (1) $ma = 0$ implies $mR \cap Ma = 0$
- (2) $ma = 0$ if and only if $m\alpha(a) = 0$.

In this work, we call the ring R α -reduced if R_R is an α -reduced module. Hence R is a reduced ring if and only if R_R is an $\mathbf{1}$ -reduced module.

In [5], Hong et al. studied α -rigid rings. For an endomorphism α of a ring R , R is called α -rigid if $\alpha\alpha(a) = 0$ implies $a = 0$ for any a in R . The relationship between α -rigid rings and α -skew Armendariz rings was studied in [6]. In fact, R is an α -Armendariz ring if and only if (1) R is an α -skew Armendariz ring and (2) $ab = 0$ if and only if $a\alpha(b) = 0$ for any a, b in R . Note that α -reduced ring is α -rigid. Really, let R be an α -reduced ring and $\alpha\alpha(a) = 0$ for some a in R . Then $a^2 = 0$. Since R is reduced, we have $a = 0$. Further, by [5, Proposition 6], any α -reduced ring R is α -Armendariz. By Theorem 2.9, R is α -abelian. So, the first statement of Lemma 2.12 is a direct corollary of [5, Proposition 6].

Lemma 2.12 *If R is an α -reduced ring, then R is α -abelian. The converse holds if R is a right p.p.-ring.*

Proof. Let R be an α -abelian and right p.p.-ring. Suppose $ab = 0$ for $a, b \in R$. If $x \in aR \cap Rb$, then there exist $r_1, r_2 \in R$ such that $x = ar_1 = r_2 b$. Since R is right p.p.-ring, $ab = 0$ implies that $b \in r(a) = eR$ for some idempotent $e^2 = e \in R$. Then $b = eb$ and $xe = ar_1 e = r_2 be$. Since R is α -abelian and $ae = 0$, we have $ar_1 e = aer_1 = r_2 be = r_2 eb = r_2 b = 0$. Hence $aR \cap Rb = 0$, that is, R is α -reduced. \square

Corollary 2.13 *If R is a reduced ring, then R is abelian. The converse holds if R is a right p.p.-ring.*

According to Lambek [10], a ring R is called *symmetric* if whenever $a, b, c \in R$ satisfy $abc = 0$, we have $bac = 0$; it is easily seen that this is a left-right symmetric concept. We now introduce α -symmetric rings as a generalization of symmetric rings.

Definition 2.14 *The ring R is called α -symmetric if, for any $a, b, c \in R$,*

- (i) $abc = 0$ implies $acb = 0$,
- (ii) $ab = 0$ if and only if $a\alpha(b) = 0$.

It is clear that a ring R is symmetric if and only if it is **1**-symmetric.

Theorem 2.15 *Let R be a right p.p.-ring. Then the following are equivalent:*

- (1) R is α -reduced.
- (2) R is α -symmetric.
- (3) R is α -semicommutative.
- (4) R is α -Armendariz.
- (5) R is α -Armendariz of power series type.
- (6) R is α -abelian.

Proof. (1) \Leftrightarrow (6) From Lemma 2.12.

(4) \Leftrightarrow (6) Clear from Theorem 2.9.

(3) \Leftrightarrow (6) From Lemma 2.5.

(5) \Leftrightarrow (6) From Proposition 2.11.

(2) \Rightarrow (3) Let $a, b \in R$ with $ab = 0$. By hypothesis, $abc = 0$ implies $acb = 0$ for all $c \in R$. Hence $aRb = 0$ and so R is α -semicommutative.

(3) \Rightarrow (2) Assume that $abc = 0$, for any $a, b, c \in R$. Since R is right p.p.-ring, $c \in r(ab) = eR$ for some idempotent $e \in R$. Then $c = ec$ and $abe = 0$, so $acbe = 0$. We have already proved that semicommutativity implied being abelian, then $acbe = aecb$. Now $acb = aecb = acbe = 0$. It completes the proof. \square

Corollary 2.16 *Let R be a Baer ring. Then the following are equivalent:*

- (1) R is α -reduced.
- (2) R is α -symmetric.
- (3) R is α -semicommutative.
- (4) R is α -Armendariz.
- (5) R is α -Armendariz of power series type.
- (6) R is α -abelian.

One may suspect that if R is an abelian ring, then $R[x, \alpha]$ is abelian also. But this is not the case.

Example 2.17 Let F be any field, $R = \left\{ \begin{pmatrix} a & b & 0 & 0 \\ 0 & a & 0 & 0 \\ 0 & 0 & u & v \\ 0 & 0 & 0 & u \end{pmatrix} \mid a, b, u, v \in F \right\}$ and $\alpha : R \rightarrow R$ be defined by

$$\alpha \left(\begin{pmatrix} a & b & 0 & 0 \\ 0 & a & 0 & 0 \\ 0 & 0 & u & v \\ 0 & 0 & 0 & u \end{pmatrix} \right) = \begin{pmatrix} u & v & 0 & 0 \\ 0 & u & 0 & 0 \\ 0 & 0 & a & b \\ 0 & 0 & 0 & a \end{pmatrix}, \text{ where } \begin{pmatrix} a & b & 0 & 0 \\ 0 & a & 0 & 0 \\ 0 & 0 & u & v \\ 0 & 0 & 0 & u \end{pmatrix} \in R$$

Since R is commutative, R is abelian. We claim that $R[x, \alpha]$ is not an abelian ring. Let e_{ij} denote the 4×4 matrix units having alone 1 as its (i, j) -entry and all other entries 0. Consider $e = e_{11} + e_{22}$ and $f = e_{33} + e_{44} \in R$ and $e(x) = e + fx \in R[x, \alpha]$. Then $e(x)^2 = e(x)$, $ef = fe = 0$, $e^2 = e$, $f^2 = f$, $\alpha(e) = f$, $\alpha(f) = e$. An easy calculation reveals that $e(x)e_{12} = e_{12} + e_{34}x$, but $e_{12}e(x) = e_{12}$. Hence $R[x, \alpha]$ is not an abelian ring.

Lemma 2.18 *If R is an α -abelian ring, then the idempotents of $R[x, \alpha]$ belong to R , therefore $R[x, \alpha]$ is an abelian ring.*

Proof. Let R be α -abelian and $e(x) = \sum_{i=0}^t e_i x^i$ be an idempotent in $R[x, \alpha]$. Since $e(x)^2 = e(x)$, we have

$$e_0^2 = e_0 \quad (1)$$

$$e_0 e_1 + e_1 \alpha(e_0) = e_1 \quad (2)$$

$$e_0 e_2 + e_1 \alpha(e_1) + e_2 \alpha^2(e_0) = e_2 \quad (3)$$

...

Since R is α -abelian, R is abelian, and so every idempotent is central. By Lemma 2.3, $\alpha(e) = e$ for every idempotent $e \in R$. Then (2) becomes $e_0 e_1 + e_1 e_0 = e_1$ and so $e_1 = 0$. Since e_0 is central idempotent, (3) becomes $e_0 e_2 + e_2 e_0 = e_2$ and so $e_2 = 0$. Similarly, it can be shown that $e_i = 0$ for $i = 1, 2, \dots, t$. This completes the proof. \square

Lemma 2.19 *If $R[x, \alpha]$ is an abelian ring, then $\alpha(e) = e$ for every idempotent $e \in R$.*

Proof. Since $R[x, \alpha]$ is abelian, we have $f(x)e(x) = e(x)f(x)$ for any $f(x), e(x)^2 = e(x) \in R[x, \alpha]$. In particular, $xe = ex$ for every idempotent $e \in R$. Hence $xe = ex = \alpha(e)x$ and so $\alpha(e) = e$. \square

Lemma 2.20 *If $R[x, \alpha]$ is an abelian ring, then the idempotents of $R[x, \alpha]$ belong to R .*

Proof. Similar to the proof of Lemma 2.18. \square

Theorem 2.21 *If R is an α -abelian ring, then $R[x, \alpha]$ is abelian. The converse holds if $R[x, \alpha]$ is a right p.p.-ring.*

Proof. If R is α -abelian, by Lemma 2.18, $R[x, \alpha]$ is abelian. Suppose that $R[x, \alpha]$ be an abelian and right p.p.-ring. It is clear that $ae = ea$ for any $a, e^2 = e \in R$. Suppose $ab = 0$ for any $a, b \in R$. Since R is right p.p.-ring, we have $b \in r(a) = eR$, $b = eb$. So $a\alpha(b) = a\alpha(eb) = ae\alpha(b) = 0$. Conversely, let $a\alpha(b) = 0$. Then $axb = 0$. Since $R[x, \alpha]$ is right p.p.-ring, we have $b \in r_{R[x, \alpha]}(ax) = eR[x, \alpha]$ for some idempotent $e \in R[x, \alpha]$. So $b = eb$, $axe = 0$. By Lemma 2.20, $e \in R$. Hence $ae = 0$ and $ab = aeb = 0$. Therefore R is α -abelian. \square

Lemma 2.22 *Let R be an α -abelian ring. If for any countable subset X of R , $r(X) = eR$, where $e^2 = e \in R$, then*

(1) $R[[x, \alpha]]$ is a right p.p.-ring.

(2) If α is an automorphism of R , then $R[[x, x^{-1}, \alpha]]$ is a right p.p.-ring.

Proof. Let $a \in R$. Since $\{a\}$ is countable subset of R , $r(a) = eR$, i.e., R is a right p.p.-ring. Then from Theorem 2.15, R is α -Armendariz of power series type. By [11, Theorem 2.11.(1)(c), Theorem 2.11.(2)(c)], $R[[x, \alpha]]$ and $R[[x, x^{-1}, \alpha]]$ are right p.p.-rings. \square

Theorem 2.23 *Let R be an α -abelian ring. Then we have:*

(1) R is a right p.p.-ring if and only if $R[x, \alpha]$ is a right p.p.-ring.

(2) R is a Baer ring if and only if $R[x, \alpha]$ is a Baer ring.

(3) R is a right p.q.-Baer ring if and only if $R[x, \alpha]$ is a right p.q.-Baer ring.

(4) R is a Baer ring if and only if $R[[x, \alpha]]$ is a Baer ring.

Let $\alpha \in \text{Aut}(R)$.

(5) R is a Baer ring if and only if $R[x, x^{-1}, \alpha]$ is a Baer ring.

(6) R is a right p.p.-ring if and only if $R[x, x^{-1}, \alpha]$ is a right p.p.-ring.

(7) R is a Baer ring if and only if $R[[x, x^{-1}, \alpha]]$ is a Baer ring.

Proof. (1) “ \Rightarrow ”: Let $f(x) = a_0 + a_1x + \dots + a_tx^t \in R[x, \alpha]$. We claim that $r_{R[x, \alpha]}(f(x)) = eR[x, \alpha]$, where $e = e_0e_1\dots e_t$, $e_i^2 = e_i$ and $r_R(a_i) = e_iR$, $i = 0, 1, \dots, t$. By hypothesis and Lemma 2.3, $f(x)e = a_0e_0e_1\dots e_t + a_1e_1e_0e_2\dots e_tx + \dots + a_te_te_0e_1\dots e_{t-1}x^t = 0$. Then $eR[x] \subseteq r_{R[x, \alpha]}(f(x))$. Let $g(x) = b_0 + b_1x + \dots + b_nx^n \in r_{R[x, \alpha]}(f(x))$. Then $f(x)g(x) = 0$. Since R is an abelian and right p.p.-ring, by Theorem 2.9, R is Armendariz. So $a_ib_j = 0$ and this implies $b_j \in r_R(a_i) = e_iR$, and then $b_j = e_ib_j$ for any i . Therefore $g(x) = eg(x) \in eR[x, \alpha]$. This completes the proof of (1) “ \Rightarrow ”.

” \Leftarrow ”: Let $a \in R$. Then there exists $e(x)^2 = e(x) \in R[x, \alpha]$ such that $r_{R[x, \alpha]}(a) = e(x)R[x, \alpha]$. Then the constant term, e_0 say, of $e(x)$ is non-zero, and e_0 is an idempotent in R . So $e_0R \subset r_R(a)$. Now let $b \in r_R(a)$. Since $r_R(a) \subset r_{R[x, \alpha]}(a)$, $ab = 0$ implies that $b = e(x)b$ and so $b = e_0b$. Hence $r_R(a) \subset e_0R$, that is, $r_R(a) = e_0R$. Therefore R is a right p.p.-ring.

(2) ” \Rightarrow ”: Since R is Baer, R is a right p.p.-ring. By Lemma 2.5, R is Armendariz. Then from [11, Theorem 2.5.1(a)], $R[x, \alpha]$ is Baer.

” \Leftarrow ”: Let $R[x, \alpha]$ be a Baer ring and X be a subset of R . There exists $e(x)^2 = e(x) = e_0 + e_1x + \dots + e_nx^n \in R[x, \alpha]$ such that $r_{R[x, \alpha]}(X) = e(x)R[x, \alpha]$. We claim that $r_R(X) = e_0R$. If $a \in r_R(X)$, then $a = e(x)a$ and so $a = e_0a$. Hence $r_R(X) \subset e_0R$. Since $Xe(x) = 0$, we have $Xe_0 = 0$, that is, $e_0R \subset r_R(X)$. Then R is a Baer ring.

(3) ” \Rightarrow ”: Let $f(x) = a_0 + a_1x + \dots + a_tx^t \in R[x, \alpha]$. We prove $r_{R[x, \alpha]}(f(x)R[x, \alpha]) = e(x)R[x, \alpha]$, where $e(x) = e_0e_1\dots e_t$, $r_R(a_iR) = e_iR$. Since R is abelian, for any $h(x) \in R[x, \alpha]$ $f(x)h(x)e(x) = 0$. Then $e(x)R[x, \alpha] \subset r_{R[x, \alpha]}(f(x)R[x, \alpha])$. Let $g(x) = b_0 + b_1x + \dots + b_nx^n \in r_{R[x, \alpha]}(f(x)R[x, \alpha])$. Then $f(x)R[x, \alpha]g(x) = 0$ and so, $f(x)Rg(x) = 0$. From last equality we have $a_0Rb_0 = 0$. Hence $b_0 \in r_R(a_0R) = e_0R$. It follows that $b_0 = e_0b_0$. Also for any $r \in R$, the coefficient of x is equal to $a_0rb_1 + a_1\alpha(rb_0)$.

Hence $a_0rb_1 + a_1\alpha(rb_0) = 0$. Multiplying the equation $a_0rb_1 + a_1\alpha(rb_0) = 0$ from the right by e_0 , we have $a_1\alpha(rb_0e_0) = 0$, that is, $a_1\alpha(rb_0) = 0$. Since R is α -abelian, $a_1rb_0 = 0$. This implies $a_1Rb_0 = 0$. Then $b_0 \in r_R(a_1R) = e_1R$ and $b_1 \in r_R(a_0R) = e_0R$. So, $b_0 = e_1b_0$ and $b_1 = e_0b_1$. Again for any $r \in R$, $a_0rb_2 + a_1rb_1 + a_2rb_0 = 0$. Multiplying this equality from right by e_0e_1 and using previous results, we have $a_2rb_0 = 0$. Then $b_0 \in r_R(a_2R) = e_2R$. So $b_0 = e_2b_0$. Continuing this process we have $b_i = e_jb_i$ for any i, j . This implies $g(x) = e_0e_1\dots e_tg(x)$. So, $R[x, \alpha]$ is a right p.q.-Baer ring.

" \Leftarrow ": Let $a \in R$. Then $r_{R[x, \alpha]}(aR[x, \alpha]) = e(x)R[x, \alpha]$, where $e(x)^2 = e(x) \in R[x, \alpha]$. By Lemma 2.18, $e(x) = e_0 \in R$. Since $aR[x, \alpha]e(x) = 0$, $aR[x, \alpha]e_0 = 0$ and $aRe_0 = 0$. So, $e_0R \subset r_R(aR)$. Let $r \in r_R(aR) = r_R(aR[x, \alpha]) \subset r_{R[x, \alpha]}(aR[x, \alpha]) = e(x)R[x, \alpha]$. Then $e(x)r = r$. This implies $e_0r = r$ and so $r \in e_0R$. Therefore $r_R(aR[x, \alpha]) = e_0R$, i.e., R is a right p.q.-Baer ring.

(4) By Corollary 2.16, every abelian and Baer ring is Armendariz of power series type, so the proof follows from [11, Theorem 2.5 (1)(b)].

(5) By Corollary 2.16, R is α -Armendariz, then proof follows from [11, Theorem 2.5 (2)(a)].

(6) Since every α -abelian and right p.p.-ring is α -Armendariz by Theorem 2.9, the proof follows from [11, Theorem 2.11 (2)(a)].

(7) By Corollary 2.16, every abelian and Baer ring is Armendariz of power series type, it follows from [11, Theorem 2.5 (2)(b)]. □

Acknowledgment The authors express their gratitudes to the referee for (his/her) valuable suggestions.

References

- [1] Agayev, N., Harmanci, A.: On semicommutative modules and rings, Kyungpook Math. J., **47**, 21-30 (2007).
- [2] Birkenmeier, G.F., Kim, J.Y., Park, J.K.: On extensions of Baer and quasi-Baer Rings, J. Pure Appl. Algebra, **159**, 25-42 (2001).
- [3] Buhphang, A.M., Rege, M.B.: Semi-commutative modules and Armendariz modules, Arab. J. Math. Sci., **8**, 53-65 (2002).
- [4] Clark, E.W.: Twisted matrix units semigroup algebras, Duke Math. J., **34**, 417-424 (1967).
- [5] Hong, C.Y., Kim, N.K., Kwak, T.K.: Ore extensions of Baer and p.p.-rings, J. Pure and Appl. Algebra, **151(3)**, 215-226 (2000).
- [6] Hong, C.Y., Kim, N.K., Kwak, T.K.: On skew Armendariz rings, Comm. Algebra, **31(1)**, 103-122 (2003).
- [7] Huh, C., Lee, Y., Smoktunowicz, A.: Armendariz Rings and Semicommutative Rings, Comm. Algebra, **30(2)**, 751-761 (2002).
- [8] Kaplansky, I.: Rings of Operators, Math. Lecture Note Series, Benjamin, New York, 1965.
- [9] Kim, N.K., Lee, Y.: Armendariz rings and reduced rings, J. Algebra, **223**, 477-488 (2000).
- [10] Lambek, J.: On the representation of modules by sheaves of factor modules, Canad. Math. Bull., **14(3)**, 359-368 (1971).

- [11] Lee, T.K., Zhou, Y.: Reduced modules, Rings, modules, algebras, and abelian groups, 365-377, Lecture Notes in Pure and Appl. Math., 236, Dekker, New York, 2004.
- [12] Rege, M.B., Chhawchharia, S.: Armendariz Rings, Proc. Japan Acad. Ser. A Math.Sci., **73**, 14-17 (1997).

Nazim AGAYEV
Department of Computer Engineering,
European University of Lefke,
Gemikonagi-Lefke, Mersin 10, Cyprus
e-mail: nagayev@ eul.edu.tr

Received 05.11.2007

Abdullah HARMANCI
Mathematics Department
Hacettepe University
06550 Ankara, Turkey
e-mail: harmanci@hacettepe.edu.tr

Sait HALICIOĞLU
Department of Mathematics
Ankara University
06100 Ankara, Turkey
e-mail: halici@science.ankara.edu.tr