

1-1-2010

Some sufficient conditions for starlikeness and convexity

MAMORU NUNOKAWA

SHIGEYOSHI OWA

YAŞAR POLATOĞLU

MERT ÇAĞLAR

EMEL YAVUZ DUMAN

Follow this and additional works at: <https://dctubitak.researchcommons.org/math>



Part of the [Mathematics Commons](#)

Recommended Citation

NUNOKAWA, MAMORU; OWA, SHIGEYOSHI; POLATOĞLU, YAŞAR; ÇAĞLAR, MERT; and DUMAN, EMEL YAVUZ (2010) "Some sufficient conditions for starlikeness and convexity," *Turkish Journal of Mathematics*: Vol. 34: No. 3, Article 4. <https://doi.org/10.3906/mat-0807-42>
Available at: <https://dctubitak.researchcommons.org/math/vol34/iss3/4>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Mathematics by an authorized editor of TÜBİTAK Academic Journals.

Some sufficient conditions for starlikeness and convexity

*Mamoru Nunokawa, Shigeyoshi Owa, Yaşar Polatoğlu,
 Mert Çağlar and Emel Yavuz Duman*

Abstract

There are many results for sufficient conditions of functions $f(z)$ which are analytic in the open unit disc \mathbb{U} to be starlike and convex in \mathbb{U} . In view of the results due to S. Ozaki, I. Ono and T. Umezawa (1956), P.T. Mocanu (1988), and M. Nunokawa (1993), some sufficient conditions for starlikeness and convexity of $f(z)$ are discussed.

Key word and phrases: Univalent, starlike, convex.

1. Introduction and Preliminaries

Let \mathcal{A} be the class of functions $f(z)$ of the form

$$f(z) = z + \sum_{n=2}^{\infty} a_n z^n, \quad (1,1)$$

which are analytic in the open unit disc $\mathbb{U} = \{z \in \mathbb{C} \mid |z| < 1\}$. Let \mathcal{S} denote the subclass of \mathcal{A} consisting of functions $f(z)$ which are univalent in \mathbb{U} . Also, let \mathcal{S}^* and \mathcal{K} be the subclasses of \mathcal{S} consisting of all starlike functions $f(z)$ in \mathbb{U} and of all convex functions $f(z)$ in \mathbb{U} , respectively.

To discuss our problems, we have to recall here the following results.

Theorem A ([3]) *If $f(z) \in \mathcal{A}$ satisfies $|f''(z)| < 1$ ($z \in \mathbb{U}$), then $f(z) \in \mathcal{S}$.*

Theorem B ([1]) *If $f(z) \in \mathcal{A}$ satisfies*

$$|f'(z) - 1| < \frac{\sqrt{20}}{5} \quad (z \in \mathbb{U}), \quad (1.2)$$

then $f(z) \in \mathcal{S}^$.*

Theorem C ([1]) *If $f(z) \in \mathcal{A}$ satisfies*

$$|\arg f'(z)| < \frac{\pi}{2} \alpha_0 \quad (z \in \mathbb{U}), \quad (1.3)$$

then $f(z) \in \mathcal{S}^*$, where $\alpha_0 = 0.6165\dots$ is the unique root of the equation

$$2 \tan^{-1}(1 - \alpha) + (1 - 2\alpha)\pi = 0. \tag{1.4}$$

2. Starlikeness and convexity

We begin with the statement and the proof of the following result for sufficient condition of $f(z)$ to be in the class \mathcal{S}^* .

Theorem 1 *If $f(z) \in \mathcal{A}$ satisfies*

$$|f''(z)| \leq \frac{\sqrt{20}}{5} = 0.8944\dots \quad (z \in \mathbb{U}), \tag{2.1}$$

then $f(z) \in \mathcal{S}^*$.

Proof. Noting that

$$\begin{aligned} |f'(z) - 1| &= \left| \int_0^z f''(t) dt \right| \\ &\leq \int_0^{|z|} |f''(\varphi e^{i\theta})| d\varphi \\ &\leq \frac{\sqrt{20}}{5} \int_0^{|z|} d\varphi = \frac{\sqrt{20}}{5} |z| < \frac{\sqrt{20}}{5}, \end{aligned}$$

Theorem B gives us that $f(z) \in \mathcal{S}^*$. □

Remark 1 In view of the result by Obradović [4], we see that the sharp bound in (2.1) is 1.

Next we derive the following theorem.

Theorem 2 *If $f(z) \in \mathcal{A}$ satisfies*

$$|f''(z)| \leq \frac{\sqrt{5}}{5} = 0.4472\dots \quad (z \in \mathbb{U}), \tag{2.2}$$

then $f(z) \in \mathcal{K}$.

Proof. It follows that

$$\begin{aligned} |(zf'(z))' - 1| &= |f'(z) + zf''(z) - 1| \\ &\leq |f'(z) - 1| + |zf''(z)| \\ &\leq \left| \int_0^z f''(t) dt \right| + |zf''(z)| \\ &\leq \int_0^{|z|} |f''(t) dt| + \frac{\sqrt{5}}{5} |z| \\ &\leq \frac{2\sqrt{5}}{5} |z| < \frac{\sqrt{20}}{5}. \end{aligned}$$

Therefore, using Theorem B, we see that $zf'(z) \in \mathcal{S}^*$, or $f(z) \in \mathcal{K}$. □

Remark 2 By virtue of the result due to Mocanu [2], we know that the sharp bound in (2.2) is $\frac{1}{2}$.

Finally, applying the result due to Nunokawa [2], we can prove

Theorem 3 *If $f(z) \in \mathcal{A}$ satisfies*

$$|\arg(f'(z) + zf''(z))| < \frac{\pi}{2} \left(\alpha_1 + \frac{2}{\pi} \tan^{-1} \alpha_1 \right) \quad (z \in \mathbb{U}), \tag{2.3}$$

then $f(z) \in \mathcal{K}$, where $\alpha_1 = 0.3834\dots$ is the root of the equation

$$2\alpha_1 + \frac{2}{\pi} \tan^{-1} \alpha_1 = 1.$$

Proof. Note that

$$\arg(f'(z) + zf''(z)) = \arg f'(z) + \arg \left(1 + \frac{zf''(z)}{f'(z)} \right).$$

If there exists a point $z_0 \in \mathbb{U}$ such that

$$|\arg f'(z)| < \frac{\pi}{2} \alpha_1 \quad (|z| < |z_0|),$$

and

$$|\arg f'(z_0)| = \frac{\pi}{2} \alpha_1,$$

then Nunokawa's theorem in [2] gives us that

$$\frac{z_0 f''(z_0)}{f'(z_0)} = i\alpha_1 k,$$

where

$$k \geq \frac{1}{2} \left(a + \frac{1}{a} \right) \quad (\text{when } \arg f'(z_0) = \frac{\pi}{2} \alpha_1),$$

$$k \leq -\frac{1}{2} \left(a + \frac{1}{a} \right) \quad (\text{when } \arg f'(z_0) = -\frac{\pi}{2} \alpha_1),$$

and

$$f'(z_0)^{1/\alpha_1} = \pm ia \quad (a > 0).$$

Therefore, if $\arg f'(z_0) = \frac{\pi}{2} \alpha_1$, then we have

$$\begin{aligned} \arg f'(z_0) + \arg \left(1 + \frac{z_0 f''(z_0)}{f'(z_0)} \right) &= \frac{\pi}{2} \alpha_1 + \arg(1 + i\alpha_1 k) \\ &\geq \frac{\pi}{2} \alpha_1 + \tan^{-1} \alpha_1 \\ &= \frac{\pi}{2} \left(\alpha_1 + \frac{2}{\pi} \tan^{-1} \alpha_1 \right), \end{aligned}$$

which contradicts (2.3). If $\arg f'(z_0) = -\frac{\pi}{2}\alpha_1$, then, applying the same method for the previous case, we also have

$$\arg f'(z_0) + \arg \left(1 + \frac{z_0 f''(z_0)}{f'(z_0)} \right) \leq -\frac{\pi}{2} \left(\alpha_1 + \frac{2}{\pi} \tan^{-1} \alpha_1 \right),$$

which contradicts (2.3). Therefore, there exists no $z_0 \in \mathbb{U}$ such that $|\arg f'(z_0)| = \frac{\pi}{2}\alpha_1$. This implies that

$$|\arg f'(z)| < \frac{\pi}{2}\alpha_1 \quad (z \in \mathbb{U}).$$

Furthermore, since

$$\begin{aligned} \left| \arg \left(1 + \frac{z f''(z)}{f'(z)} \right) \right| - |\arg f'(z)| &\leq |\arg(f'(z) + z f''(z))| \\ &< \frac{\pi}{2} \left(\alpha_1 + \frac{2}{\pi} \tan^{-1} \alpha_1 \right) \quad (z \in \mathbb{U}), \end{aligned}$$

we conclude that

$$\left| \arg \left(1 + \frac{z f''(z)}{f'(z)} \right) \right| < \frac{\pi}{2} \left(2\alpha_1 + \frac{2}{\pi} \tan^{-1} \alpha_1 \right) = \frac{\pi}{2} \quad (z \in \mathbb{U}),$$

or

$$\operatorname{Re} \left(1 + \frac{z f''(z)}{f'(z)} \right) > 0 \quad (z \in \mathbb{U}).$$

This completes the proof of Theorem 3. □

3. Appendix

Let us consider a function $f(z) \in \mathcal{A}$ which satisfies

$$f'(z) = \frac{n}{n+z}, \quad f'(0) = 1,$$

where n is a sufficiently large positive integer. Then we see that, for $z_0 = \frac{\sqrt{2}}{2}(1+i)$,

$$\begin{aligned} \lim_{\substack{z \rightarrow z_0 \\ n \rightarrow n_0}} \arg \left(\frac{z f'(z)}{f(z)} \right) &= \lim_{\substack{z \rightarrow z_0 \\ n \rightarrow n_0}} \arg \left(\frac{z}{(n+z) \log(n+z)} \right) \\ &= \lim_{\substack{z \rightarrow z_0 \\ n \rightarrow n_0}} (\arg z - \arg(n+z) - \arg(\log(n+z))) \\ &= \frac{3}{4}\pi > \frac{\pi}{2}. \end{aligned}$$

This shows that, for arbitrary real α which is sufficiently close to 1 but less than 1, if $f(z) \in \mathcal{A}$ satisfies

$$\operatorname{Re} f'(z) > \alpha \quad (z \in \mathbb{U}),$$

then $f(z)$ is not necessarily starlike in \mathbb{U} .

References

- [1] Mocanu, P. T.: Some starlikeness conditions for analytic functions, *Rev. Roumaine Math. Pures. Appl.*, **33**, 117–124, (1988).
- [2] Mocanu, P. T.: Some simple criteria for starlikeness and convexity, *Libertas Mathematica*, **13**, 27–40, (1993).
- [3] Nunokawa, M.: On the order of strongly starlikeness of strongly convex functions, *Proc. Japan Acad., Sec.A* **68**, 234–237, (1993).
- [4] Obradović, M.: Simple sufficient conditions for univalence, *Matemat. Vesnik*, **49**, 242–244, (1997).
- [5] Ozaki, S., Ono, I. and Umezawa, T.: On a general second order derivative, *Sci. Rep. Tokyo Kyoiku Daigaku, Sec.A* **5**, 111–114, (1956).

Mamoru NUNOKAWA
Emeritus Professor
University of Gunma, Hoshikuki 798-8
Chuo, Chiba 260-0808, JAPAN

Received 23.07.2008

Shigeyoshi OWA
Department of Mathematics,
Kinki University, Higashi-Osaka,
Osaka 577-8502, JAPAN
e-mail: owa@math.kindai.ac.jp

Yaşar POLATOĞLU
Department of Mathematics and Computer Science,
T.C. İstanbul Kültür University,
34156 İstanbul-TURKEY
e-mail: y.polatoglu@iku.edu.tr

Mert ÇAĞLAR
Department of Mathematics and Computer Science,
TC İstanbul Kültür University,
34156 İstanbul-TURKEY
e-mail: m.caglar@iku.edu.tr

Emel YAVUZ DUMAN
Department of Mathematics and Computer Science,
T.C. İstanbul Kültür University,
34156 İstanbul-TURKEY
e-mail: e.yavuz@iku.edu.tr