

1-1-2003

## Effect of Cypermethrin and Permethrin on Cholinesterase Activity and Protein Contents in *Rana tigrina* (Amphibia)

MUHAMMAD ZAHEER KHAN

RAHILA TABASSUM

SYED NAIM-UL-HASAN NAQVI

ERUM Z. SHAH

FARHANA TABASSUM

*See next page for additional authors*

Follow this and additional works at: <https://journals.tubitak.gov.tr/zoology>

 Part of the [Zooology Commons](#)

---

### Recommended Citation

KHAN, MUHAMMAD ZAHEER; TABASSUM, RAHILA; NAQVI, SYED NAIM-UL-HASAN; SHAH, ERUM Z.; TABASSUM, FARHANA; AHMED, IMTIAZ; FATIMA, FARINA; and KHAN, MUHAMMAD FARHANULLAH (2003) "Effect of Cypermethrin and Permethrin on Cholinesterase Activity and Protein Contents in *Rana tigrina* (Amphibia)," *Turkish Journal of Zoology*. Vol. 27: No. 3, Article 11. Available at: <https://journals.tubitak.gov.tr/zoology/vol27/iss3/11>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Zoology by an authorized editor of TÜBİTAK Academic Journals. For more information, please contact [academic.publications@tubitak.gov.tr](mailto:academic.publications@tubitak.gov.tr).

---

## Effect of Cypermethrin and Permethrin on Cholinesterase Activity and Protein Contents in *Rana tigrina* (Amphibia)

### Authors

MUHAMMAD ZAHEER KHAN, RAHILA TABASSUM, SYED NAIM-UL-HASAN NAQVI, ERUM Z. SHAH, FARHANA TABASSUM, IMTIAZ AHMED, FARINA FATIMA, and MUHAMMAD FARHANULLAH KHAN

## Effect of Cypermethrin and Permethrin on Cholinesterase Activity and Protein Contents in *Rana tigrina* (Amphibia)

Muhammad Zaheer KHAN, Rahila TABASSUM  
Department of Zoology, University of Karachi, KARACHI

Syed Naeemul Hasan NAQVI  
Baqai Medical University, KARACHI

Erum Zehra SHAH, Farhana TABASSUM, Imtiaz AHMAD, Farina FATIMA  
Department of Zoology, University of Karachi, KARACHI

Muhammad Farhanullah KHAN  
Pakistan Agricultural Research Council, KARACHI

Received: 18.01.2002

**Abstract:** The effect of different concentrations (0.1% and 1%) of cypermethrin and permethrin were examined on the kidney and liver of *Rana tigrina*. After treatment with these pesticides, cholinesterase activity and protein contents were determined, in the kidney and liver. It was observed that protein contents and cholinesterase activities decreased in the treated animals.

**Key Words:** Induced effect, cypermethrin, permethrin, *Rana tigrina*, cholinesterase

### Cypermethrin ve Permethrin'in *Rana tigrina*'da (Amphibia) Kolinesterazın Etkinliğine ve Protein İçeriğine Etkisi

**Özet:** Cypermethrin ve permethrinin farklı derişimlerinin etkisi (% 0,1 ve % 1) *Rana tigrina*'nın böbrek ve karaciğeri üzerinde araştırıldı. Bu böcek ilaçlarının uygulanmasından sonra böbrek ve karaciğerdeki kolinesteraz etkinliği ve protein içeriği belirlendi. İşlemden geçirilen hayvanlarda protein içeriğinin ve kolinesteraz etkinliğinin azaldığı gözlemlendi.

**Anahtar Sözcükler:** Tetiklenen etki, cypermethrin, permethrin, *Rana tigrina*, kolinesteraz

### Introduction

Pesticides affect all members of an ecosystem, from the smallest invertebrates to birds and humans. Most toxic pesticides in both urban and agricultural settings are responsible for the deaths of many birds and fish and smaller aquatic animals that fish depend on for food. Amphibians are an important component of the food chain. They are cold-blooded animals and cannot regulate their body temperature. They live near water reservoirs or in water, so they are indirectly affected by pesticides used in agriculture for crop protection. In California many frogs and toads grow extra legs and eyes and do not survive to adulthood due to various effects of toxic chemicals.


No significant work has been done on the relationship between amphibians and pesticides, but some work on other vertebrates has been reported,

such as Meeks, (1968) and Fleet et al. (1972), who worked on pesticide residues in turtles and snakes. Hutson and Casida (1978) and Shono et al. (1979) isolated cypermethrin and deltamethrin in rats and mice. Harfenist et al. (1989) studied the effect of 11 pyrethroids on frogs. Mineau, (1993), O,Hara et al. (1991), Simon et al. (1999), Bishop et al. (2000), Parsons et al. (2000) and Khan, (2000) reported the effects of pesticides on different non-target organisms. In the present research the effects of cypermethrin and permethrin on the liver and kidney and cholinesterase activity and protein contents were studied. Some previous researchers have observed the effects of different pesticides on enzymatic activities of non-target species of vertebrates i.e. Custer et al. (1985), Hill, (1989), Cooper, (1991), Mineau, (1993) and Parsons et al. (2000).

## Materials and Methods

In the present research the subject organism was the adult frog *Rana tigrina*. The frogs were collected from Karachi University Campus and kept in the departmental pond. 1 ml cypermethrin and permethrin in concentrations of 0.1% and 1% (in acetone as solvent) each were injected subcutaneously in the abdominal region of the subject species with the help of an insulin syringe. After treatment with these chemicals the enzyme cholinesterase and protein contents were estimated in the kidney and liver of the frogs after 24 h post-treatment. The kidney and liver were taken as per Shakoori and Ahmad's (1973) techniques. Cholinesterase activity was estimated by a Randox Kit No. CE-190. The method is based upon the hydrolysis of acetylcholine by the action of cholinesterase, following Knedel and Boettger (1967). The reaction between thiocholine and dithiobis (nitrobenzoate) gives 2-nitro-5-mercaptobenzoate, a yellow compound which can be measured at 405 nm.

Principle:



In cholinesterase estimation three tubes were used for treated (i.e. liver and kidney) and one tube for untreated (i.e. also liver and kidney) assays. The first was solution 1 (i.e. buffer tube), the second was sample (kidney and liver) and third was solution 2 (i.e. substrate). A buffer solution of 1.5 ml was placed in the sample 1 tube, while a 0.01 ml sample and a 0.05 ml substrate were placed in the sample tube and the solution 2 tube, respectively. These were then mixed thoroughly, the initial absorbance read and the timer started simultaneously. This was read after 30, 60 and 90 s. The mean absorbance change per min was then determined.

Protein estimation was carried out by the Biuret method after Ahmad et al. (2000) based on Khan (1998).

## Results

The effects of different concentrations (0.1% and 1%) of cypermethrin and permethrin on the kidney and liver of *Rana tigrina* were examined. After treatment with these pesticides cholinesterase activity and protein contents were determined in the kidney and liver.

## Total protein contents

Total protein contents were estimated in the kidney and liver of *Rana tigrina* at 24 h post-treatment with the selected chemicals. A batch of untreated (control) sample was also kept for comparison purposes.

In cypermethrin-treated samples the reduction in protein content was 79% and 69% whereas in the permethrin-treated samples 61% and 16% decreases in the kidney were observed. In the liver the reduction was 75% and 44% in the cypermethrin-treated samples at concentrations of 0.1 and 1%, respectively.

## Cholinesterase activity

Cholinesterase activity was estimated in the kidney and liver of treated organisms. After 0.1% and 1% of cypermethrin and permethrin treatment, the inhibition was found to be 36% and 35% in the cypermethrin- and 35% and 6.76% in the permethrin-treated kidney, respectively. In the liver of cypermethrin-treated frogs the inhibition was 52% and 47%, whereas in permethrin, treated animals the inhibition was 29% and 23%, respectively.

## Discussion

Cholinesterase activity and the protein contents of kidney and liver decreased after cypermethrin and permethrin treatment (0.1% and 1%). Custer et al. (1985) described how after parathion spraying brain cholinesterase activity reduced up to 7%, 40%, 54% and 57% in blackbirds, pheasants, mice and coots respectively. Fairbrother et al. (1989) reported that 12 h after the second application of azinphos-methyl, plasma ChE levels in adult tree swallows were significantly inhibited, by 41%, as compared to the controls. ChE activity in another group of adult swallows dropped by 21% and 19% following the first and second sprayings.

Wilson et al. (1991) reported that hawks trapped near orchards recently treated with organophosphates diazinon and chlorpyrifos had lower levels of cholinesterase than the control group. Burgees et al. (1999) observed that organophosphate insecticides decreased cholinesterase activity in birds, and that these insecticides also produced tremors, loss of muscular control and paralysis. Flickinger et al. (1991) described how after parathion spraying geese collected from wheat fields exhibited up to 77% depression of their brain ChE activities 77%. Mineau (1993) reported that after

exposure to carbamate and organophosphate, cholinesterase activity in wild birds decreased. Taylor et al. (1999) observed the effects of sublethal doses (0.01 mg/g toad) and (0.0011 mg/g toad) of field grade malathion. The brain cholinesterase levels, decreased by 22% and 17% respectively. Parson et al. (2000) studied the effects of organophosphate and carbamate on non-target wildlife. These pesticides inhibited cholinesterase activity. In the present study cypermethrin and permethrin produced an inhibitory effect in the kidney and liver cholinesterase activity of the frog (*Rana tigrina*). The present findings are therefore in line with earlier results with reference to all the used compounds (organophosphate, organochlorine, carbamate and pyrethroids) applied directly or indirectly. In the present research cypermethrin and permethrin were applied to the frog body directly (0.1% and 1% conc.), and both compounds reduced cholinesterase activity in the kidney up to 36% and 35% after cypermethrin and 35% and 6.76% after permethrin treatment, whereas in the liver the percentage inhibition was 52% and 47% after cypermethrin treatment while in the case of permethrin 29% and 23% decreases were noted. These results appear to be in agreement with previous observations.

Total protein content also decreased in non-target vertebrate fauna after pesticide treatment, indicating pesticide-produced changes in the biochemical systems of non-target organisms. Proteins play an important role in the life of all living organisms. Pesticides disturb protein synthesis. In the present case the total protein contents of the liver and kidney in the frog (*Rana tigrina*) decreased after cypermethrin and permethrin (0.1% and 1% concentration at 1 ml dose) treatment. The reduction of protein content in the kidney was up to 79% and 69% after cypermethrin treatment and 61% and 16% after permethrin treatment. In the liver the reduction was 76% and 44% after cypermethrin treatment and 89% and 87% in permethrin-treated samples. Naqvi et al. (1986), Javid (1989), Nizam (1993), Saleem et al. (1998), Ahmad et al. (2000) and Tabassum and Naqvi (2001) also observed a reduction in total protein contents after pesticide application in different insects. Khan (2000) and Fatima (2001) also reported a reduction in the liver and kidney of reptiles. The present investigations also appear to be in line with the earlier findings. The present results therefore confirm the Khan, (2000) and Fatima, (2001) findings in this respect.

## References

- Ahmad, I. Perveen, A., Khan, M.F., Akhater, K. and Azmi, M.A. 2000. Determination of toxicity of early immature neem berries extract as compared to profenophos against *Tribolium castaneum*. Pakistan Zoological Congress (Ed., Shakoori, A.R), Vol. 20, pp. 93-100.
- Ahmad, I., Shamshad, A. and Tabassum, R. 2000. Effect of neem extract in comparison with cypermethrin (10 EC) and methylparathion (50 EC) on cholinesterase and total protein content of adult *Tribolium castaneum* (PARC strain). Bull. Pure & Appl. Sci., 19A(1): 55-61.
- Bishop, C.A., Collin, B., Mineau, P., Burgees, N.M., Red, W.F. and Risley, C. 2000. Reproduction activity of nesting birds in pesticide sprayed apple orchards in Southern Ontario, Canada. Environ. Toxicol. Chem., 19(3): 588-599.
- Burgees, N.M., Hunt, K.A., Bishop, C. and Weseloh, D.V. 1999. Cholinesterase inhibition in tree swallow (*Tachycineta* and eastern bluebird (*Sialia sialis*) exposed *bicolor*) to organophosphorus insecticides in apple orchards in Ontario, Canada. Environ Toxicol & Chem., 18(4): 708-716.
- Cooper, K. 1991. Effects of Pesticides on Wildlife, Handbook of Pesticides Toxicology. Academic Press, New York.
- Custer, T., Hill, E. and Ohlendorf, H. 1985. Effects on wildlife of ethyl and methyl parathion applied to California rice fields. California, Fish and Game, 71: 220-224.
- Fairbrother, A., Bennett, R.S. and Bennet, J.K. 1989. Sequential sampling of plasma cholinesterase in mallards (*Anas platyrhynchos*) as an indicator of exposure to cholinesterase inhibitors. Environ. Toxicol. Chem., 8: 117-122.
- Fatima, F. 2001. Bioecology of *Calotes versicolor* with special reference to induce effect of pyrethroid and organophosphate M.Phil thesis, University of Karachi, 147 pp.
- Fleet, R.R., Clark, D.R. Jr. and Plapp, F.W. Jr. 1972. Residues of DDT and dieldrin in snakes from two Texas agro systems. Biosci., 22: 664-665.
- Flickinger, E.L., Juenger, G., Roffe, T.J., Smith, M.R. and Irwin, R.J. 1991. Poisoning of Canada geese in Texas by parathion sprayed for control of Russian wheat aphid. J. Wildlife Dis., 27(2): 265-268.
- Harfenist, A., Power, T., Clark, K.L. and Peakall, D.B. 1989. A review and evaluation of the amphibian toxicological literature. Technical Report Series, No. 61, Canadian Wildlife Service, Ottawa.
- Hill, E.F. 1989. Sex and storage affect cholinesterase activity in blood plasma of Japanese quail. J. Wildlife Dis., 25(4): 580-585.
- Hutson, D.H. and Casida, J.E. 1978. Taurine conjugation in metabolism of 3-phenoxybenzoic acid and the pyrethroid insecticide cypermethrin in mouse. Xenobiotica, 8(9): 565-571.

- Javaid, M.A. 1989. Biochemical investigations on the effect of azadirachtin on the development of *Heliothis armigera* (Hub). M.Phil. Thesis, Quaid-e-Azam University, Islamabad, 270 pp.
- Khan, M.F. 1998. Determination of resistance in the cotton pest *Earias fabia stoll* against neem product in comparison with cypermethrin and monocrotophos. Ph.D. thesis, University of Karachi, 360 pp.
- Khan, M.Z. 2000. Determination of induced effect in *Agama* against permethrin and neem fractions and their effect on proteinic and enzymatic pattern. Ph.D. thesis, University of Karachi, 142 pp.
- Knedel, M. and Boettger, R. 1967. Kinetic method for determination of pseudocholinesterase (acylcholine acylhydrolase) activity. *Klin. Wochenschr.*, 45(6): 325-327.
- Meeks, R.L. 1968. The accumulation of 38 Cl ring-labeled DDT in a freshwater marsh. *J. Wild. Manag.*, 32: 576-598.
- Mineau, P. 1993. The hazard of carbofuran to birds and other vertebrate wildlife. Technical Report Series No. 177, Canada Wildlife Service.
- Naqvi, S.N.H., Shafi, S. and Zia, N. 1986. Effect of diflubenzuron and penfluron on the protein pattern of *Blattella germanica*. *Pak. J. Entomol. Kar.*, 1: 81-86.
- Nizam, S. 1993. Effect of allelochemicals against 3rd instar larvae of *Musca domestica* L. (Malir strain). Ph.D. thesis, University of Karachi, 369 pp.
- O'Hara, T.M., Krahn, M.M., Boyd, D., Becker, P.R. and Philo, L.M. 1991. Organochlorine contaminant levels in Eskimo harvested bowhead whales of arctic Alaska. *J. Wildlife Dis.*, 35(9): 741-752.
- Parson, K.C., Matz, A.C., Hooper, M.J. and Pokras, M.A. 2000. Monitoring wading bird exposure to agricultural chemicals using serum cholinesterase activity. *Environ. Toxicol. Chem.*, 19(5): 1317-1323.
- Saleem, M.A., Shakoori, A.R. and Mantte, D. 1998. In vivo ripcord induced macromolecular abnormalities in *Tribolium castaneum* larvae. *Pak. J. Zool.*, 30(3): 233-243.
- Shakoori, A.R. and Ahmad, M.S. 1973. Studies on the liver of chicken *Gallus domesticus*. Liver growth and nucleic acid content. *Pakistan J. Zool.* 5:111-117.
- Shono, T., Ohsawa, K. and Casida, J.E. 1979. Metabolism of trans-cypermethrin and cis permethrin trans-cypermethrin and cis-cypermethrin and decamethrin by microsomal enzymes. *J. Agric. Food Chem.*, 27(2): 316-325.
- Simon, L.M., Laszlo, K., Kotorman, M., Vertesi, A., Bagi, K. and Nemesok, J. 1999. Effects of synthetic and methidation on activities of some digestive enzymes in carp (*Cyprinus carpio* L.). *J. Environ. Sci. Health Part-B Pest Food Contam Agric. Wastes*, 34(5): 819-828.
- Tabassum, R. and Naqvi, S.N.H., Effect of dimilin (IGR), NC and Nfc (neem extracts) on nucleic acid and protein contents of *Callosobruchus analis* (F) of 21st Pakistan Zoological Congress (Ed., Shakoori, A.R) Inpress.
- Taylor, S.K., Williams, E.S. and Mills, K.W. 1999. Effects of malathion on disease susceptibility in wood house's toads. *J. Wildlife Dis.*, 35(3): 536-541.
- Wilson, B., Hooper, M., Littrell, E., Detrich, P., Mansen, M., Weisskopf, C. and Seiber, J. 1991. Orchard dormant sprays and exposure of red-tailed hawks to organophosphates. *Bull. Environ. Contam. Toxicol.*, 47: 717-724.