

1-1-1998

Spectrophotometric Determination of Chlorophyll - A, B and TotalCarotenoid Contents of Some Algae Species Using Different Solvents

Şükran DERE

Tohit GÜNEŞ

Rıdvan SIVACI

Follow this and additional works at: <https://journals.tubitak.gov.tr/botany>



Part of the [Botany Commons](#)

Recommended Citation

DERE, Şükran; GÜNEŞ, Tohit; and SIVACI, Rıdvan (1998) "Spectrophotometric Determination of Chlorophyll - A, B and TotalCarotenoid Contents of Some Algae Species Using Different Solvents," *Turkish Journal of Botany*. Vol. 22: No. 1, Article 3. Available at: <https://journals.tubitak.gov.tr/botany/vol22/iss1/3>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Botany by an authorized editor of TÜBİTAK Academic Journals. For more information, please contact academic.publications@tubitak.gov.tr.

Spectrophotometric Determination of Chlorophyll - A, B and Total Carotenoid Contents of Some Algae Species Using Different Solvents

Şükran DERE

Uludağ University, Faculty of Arts and Science, Department of Biology, Bursa-TURKEY

Tohit GÜNEŞ

Gazi University, Faculty of Education, Department of Biology, Kırşehir-TURKEY

Rıdvan SIVACI

Cumhuriyet University, Faculty of Arts and Science, Department of Biology, Sivas-TURKEY

Received: 14 / 6 / 1996

Accepted: 25 / 7 / 1997

Abstract: In our study, chlorophyll a, chlorophyll b and the total amounts of carotenoids in *Cladophora glomerata* (L.) Kuetzing, *Ulva rigida* L., *Codium tomentosum* (Huds.) Stackh and *Cladostephus verticillatus* Ag. species were determined. Methanol, acetone and diethyl ether were used as solvents. The chlorophyll amount in *Cladophora glomerata* L. Kuetzing, species and carotenoids in *Cladostephus verticillatus* Ag. were found the highest. It was observed that sonication had no much contribution to the extraction.

Key Words: Algae, Chlorophyll, Carotenoid, Solvents

Farklı Çözücüler Kullanılarak Bazı Alg Türlerinde Klorofil - A, B ve Toplam Karotenoid İçeriğinin Spektrofotometrik Saptanması

Özet: Çalışmamızda *Cladophora glomerata* (L.) Kuetzing., *Ulva rigida* L., *Codium tomentosum* (Huds.) Stackh ve *Cladostephus verticillatus* Ag. türlerinde klorofil a, klorofil b ve toplam karotenoid miktarları saptandı. Çözücü olarak metanol, aseton ve dietil eter kullanıldı. *Cladophora glomerata* L. Kuetzing, türünde klorofil a miktarı, *Cladostephus verticillatus* Ag. da ise karoten miktarı en yüksek bulunmuştur. Sonikasyonun ekstraksiyona fazla bir katkısının olmadığı gözlenmiştir.

Anahtar Sözcükler: Alg, Klorofil, Karotenoid, Çözücüler

Introduction

The spectrophotometric definition of photosynthetic pigments that cause light energy to turn into chemical energy in all photosynthetic organisms was first determined by Stokes in 1864 (1). Later, the examples obtained from *Fucus* L. and *Laminaria* L. were classified as blue chlorophyll (chlorophyll a), green chlorophyll (chlorophyll b), chlorofucin (chlorophyll c1, chlorophyll c2) and orange-yellow (xanthophyll) according to the pigment colors (2).

The absorbance properties of pigments facilitate the qualitative and quantitative analysis of them. In the definition of the pigment content of freshwater and sea algae various methods and solvents have been used (3,4) and it was determined that the pigment

level was influenced by limiting factors such as high-light, lack of nitrogen and limited nutrient (5,6).

In earlier studies, it was suggested that the content of chlorophyll a relating to the pigment level was almost the same in all algae groups, but chlorophyll b and c changed, and so did the carotene level depending on the algae species and environmental conditions, and especially there was an increase in the carotene level in stress conditions (7, 8, 9). Although the role of carotene pigments in algae was not exactly known, it has been suggested that they functioned as a passive light protecting filter and have got the role of accessory pigments transferring energy and oxygen (10, 11, 12). Vechetel and et al. (1992) in their studies determined that carotene pigments were the most

important photosynthetic pigments, and they prevented chlorophyll and thylakoid membrane from the damage of absorbed energy by photooxidation (13).

In our study, the pigment levels of four algae species were determined by using various solvents (methanol, acetone, diethyl ether) and the contribution of these solvents to the extraction in various species was examined comparatively. In addition, after homogenization it was examined whether sonication had any contribution to the extraction. Furthermore, the determination of suitable methods and solvents that could be used in studies on pigments was aimed.

Material and Method

Of the samples we used for the studies in the laboratory, freshwater form *Cladophora glomerata* (L.) Kützing (*Chlorophyta*) was collected from the Yeşilirmak (Tokat), sea forms *Codium tomentosum* (Huds.) Stackh (*Chlorophyta*), *Ulva rigita* L. (*Chlorophyta*) and *Cladostephus verticillatus* Ag. (*Phaeophyta*) were collected from the Aegean sea (Narlidere-Izmir). The samples were brought to the laboratory in the natural water in which they had lived, and one day later the extraction processes were carried out.

Extraction Processes

The weighed samples, having been put separately in 95% diethyl ether, 96% methanol and 100% acetone (50 ml for each gram), were homogenized with the B-Brawn type homogenizer at 1000 rpm for one minute. The homogenate was filtered through two layer cheese cloths, and was centrifuged using the Nüve Fjü 650 model centrifuge at 2500 rpm for ten minutes. The supernatant was separated and the absorbances were read at 400-700 nm on Shimadzu UV-260 spectrophotometer. It was recorded that Chlorophyll a showed the maximum absorbance at 662 nm, chlorophyll b at 646 nm and total caroten at 470 nm and the amount of these pigments was calculated

Table 1. The formulas that used in the calculation of chlorophyll a, chlorophyll b and total caroten levels

| | |
|---------------|--|
| Diethyl ether | $C_a = 10.05 A_{662} - 0.766 A_{644}$ $C_b = 16.37 A_{644} - 3.140 A_{662}$ $C_{x+c} = 1000 A_{470} - 1.280 C_a - 56.7 C_b / 230$ |
| Methanol | $C_a = 15.65 A_{666} - 7.340 A_{653}$ $C_b = 27.05 A_{653} - 11.21 A_{666}$ $C_{x+c} = 1000 A_{470} - 2.860 C_a - 129.2 C_b / 245$ |
| Acetone | $C_a = 11.75 A_{662} - 2.350 A_{645}$ $C_b = 18.61 A_{645} - 3.960 A_{662}$ $C_{x+c} = 1000 A_{470} - 2.270 C_a - 81.4 C_b / 227$ |

C_a = Chlorophyll a, C_b = Chlorophyll b, C_{x+c} = Total carotene

according to the formulas of Lichtentaler and Wellburn (1985) (14). The formulas were showed in Table 1.

Instead of the wave-lengths determined by these researchers, 662, 646 and 470 nm at which we observed maximum absorbance were used. The experiments were repeated three times. The results of the above experiments were analysed statistically using Snedecor's F-test for analysis of variance and to determine statistically significant differences between means "Multiple Range Test" was applied (15, 16).

Results

Chlorophyll a content in four algae species whose pigment levels were studied was determined the highest in *Cladophora glomerata* species (Table 2). Although chlorophyll a levels in the other three sea forms were very close to each other, it was found a little higher in *Ulva rigita* in comparison with the other two species (Table 2). In terms of chlorophyll b content no significant difference was observed between the species ($P < 0.01$), but no absorbance could be de-

Table 2. Pigment levels of the species in different solvents ($\mu\text{g/gfw}$)

| Species | Methanol | | | Diethyl ether | | | Acetone | | |
|-----------------------------------|------------|------------|------------|---------------|------------|------------|------------|------------|------------|
| | C_a | C_b | Carotene | C_a | C_b | Carotene | C_a | C_b | Carotene |
| <i>Cladophora glomerata</i> | 60.7±0.05x | 23.0±1.05x | 19.2±0.95x | 53.8±2.50x | 20.1±0.65x | 18.8±0.34x | 56.2±0.46x | 20.3±0.80x | 20.1±1.05x |
| <i>Ulva rigita</i> | 54.6±1.40y | 24.1±0.80x | 20.8±1.15x | 48.6±0.72y | 21.2±0.10x | 21.6±0.72x | 48.9±0.72y | 23.3±0.95x | 20.6±0.95x |
| <i>Codium tomentosum</i> | 49.0±2.60y | 21.8±0.30x | 24.7±0.45x | 46.2±0.05y | 20.1±2.50x | 20.5±0.96x | 47.4±0.45y | 20.2±1.05x | 22.4±0.60x |
| <i>Cladostephus verticillatus</i> | 51.3±1.00y | - | 33.8±1.30y | 47.9±0.65y | - | 28.7±0.25y | 48.2±0.20y | - | 29.5±0.65y |

* Data shown with the same symbols in the vertical column are not different from each other on 0.01 statistical levels

± Standard error

terminated because of the absence of chlorophyll b in *Cladostephus verticillatus* species. The total carotene content was determined the highest in *Cladostephus verticillatus* and the lowest in *Cladophora glomerata* (Table 2). It was observed that the pigments were not extracted completely from *Cladostephus verticillatus* species. The total carotene content was determined the highest in *Cladostephus verticillatus* species and *Codium tomentosum* species as a result of the extraction, and particularly these two species contained pigments in certain proportions after homogenization.

It was determined that the solvents used were important in the pigment extraction, and the best solvent was methanol (Table 4). It was observed that the extraction with methanol was nearly complete in *Cladophora glomerata* and *Ulva rigita* because of the diversity of the cell wall structures, (Table 2, 3). Although it was seen that acetone was a better solvent in comparison with diethyl ether, no significant difference was determined between two solvents.

It was observed that the 5 minute sonication carried out in addition to homogenization was not very efficient on the isolation of the pigments during the extraction process. It was determined that it increased the extraction only a little in *Cladostephus verticillatus*, *Codium tomentosum* (Table 3).

Discussion

Although it was suggested in earlier studies that the chlorophyll a level was the same in all algae groups (7, 8, 9), in the present study it has been found that the level of chlorophyll a in fresh water form *Cladophora glomerata* was rather high in comparison with the other three species. The chlorophyll a level in *Ulva rigita* was also found higher than the other two species. Because of the variety of the cell wall structures of those two species, a well performed homogenization and a complete extraction facilitated the isolation of the pigments. In addition we have got an opinion that there was no much change at the pig-

ment level in connection with the term when the samples were taken because *Cladophora glomerata* lives in freshwater and in an environment where much light and temperature stratifications are not seen (17). In previous studies it was determined that there were changes at the pigment level of algae species that live in an environment where light stratification is seen (5, 6). That is why we have got an opinion that the reason for the chlorophyll a level to be lower in other species is linked with light and temperature stratification during the term when the examples were taken.

Furthermore, because the accumulation of CaCO_3 was high on the cell walls of *Codium tomentosum* and *Cladostephus verticillatus* species that we used in our study (18), it was observed that homogenization was difficult and all of the pigments were not extracted. So we believe that the use of more suitable methods in the studies for breaking up cell walls will be useful for extraction to be complete.

It is known that carotene pigments are at various amount in algae groups. In the earlier studies it was proved that there were more carotene and xanthophyll pigments in *Phaeophyta* (19, 20). In the samples studied the carotene content in *Cladostephus verticillatus*, only species belonging to *Phaeophyta* group, was found higher in comparison with the other three species. The absence of chlorophyll b in this species may be due to more carotene pigments that have short wave-length and can absorb the light. It was pointed out that in the studies with *Chlorella* Beyerinck the carotene pigment level might be different even in the same species and it depends on various environmental conditions (21). According to these results we suggest that the comparison of the caroten levels of species used is not accurate because each one of the species examined in this study was collected from different mediums.

The solvents used in the pigment extraction have an important effect. In the studies with *Scenedesmus quadricauda* (Turp) de Bre'bisson in de Bre'bisson and

Table 3. Effect of sonication on extraction of pigments ($\mu\text{g/gfw}$)

| Species* | Methanol | | | Diethyl ether | | | Acetone | | |
|-----------------------------------|---------------|---------------|-----------|---------------|---------------|-----------|---------------|---------------|-----------|
| | Cl_a | Cl_b | Carotene | Cl_a | Cl_b | Carotene | Cl_a | Cl_b | Carotene |
| <i>Cladophora glomerata</i> | 61.4±1.05 | 23.6±0.72 | 19.4±1.05 | 53.7±1.65 | 20.8±1.67 | 19.1±0.05 | 57.0±2.67 | 20.4±1.45 | 19.7±2.45 |
| <i>Ulva rigita</i> | 54.7±1.40 | 24.1±0.30 | 21.1±0.65 | 49.0±0.25 | 21.3±0.95 | 21.5±0.65 | 49.3±1.95 | 24.2±0.30 | 21.4±0.95 |
| <i>Codium tomentosum</i> | 50.9±0.95 | 20.9±1.40 | 27.9±0.45 | 48.2±0.95 | 21.2±1.25 | 21.4±1.67 | 49.9±0.95 | 21.6±3.20 | 22.4±0.72 |
| <i>Cladostephus verticillatus</i> | 53.1±0.25 | - | 36.2±1.40 | 50.8±0.65 | - | 30.6±1.12 | 51.8±0.25 | - | 31.9±2.60 |

± Standard error

Table 4. Effect of solvents on pigment levels

| Species | Pigment | Methanol* | Diethyl Ether | Acetone |
|-----------------------------|-----------------|------------|---------------|------------|
| Chladophora glomerata | Cl _a | 60.7±0.05a | 53.8±2.50b | 56.2±0.46b |
| | Cl _b | 23.0±1.05a | 20.1±0.65b | 20.3±0.80b |
| | Carotene | 19.2±0.95a | 18.8±0.34a | 20.1±1.05a |
| Ulva rigita | Cl _a | 54.6±1.40a | 48.6±0.72b | 48.9±0.72b |
| | Cl _b | 24.1±0.80a | 21.2±0.10b | 23.3±0.95a |
| | Carotene | 20.8±1.15a | 21.6±0.72a | 20.6±0.95a |
| Codium tomentosum | Cl _a | 49.0±2.60a | 46.2±0.05b | 47.4±0.45b |
| | Cl _b | 21.8±0.30a | 20.1±2.50b | 20.2±1.05b |
| | Carotene | 24.7±0.45a | 20.5±0.96b | 22.4±0.60c |
| Chladostephus verticillatus | Cl _a | 51.3±1.00a | 47.9±0.65b | 48.2±0.20b |
| | Cl _b | - | - | - |
| | Carotene | 33.8±1.30a | 28.7±0.25b | 29.5±0.65b |

* Data shown with the same symbols in the horizontal column are not different from each other on 0.01 statistical levels
± Standard error

References

1. Stokes, G.G., On the Supposed Identity of Biliverdin with Chlorophyll with Remarks on the Constitution of Chlorophyll. Proc. Roy. Soc. 13, 144-145 (1864).
2. Sorby, H.C., On Comparative Vegetable Chromatology. Proc. Roy. Soc. 21: 442-483 (1873).
3. Nusch, E.A., Comparison of Methods for Chlorophyll and Phaeopigment Determination. Arch. Hydrobiol. Limnol. 14: 14-36 (1980).
4. Sartory, D.P., Extraction of Chlorophyll A from Freshwater Phytoplankton for Spectrophotometric Analysis. M. Sc. Thesis. Univ. Orange free state, Bloemfontein, Republic of S. Africa. (1982).
5. Greene, R.M., Gerder, R.J., Falkowski, P.G., Effect of Iron Limitation on Photosynthesis in a Marine Diatom. Limnol. Oceanogr. 36: 1772-1782 (1991).
6. Collier, J.L., Grossman, A.R., Chlorosis Induced by Nutrient Deprivation in *Synechococcus* sp. J. Bacteriol. 174: 4718-4726 (1992).
7. Donkin, P., Ketocarotenoid Biosynthesis by *Haematococcus lacustris*. Phytochemistry 15: 711-715 (1976).
8. Martin, J.H., Gordon, R.M. and Fitzwater, S.E., The Case for Iron Limnol. Oceanogr. 36: 1793-1803 (1991).
9. Grung, M.F., D'souza, M.L., Borowitzka, M., Algal Carotenoids 1. Secondary Carotenoids. J. Appl. Phycol. 4. 165-171 (1992).
10. Yong, Y.Y.R., Lee, Y.K., Carotenoids Play a Photoprotective Role in the Cytoplasm of *Haematococcus lacustris* (Chrolophyta). Phycologia 30. 257-261 (1991).
11. Lichtenthaler, H.K. Chlorophylls and Carotenoids, Pigments of Photosynthetic Biomembranes. Methods in Enzymology 148: 350-382 (1987).
12. Bidigare, R.R., Ondrusek, M.E., Kennicutt, M.C., Harvey, H.R., Evidence for a Photoprotective Function for Secondary Carotenoids of Snow Algae. J. Phycol. 29, 427-434 (1993).
13. Vechetel, B.W., Ruppel, H.G., Lipid Bodies in *Eremosphaera viridis* De Bary (Chlorophyceae). Plant and Cell Phys. 31, 41-48 (1992).
14. Lichtenthaler, H.K., Wellburn, A.R., Determination of Total Carotenoids and Chlorophylls A and B of Leaf in Different Solvents. Biol. Soc. Trans. 11. 591-592 (1985).
15. Snedecor, G.W., Cochran, W.G., Statistical Methods Iowa State Univ. Press Ames. (1980).
16. Duncan, D.B., Multiple Range and Multiple F Tests. Biometrics 11, 1-41 (1955).
17. Palmer, C.M., Algae and Water Pollution. Castle Home Pub. 37-42 (1980).
18. Fritsch, F.E., The Structure and Reproduction of the Algae. Vol. I 408-410 (1971).
19. Fritsch, F.E., The Structure and Reproduction of the Algae. Vol. II 261-269 (1971).
20. Tanaka, A., Tsuji, H., Changes in Chlorophyll A and B Content in Dark-Incubated Plants. Plant Phys. 68: 567-570 (1981).
21. Rise, M., Cohen, E., Vishkautsan, M., Arad, S., Accumulation of Secondary Carotenoids in *Chlorella zofingiensis*. J. Plant Phys. Vol. 144: 287-292 (1994).

Godey and *Selenastrum capricornutum* Reinsch ethanol, methanol and acetone were used, and it was shown that the best solvent was ethanol (22). Yet in our study it was determined that the best solvent used for algae species was methanol. We think that this difference depends on the kind of the plant and its cell wall structure. It was observed that methanol was a better solvent for these species because the cell wall structures of *Codium tomentosum* and *Cladostephus verticillatus* we used were in porous structure and there was too much CaCO₃ accumulation (18). It has been seen that acetone, chloroform and diethyl ether had less effect on homogenization.

In fact, acetone, chloroform, diethyl ether, dimethyl formamide and methanol were used in the studies with high plant leaves, and it was determined that the extraction rate was various in every solvent (23). That is why we believe that the selection of the method and the solvent to be used in the studies in connection with pigments according to the species will be more useful. Although the methanol was a good extractant for some algae species, it should not be forgotten that it was toxic.

22. Sartory, D.P., Grobbelaar, J.U., Extraction of Chlorophyll a from Freshwater Phytoplankton for Spectrophotometric Analysis. *Hydrobiologia* 114: 177-187 (1984).
23. Wellburn, A.R., The Spectral Determination of Chlorophylls A and B, as well as Total Carotenoids, Using Various Solvents with Spectrophotometers of Different Resolution. *J. Plant Phys.* Vol. 144, 307-313 (1994).