

1-1-1998

Two New Species and A New Hybrid from Anatolia

Hayri DUMAN

K. Hüsnü Can BAŞER

Zeki AYTAÇ

Follow this and additional works at: <https://journals.tubitak.gov.tr/botany>



Part of the [Botany Commons](#)

Recommended Citation

DUMAN, Hayri; BAŞER, K. Hüsnü Can; and AYTAÇ, Zeki (1998) "Two New Species and A New Hybrid from Anatolia," *Turkish Journal of Botany*. Vol. 22: No. 1, Article 8. Available at: <https://journals.tubitak.gov.tr/botany/vol22/iss1/8>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Botany by an authorized editor of TÜBİTAK Academic Journals. For more information, please contact academic.publications@tubitak.gov.tr.

Two New Species and A New Hybrid from Anatolia

Hayri DUMAN

Gazi University, Faculty of Science and Art, Department of Biology, Ankara-TURKEY

K. Hüsnü Can BAŞER

Anadolu University, Medicinal Plants research Centre (TBAM), Eskişehir-TURKEY

Zeki AYTAÇ

Gazi University, Faculty of Science and Art, Department of Biology, Ankara-TURKEY

Received: 13 / 12 / 1996

Accepted: 8 / 8 / 1997

Abstract: Two new species and a new hybrid of *Labiatae* from south Anatolia are described and illustrated: *Sideritis caesarea* Duman, Aytaç et Başer, *Sideritis vuralii* Duman et Başer, *Origanum x adanense* Başer et Duman.

Key Words: *Sideritis*, *Origanum*, *Labiatae*, New species, Turkey

Anadolu'dan İki Yeni Tür ve Bir Yeni Hibrid

Özet: Anadolu'dan iki yeni tür ve bir yeni hibrid (*Labiatae*) tanımlandı ve şekilleri verildi. Bunlar; *Sideritis caesarea* Duman, Aytaç et Başer, *Sideritis vuralii* Duman et Başer, *Origanum x adanense* Başer et Duman'dır.

Anahtar Sözcükler: *Sideritis*, *Origanum*, *Labiatae*, Yeni tür, Türkiye

Introduction

Since *Sideritis* L. was revised by A. Huber-Morath for Flora of Turkey (1), four new species have been described from Turkey, *S. huber-morathii* Greuter et Burdet, *S. athoa* Papanikolau et Kokkini (2), *S. akmanii* Z. Aytaç, M. Ekici et A. Dönmez, *S. gulendamii* H. Duman et F. A. Karavelioğulları (3) and one species (*S. scardica* Griseb. subsp. *scardica*) was added new record for Turkey (4). *S. caesarea* and *S. vuralii* are described as new species in this paper and this increases the number of *Sideritis* species known from Turkey to 45. After *Origanum* L. was revised by J. H. Letswaart for Flora of Turkey (1), *O. munzurense* Kit Tan et Sorger (2) and *O. husnucan-baserii* H. Duman, Z. Aytaç et A. Duran (3) have been described new to science from Turkey. 23 species are now known from Turkey. 17 hybrids of *Origanum* are known (5), five of which grow in Turkey (1). *O. x adanense* (*O. bargyli* Mouterde x *O. laevigatum* Boiss.) is described as a new hybrid in this paper.

Sideritis caesarea Duman, Aytaç et Başer spec. nov. (Sect. *Empedoclia* (Rafin.) Benth.) Fig. 1, Map 1.


Affinis *S. germanicopolitanae* Bornm. et *S. libanot-*

ica Labill. subsp. *kurdicae* (Bornm.) Hub. Mor. sed ab *S. germanicopolitanae foliis medianis caulinis* 3.5-8 x 0.3-1 cm (non 5-12 x 1-2 cm), bracteis medianis 1-1.5 x 0.7-1.3 cm (non 1-6 cm), calycibus 9-10 mm (non 9-13 mm), calycibus pilis longis simplicibus et brevibus glandulosis, corollis 10-12 mm (non 12-14 mm) distincta. Ab *S. libanotica* subsp. *kurdicae* foliis caulinis 2-4- jugis (non 4-8- jugis), bracteis pilis glanduliferis et eglandulosis pilosis (haud dense pilosis pilis eglandulosis tantum), calyce 9-10 mm, pilis glanduliferis et eglandulosis obtecto (non circa 6 mm et dense albopiloso), calycis dentibus paulo longiore 3-4 mm (non 2.5-3 mm) differt.

Perennial, 30-50 cm, erect, simple or branched from the base. Stems shortly white villous-tomentose to glabrescent below, glabrescent to with short eglandular and glandular hairs above. Lower leaves distinctly petiolate, petioles 1-3 cm, glabrescent to villous-tomentose; lamina oblanceolate, 3-5 x 0.5-1 cm, finely serrate to crenate or entire, glabrescent to villous-tomentose, reticulately nerved. Middle cauline leaves linear to oblanceolate, sessile or shortly petiolate, 3.5-8 x 0.3-1 cm, entire or obscurely crenate-serrate, acute, reticulately nerved. Internodes to 10 cm long.



Figure 1. *Sideritis caesarea*: A-habit, B-bract, C-calyx, D-flower, E-corolla



Map 1. Distribution of Δ *Sideritis caesarea*, \circ *Sideritis vuralii*, * *Origanum adanensis*

Inflorescence simple or 2-3-branched, 10-25 cm, glabrescent to shortly glandular and eglandular hairy; verticillasters 4-6 (-10), 6-flowered, 2-6 cm distant. Middle bracts orbicular to cordate, acuminate, 1-1.5 x 0.7-1.3 cm including 3-7 mm acumen, glandular or eglandular, margine ciliate, prominently reticulate nerved. Calyx 9-10 mm, purplish; tube c. 6 mm, long simple and short glandular hairy; teeth triangular-lanceolate, acute, 3-4 mm, with glandular and eglandular hairs. Corolla yellow, 10-12 mm, hairy outside and inside, without brown markings, limb 2-3 mm. Nutlets triangular-ovoid, rounded at apex, glabrous, minutely tuberculate. *Fl.* and *Fr.* July-September. *High mountain steppe*, 1500-2400 m.

Type: Turkey, B6 Kayseri: Sarız, Yalak (Yeşilkent), Binboğa Dağı, Tekkekayası mevkii, 1800-2200 m, yüksek dağ stebi, 4.viii.1991. Z.Aytaç 4310- H. Duman (holo- GAZI; iso- ESSE, ANK).

Central Anatolia. Endemic. Irano-Turanian element.

A5 Yozgat: Çekerek, Deveci Dağı, c. 1900 m, stony place, 14. vii.1979. *R. İlarıslan* -527 ANKI, B6 Kayseri: Hınzır Dağı, above Ganişeyh, Kuşkuran Deresi, 1900 m, 23.x.1979. N. Çelik- 947 HUBI; ibid. 23.vii.1980, N. Çelik- 1545 ANKI; Hınzır Dağı, above Kaynar, Erikli Dere, 2050 m, 12.x.1979. N. Çelik-822 HUBI; Hınzır Dağı, Akkışla, above Ganişeyh, Taşlı Dere, 1800 m, 19.viii.1979. N. Çelik-740 HUBI; Hınzır Dağı, Akkışla, above Ortaköy, 1950 m, 15.ix.1979. N. Çelik-890 HUBI; Pınarbaşı, east of Hınzır Dağı, Kaynar, Bademli Dere, 2000 m, 19.vii.1979. N. Çelik-650 HUBI; Pınarbaşı, south of Hınzır Dağı, Kaynar, 2000 m, 18.vii.1979. N. Çelik-586 HUBI; ibid viii.1994. Aytaç 6727 Ekici & Dönmez; GAZI, Sarız, Binboğa Dağı. G. Tümen ESSE 95711; ibid. viii.1992. G. Tümen ESSE 98311; ibid.

vii.1995. G. Tümen ESSE 11952I; ibid. Çağsak Köyü, vii.1993. G. Tümen ESSE 10432I; B6 K. Maraş: Göksun, Fındık Köyü, Kandil Dağı, 1600-1800 m, clearing forest, 27.vii.1981. B. Yıldız- 3114 HUBI; Göksun, north of Fındık, Bostandere, 1500m, clearing forest, 24.x.1980. B. Yıldız -2348 HUBI; B6 K. Maraş: Binboğa Dağı, 2300-2400 m, 17.vii.1952. P.H.Davis 20102 ANKI, KI, EI; B6 Kayseri: Bakır Dağı, between Kisge and Bakıroluğu Y., 1700 m, in open *Abietum*, 28.vi.1952, Davis 19477, Dodds&Çetik, KI.

This species is similar to *S. germanicopolitana* and *S. libanotica* subsp. *kurdica*. From *S. germanicopolitana*, it differs middle cauline leaves 3.5-8x0.3-1 cm (not 5-12x1-2 cm); middle bracts 1-1.5 cm (not 1-6 cm); calyx 9-10 mm, long simple and short glandular hairs (not 9-13 mm, eglandular hairs); corolla 10-12 mm (not 12-14 mm). From *S. libanotica* subsp. *kurdica*, it differs cauline leaves 2-4-paired (not 4-8-paired); bracts glandular and eglandular hairy (not only with eglandular densely hairs); calyx 9-10 mm, long simple and short glandular hairy (not ca. 6 mm long and densely white- villous), calyx teeth 3-4 mm (not 2.5-3 mm). This new species was doubtfully identified by N. Çelik, *R. İlarıslan*, *B. Yıldız* and Huber-Morath as *S. libanotica* subsp. *kurdica* before.

Sideritis vuralii Duman et Başer spec. nov. (Sect. *Empedoclia* (Rafin.) Benth) Fig.2, Map1.

Affinis *S. syriacae* L. subsp. *nusairiensi* (Post) Hub.-Mor. et *S. hololeucae* Boiss. & Heldr. sed ab *S. syriacae* subsp. *nusairiensi* bracteis medianis calycum brevioribus, 0.4-0.6 x 0.4-0.7 cm (non 0.6-1 cm), calycibus 6-7 mm (non 7.5-9 mm) longis et calyci dentibus 1-2 mm (non 2-2.5 mm) longis differt. Ab *S. hololeucae* foliis caulinis medianis inferioribusque oblan-



Figure 2. *Sideritis vuralii*: A-habit, B-bract, C-calyx, D-corolla

ceolatis usque obovatis, lamina 3-5 cm (non orbicularibus usque suborbicularibus, lamina 1-2.5 cm longa), verticillastris 4-12 (non (1-) 2-5) differt.

Perennial, 30-70 cm, erect, branched, densely adpressed white-pannose, eglandular. Basal leaves numerous, forming a rosette, distinctly petiolate, petioles 2-4 cm; lamina oblanceolate-spathulate or obovate, 1.5-5 x 0.5-2 cm, entire to finely crenulate-serrate. Middle cauline leaves oblanceolate to obovate, lamina 3-5x0.8-1.5 cm, entire to minutely crenate, obtuse, abruptly constricted into petiole to 1.5 cm. Internodes to 12 cm. Inflorescence simple or 2-6-branched, 5-20 cm; verticillasters 4-12, (5-) 6-flowered, lowermost (0.5-) 2-5 (-6) cm distant, upper sometimes crowded. Middle bracts ovate-orbicular to cordate, appearing connate because of interlocking indumentum, densely white-pannose, eglandular, always shorter than calyx, 0.4-0.6x0.4-0.7 cm, mucronate, mucro 1-3 mm. Calyx 6-7 mm, densely eglandular arachnoid pilose; teeth triangular-lanceolate, obtuse to slightly acute, 1-2 mm long. Corolla yellow, c. 10 mm, hairy outside and inside, with brown striations inside; limb c. 3 mm. Nuts triangular-ovoid, brown, glabrous, c. 2 mm. *Fl. July-August. Calcareous rocky place, 1200-1900 m.*

Type: Turkey, C4 İçel: Anamur, Kazancı-Abanoz, 1300 m, clearing forest and calcareous rocky place, 19.viii.1994. *M. Vural* 7165- *N. Adıgüzel* & *A. Dönmez* (holo- GAZI; iso- ESSE, ANK).

South Anatolia. Endemic. Mediterranean element. C4 İçel: Anamur, Kaş-Abanoz Yaylası, 1485 m, calcareous rocky place, 18.vii.1995. *H. Duman* 5846- *K.H.C. Başer* and *A. Altıntaş*; ibid. ESSE 11633I; ibid. vii.1991. *G. Tümen* ESSE 9572I; Abanoz-Akpinar Yaylası, 15.viii.1993. *G. Tümen* ESSE 10425I; ibid. 20.vi.1991. *G. Tümen* ESSE 9444I; C4 İçel: Gülnar-Ermenek, 1220 m, *Juniperus* forest, rocky place, 25.viii.1992. *M. Vural* 6110- *M. Koyuncu* & *M. Ekici* GAZI; C4 Konya: south of Ermenek, Sarıvadi, 1300 m, 30.viii.1993. *G. Tümen* ESSE 10140I; C4 Antalya: Gazipaşa, Sugözü village, Akçal Dağı, 1900 m, 5.viii.1984, *H. Sümbül* 3252 HUBI, ANKI [identified by *H. Sümbül* *S. syriaca* L. subsp. *nusairiensis* (Post) Hub.-Mor.].

Allied to *S. syriaca* L. subsp. *nusairiensis* which is distributed in C5 and C6 squares, but *S. vuralii* differs in its middle bracts always long shorter than calyx, 0.4-0.6x cm (not 0.6-1 cm) and calyx 6-7 mm (not 7.5-9 mm), teeth 1-2 mm (not 2-2.5 mm). Also allied to *S. hololeuca*, but middle and lower cauline leaves oblanceolate to obovate, lamina 3-5 cm (not orbicular

to suborbicular, lamina 1-2.5 cm), verticillasters 4-12 (not (1-)2-5) differing.

This species is named in honour of Prof. Dr Mecit Vural a well-known phytosociologist and taxonomist at Gazi University, Ankara.

Origanum x adanense Başer & Duman hybr. nov.

(*O. bargyli* Mouterde x *O. laevigatum* Boiss.) Fig. 3, Map1.

Affinis *O. laevigato* Boiss. sed spiculis 10-20 x 7-12 mm (non 5-20 x 3-6 mm), bracteis 5-8 x 2-5 mm (non 3-6 x 0.5-2 mm), calyce 2-chilo. Ab *O. bargyli* Mouterde bracteis brevioribus 5-8 x 2-5 mm (non 5-17 x 3-13) et calycibus 5-6 mm, (non 5-8 mm) differt.

Subshrub. Young shoots hirtellous. Stems 30-50 cm, ascending erect, light or purplish brown, somewhat hirtellous at the base (hairs to 1 mm long), otherwise glabrous. Branches of first order present in the upper 1/2-1/3 of the stems, up to 5 pairs per stem, (1-)3-5(-15) cm long. Leaves up to 15 pairs per stem, shortly petiolate or sessile, ovate to cordate, obtuse or acute, 8-25 x 5-15 mm, glaucous, green or purplish, glabrous or sparsely hirtellous or scabrous at least at the margins, sessile glands up to 300 per square cm. Spikes ovate to cylindrical in outline, 10-20 x 7-12 mm, erect or sometimes nodding. Bracts 4-8 pairs per spike, ovate to lanceolate or rhomboid, acute, 5-8 x 2-5 mm, glabrous or scabrous at the margins, partly purple. Flowers subsessile, pedicels 0.5-1 mm. Calyx 5-6 mm, outside glabrous, throats pilose; upper lips triangular, teeth c. 1 mm, lower lips approximately as long as upper lips, triangular-lanceolate, teeth c.2 mm. Corolla 2-lipped for c. 1/6, 11-13 mm long, pink, not saccate, outside pilose. Stamens; the upper 2 included, the lower 2 shortly protruding; filaments c. 0.8-1 mm and 2-3 mm long. *Fl. July. Pinus nigra clearings, 1100-1200m.*

Type: Turkey, C6 Osmaniye: Amanos Dağı, Yarpuz-Ağulu, 1190m, *Pinus nigra* forest, 20.vii.1995. *H. Duman* 5881- *K.H.C. Başer* and *A. Altıntaş* (holo- GAZI; iso- ESSE, ANK).

South Anatolia. Endemic. Mediterranean element. This new hybrid has become *O. laevigatum* Boiss. and *O. bargyli*. We found that the same locality with *O. bargyli*. *O. x adanense* similar to *O. laevigatum* as appearance which is distributed Amanos mountain and Cyprus. But, *O. x adanense* differs in its spicules 10-




Figure 3. *Origanum x adanense*: A-habit, B-Bract, C-dissected calyx, D-dissected corolla; *O. bargyli*: E-bract, F-dissected calyx; *O. laevigatum*: G-bract, H- dissected calyx

20 x 7-12 mm (not 5-20 x 3-6 mm), bracts 5-8 x 2-5 mm (not 3-6 x 0.5-2 mm) and Calyx 2-lipped (not 1-lipped). From *O. bargyli* Mouterde, it differs bracts smaller (not 5-17 x 3-13 mm) and calyx 5-6 mm (not 5-8 mm).

References

1. Huber-Morath, A., Sideritis L. in Davis, P. H., Flora of Turkey and the East Aegean Islands, 7:178-199, 297-313, Edinburgh Univ. Pres. (1982).
2. Davis, P. H., Mill, R.R., Tan, K.,(eds.), Flora of Turkey and the East Aegean Islands, 10:203, 206-207, Edinburgh Univ. Pres. (1988).

Acknowledgements

We thanks Prof. Dr Tuna Ekim and Prof. Dr Mecit Vural for their critical discussion, TÜBİTAK (TBAG-940) and TBAM for financial support. We are indebted to Dr. Robert Mill and Ian Hedge (Royal Botanic Garden Edinburg) for checking the latin diagnosis. Also thanks Figen Özkaya for illustration.

3. Duman, H., Aytaç, Z., Ekici, M., Karavelioğulları, F.A., Dönmez, A.A.,Duran A., Three New Species (Labiatae) From Turkey, Flora Mediteranea,5:221-228, Palermo (1995).
4. Başer, K.C.H., Vural, M., Tümen, G., Akyalçın, H., Satıl, F., Two New Records for the Flora of Turkey, Tr. J. of Botany, 19, 489-490 (1995).
5. Letswaart, J.H., A Taxonomic Revision of the Genus *Origanum* L. (Labiatae), Leiden Botanical Series. no:4, Leiden Univ. press. (1980).