

1-1-2009

Perturbation of Closed Range Operators

MOHAMMAD SAL MOSLEHIAN

GHADIR SADEGHI

Follow this and additional works at: <https://dctubitak.researchcommons.org/math>



Part of the [Mathematics Commons](#)

Recommended Citation

MOSLEHIAN, MOHAMMAD SAL and SADEGHI, GHADIR (2009) "Perturbation of Closed Range Operators," *Turkish Journal of Mathematics*: Vol. 33: No. 2, Article 5. <https://doi.org/10.3906/mat-0805-26>
Available at: <https://dctubitak.researchcommons.org/math/vol33/iss2/5>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Mathematics by an authorized editor of TÜBİTAK Academic Journals.

Perturbation of Closed Range Operators

Mohammad Sal Moslehian and Ghadir Sadeghi

Abstract

Let T, A be operators with domains $\mathcal{D}(T) \subseteq \mathcal{D}(A)$ in a normed space X . The operator A is called T -bounded if $\|Ax\| \leq a\|x\| + b\|Tx\|$ for some $a, b \geq 0$ and all $x \in \mathcal{D}(T)$. If \mathcal{A} has the Hyers–Ulam stability then under some suitable assumptions we show that both T and $S := A + T$ have the Hyers–Ulam stability. We also discuss the best constant of Hyers–Ulam stability for the operator S . Thus we establish a link between T -bounded operators and Hyers–Ulam stability.

Key Words: Hilbert space; perturbation; Hyers–Ulam stability; closed operator; semi-Fredholm operator.

1. Introduction and preliminaries

Let X, Y be normed linear spaces and T be a (not necessarily linear) mapping from X into Y . Following [5, 6] we say that T has the Hyers-Ulam stability if there exists a constant $K > 0$ with the property:

(i) For any y in the range $\mathcal{R}(T)$ of T , $\varepsilon > 0$ and $x \in X$ with $\|T(x) - y\| \leq \varepsilon$, there exists a $x_0 \in X$ such that $T(x_0) = y$ and $\|x - x_0\| \leq K\varepsilon$.

We call such $K > 0$ a Hyers-Ulam stability constant for T and denote by K_T the infimum of all Hyers-Ulam stability constants for T . If K_T is a Hyers-Ulam stability constant for T , then K_T called the Hyers-Ulam stability constant for T .

If T is linear then condition (i) is equivalent to:

(ii) For any $\varepsilon > 0$ and $x \in X$ with $\|Tx\| \leq \varepsilon$, there exists a $x_0 \in X$ such that $Tx_0 = 0$ and $\|x - x_0\| \leq K\varepsilon$.

If put $\mathcal{N}(T) := \{x \in X : Tx = 0\}$, condition (ii) is equivalent to

(iii) For any $x \in X$ there exists a $x_0 \in \mathcal{N}(T)$ such that $\|x - x_0\| \leq K\|Tx\|$.

We refer the interested reader for more results on the stability of various mappings to papers [10, 11, 12] and references therein, and for a comprehensive accounts of the Hyers-Ulam-Rassias stability of functional equations to the monographs [3, 8, 13].

In [6] the authors proved the following useful result.

2000 AMS Mathematics Subject Classification: Primary 47A55; Secondary 39B52, 34K20, 39B82.

Theorem 1.1 *Let T be a closed operator from the subspace $\mathcal{D}(T)$ of a Hilbert space \mathcal{H} into a Hilbert space \mathcal{K} . The following assertions are equivalent:*

- (i) T has the Hyers-Ulam stability;
- (ii) T has closed range.

Moreover, if one of the conditions above is true, then $K_T = \gamma(T)^{-1}$, where

$$\gamma(T) = \sup\{\gamma > 0 : \|Tx\| \geq \gamma\|x\|, \quad x \in \mathcal{D}(T) \cap (\mathcal{N}(T))^\perp\}.$$

(Here \perp denotes the orthogonal complement in Hilbert spaces.)

Let X be a Banach space and let M, N be closed linear subspaces of X . Following [9] we define the quantity

$$\delta(M, N) := \inf\left\{\frac{\text{dist}(x, N)}{\text{dist}(x, M \cap N)} : x \in M, x \notin N\right\} (\leq 1)$$

If $M \subseteq N$, then we set $\delta(M, N) = 1$. Obviously $\delta(M, N) = 1$, if $M \supseteq N$. It is well known that $\delta(M, N)$ is not symmetric with respect to (M, N) . If $\delta(M, N) = \delta(N, M)$, we say that the pair (M, N) is regular. It is known that any pair (M, N) is regular if X is a Hilbert space [9].

Let A and T be operators with their domains in a normed space X such that $\mathcal{D}(T) \subseteq \mathcal{D}(A)$, and

$$\|Ax\| \leq a\|x\| + b\|Tx\| \quad (x \in \mathcal{D}(T)), \tag{1.1}$$

where a, b are nonnegative constants. Then we say that A is relatively bounded with respect to T or simply it is T -bounded [9].

A bounded operator A is clearly T -bounded for any T with $\mathcal{D}(T) \subseteq \mathcal{D}(A)$.

In this paper, we show that if a T -bounded operator A has the Hyers-Ulam stability then under some suitable assumptions the operator T and the perturbation $S := A + T$ have the Hyers-Ulam stability. We also discuss the best constant of Hyers-Ulam stability for the operator S . Thus we establish a link between T -bounded operators and the Hyers-Ulam stability.

2. Main Results

Throughout this section \mathcal{H} and \mathcal{K} denote Hilbert spaces and A and T are operators having their domains in \mathcal{H} and their images in \mathcal{K} . We start our work with the following theorem.

Theorem 2.1 *Suppose that A is a T -bounded operator with a T -bound smaller than 1. If T is a closed operator and $S := T + A$, then the following assertions are equivalent:*

- (i) S has the Hyers-Ulam stability;
- (ii) S has closed range.

Moreover, if A is closed and the operators A and T have the Hyers-Ulam stability and $\mathcal{R}(S) = \mathcal{R}(A) + \mathcal{R}(T)$ then conditions (i) and (ii) are equivalent with the following assertions:

- (iii) $\delta(M, N) > 0$, where $M = \mathcal{R}(A)$ and $N = \mathcal{R}(T)$;
- (v) $\delta(M^\perp, N^\perp) > 0$, $M = \mathcal{R}(A)$ and $N = \mathcal{R}(T)$.

Proof. The operator S is closed since the operator A is T -bounded with a T -bound smaller than 1 and T is a closed operator (see [9, Theorem 1.1]). It follows from [6, Theorem 3.1] that operator S has the Hyers-Ulam stability if and only if S has closed range. Hence (i) \iff (ii).

Now, if $\mathcal{R}(S) = \mathcal{R}(A) + \mathcal{R}(T)$ and A and T have the Hyers-Ulam stability then $\mathcal{R}(A)$ and $\mathcal{R}(T)$ are closed and Theorems 4.2 and 4.8 of [9] show that (ii) \iff (iii) and (iii) \iff (v). \square

Remark 2.2 *If A and T are closed operators as in the above theorem, the operators A and T have the Hyers-Ulam stability, $S := T + A$, $\mathcal{R}(S) = \mathcal{R}(A) + \mathcal{R}(T)$ and we have $\mathcal{R}(A) \subseteq \mathcal{R}(T)$ or $\mathcal{R}(T) \subseteq \mathcal{R}(A)$ then $\delta(\mathcal{R}(A), \mathcal{R}(T)) > 0$. Hence the operator S has the Hyers-Ulam stability and therefore its range is closed.*

Corollary 2.3 *Suppose that A is a T -bounded operator with a T -bound smaller than 1. Let A and T be closed, $S := A + T$ and let A and T have the Hyers-Ulam stability. Suppose that at least one of the spaces $\mathcal{R}(A)$ or $\mathcal{R}(T)$ is finite dimensional and assume that $\mathcal{R}(S) = \mathcal{R}(A) + \mathcal{R}(T)$. Then operator S has the Hyers-Ulam stability and so it has closed range.*

Proof. Without loss of generality assume that $\mathcal{R}(A)$ is finite dimensional. It is known that there exists $u \in \mathcal{R}(T)$ such that $\text{dist}(u, \mathcal{R}(A)) = \|u\|$ (see [2]). Hence

$$\delta(\mathcal{R}(A), \mathcal{R}(T)) = \delta(\mathcal{R}(T), \mathcal{R}(A)) > 0.$$

Therefore operator $S = T + A$ has the Hyers-Ulam stability. \square

Corollary 2.4 *Suppose that A is a T -bounded operator with a T -bound smaller than 1. Let A and T be closed, $S := A + T$ and let A , T and S have the Hyers-Ulam stability. If $\mathcal{R}(A) \cap \mathcal{R}(T) = \{0\}$, then $\delta(\mathcal{R}(T), \mathcal{R}(A)) = 1$ and*

$$K_S \leq \min\left\{\frac{1}{\gamma(T)}, \frac{1}{\gamma(A)}\right\}.$$

Proof. Each $z \in \mathcal{R}(S)$ has a unique expression as $z = x + y$ in which $y \in \mathcal{R}(T)$ and $x \in \mathcal{R}(A)$. Consider the projection P of $\mathcal{R}(S)$ onto $\mathcal{R}(T)$ along $\mathcal{R}(A)$. Now we have

$$1 = \|P\| = \sup_{z \in \mathcal{R}(S)} \frac{\|Pz\|}{\|z\|} = \sup_{y \in \mathcal{R}(T), x \in \mathcal{R}(A)} \frac{\|y\|}{\|x + y\|} = \sup_{y \in \mathcal{R}(T)} \frac{\|y\|}{\text{dist}(y, \mathcal{R}(A))} = \delta(\mathcal{R}(A), \mathcal{R}(T))^{-1}.$$

By the definition of $\gamma(T)$, we have $\|Tv\| \geq \gamma(T)\|v\|$. Hence $\|P\|\|Tv + Av\| \geq \|P(Tv + Av)\| \geq \gamma(T)\|v\|$. So $\|Sv\| \geq \gamma(T)\|v\|$. Since $\gamma(S) \geq \gamma(T)$, by [6, Theorem 3.1], we have $K_S \leq \frac{1}{\gamma(T)}$. We can analogously show that $K_S \leq \frac{1}{\gamma(A)}$. Thus $K_S \leq \min\left\{\frac{1}{\gamma(A)}, \frac{1}{\gamma(T)}\right\}$. \square

Recall that if x, y are elements of the Hilbert space \mathcal{H} , then the bounded operator $x \otimes y$ defined on \mathcal{H} by $(x \otimes y)(z) = \langle z, y \rangle x$ is rank one if x, y are not zero. Let x_1, x_2, y be elements of \mathcal{H} such that $\|x_1\| \leq \frac{\|x_2\|}{2}$. If $A = x_1 \otimes y, T = x_2 \otimes y$ and $S = A + T$, then $\mathcal{N}(A) = \mathcal{N}(T)$ and $\|Ax\| \leq \frac{\|Tx\|}{2}$. It is clear that A, T and

S have the Hyers-Ulam stability (note that they have closed range). This motivates us toward the following theorem.

Theorem 2.5 *Suppose that A is a T -bounded operator with a T -bound b and a constant a and A has the Hyers-Ulam stability.*

If $a = 0$ and $\mathcal{N}(A) = \mathcal{N}(T)$, then T has also the Hyers-Ulam stability.

Proof. There exists a constant $K_0 > 0$ such that for every $x \in \mathcal{D}(A)$ there exists $x_0 \in \mathcal{N}(A) = \mathcal{N}(T)$ such that $\|x - x_0\| \leq K_0\|Ax\| \leq K_0b\|Tx\|$. Thus operator T has the Hyers-Ulam stability. \square

Now we show that conditions $\mathcal{N}(A) = \mathcal{N}(T)$ and $a = 0$ in Theorem 2.5 are necessary.

Example 2.6 *Consider the operators $A, T : \ell^2 \longrightarrow \ell^2$ defined by*

$$A(x_1, x_2, \dots) = (x_1, 0, 0, \dots), \quad (x_1, x_2, \dots) \in \ell^2$$

and

$$T(x_1, x_2, \dots) = (x_1, \frac{x_2}{2}, \frac{x_3}{3}, \dots), \quad (x_1, x_2, \dots) \in \ell^2.$$

It is clear that the operator A is T -bounded with constant $a = 0$. Then $\mathcal{R}(A)$ is of finite dimension. Hence the operator A has closed range. Hence A has the Hyers-Ulam stability and $\mathcal{N}(A) \neq \mathcal{N}(T)$. If we take a_n to be

$$a_n = \begin{cases} 1 & i \leq n \\ 0 & i > n \end{cases}$$

then

$$(Ta_n)(i) = \begin{cases} 1/i & i \leq n \\ 0 & i > n \end{cases}$$

and (Ta_n) converges to $b = (1, \frac{1}{2}, \frac{1}{3}, \dots)$ which does not belong to the range of T . Therefore $\mathcal{R}(T)$ is not closed, i.e, operator T does not have the Hyers-Ulam stability.

Example 2.7 *Consider the operators $A, T : \ell^2 \longrightarrow \ell^2$ defined by*

$$A(x_1, x_2, \dots) = (0, x_1, x_2, \dots), \quad (x_1, x_2, \dots) \in \ell^2$$

and

$$T(x_1, x_2, \dots) = (x_1, \frac{x_2}{2}, \frac{x_3}{3}, \dots), \quad (x_1, x_2, \dots) \in \ell^2.$$

The operator A is T -bounded with a nonzero constant a . Since $\gamma(A) > 0$, the operator A has closed range and $\mathcal{N}(A) = \mathcal{N}(T)$. The space $\mathcal{R}(T)$ is not closed, i.e, operator T does not have the Hyers-Ulam stability.

Let x_1, x_2, y be elements of \mathcal{H} such that $x_1 \perp x_2$. If $A = x_1 \otimes y, T = x_2 \otimes y$ and $S = A + T$, then $\gamma(A) = \|x_1\|\|y\|, \gamma(T) = \|x_2\|\|y\|$ and $\gamma(S) = \gamma(A) + \gamma(T)$, therefore $K_S = \gamma(S)^{-1} = \frac{1}{\gamma(A) + \gamma(T)}$. This motivates us toward the following result.

Corollary 2.8 *Suppose that A is a T -bounded operator with a T -bound b smaller than 1 and constant $a = 0$, $\mathcal{N}(A) = \mathcal{N}(T)$ and A has the Hyers-Ulam stability. Then $S := T + A$ has the Hyers-Ulam stability, if $\mathcal{R}(A) \perp \mathcal{R}(T)$. Moreover, if T is a closed operator then $\mathcal{R}(S)$ is closed and $K_S = \frac{1}{\gamma(T) + \gamma(A)}$.*

Proof. Suppose that K is a Hyers-Ulam stability constant for A . By Theorem 2.5, $K' = Kb$ is a Hyers-Ulam stability constant for T . In fact, for each $v \in \mathcal{D}(T)$ there exists $v_0 \in \mathcal{N}(T)$ such that

$$\|v - v_0\| \leq (Kb)\|Tv\| \leq K\|Tv\|$$

since b is smaller than 1.

Hence for $x \in \mathcal{D}(S) = \mathcal{D}(T)$ there exists $x_0 \in \mathcal{N}(T) = \mathcal{N}(A)$ such that

$$\|x - x_0\| \leq K(\|Ax\| + \|Tx\|) = K\|Ax + Tx\|.$$

Now we show that $\mathcal{N}(S) = \mathcal{N}(T)$. If $x \in \mathcal{N}(S) - \mathcal{N}(T)$, then $-Ax = Tx$ and so $\|Tx\| = \|Ax\| \leq b\|Tx\|$. Hence $b \geq 1$ which is a contradiction. Thus $\mathcal{N}(S) \subseteq \mathcal{N}(T)$ since $\mathcal{N}(A) = \mathcal{N}(T)$ and $\mathcal{N}(T) \subseteq \mathcal{N}(S)$. Therefore $\mathcal{N}(S) = \mathcal{N}(T)$. Thus S has the Hyers-Ulam stability.

Assume that T is a closed operator. Then so is S . Hence $\mathcal{R}(S)$ is closed. Since $\frac{\|Sx\|}{\|x\|} = \frac{\|Tx + Ax\|}{\|x\|} = \frac{\|Tx\|}{\|x\|} + \frac{\|Ax\|}{\|x\|}$ and $\mathcal{N}(T) = \mathcal{N}(S)$ we have $\gamma(S) = \gamma(T) + \gamma(A)$. Hence, by [6, Theorem 3.1], $K_S = \frac{1}{\gamma(T) + \gamma(A)}$. \square

The following result can be regarded as a special case of [1, Theorem 2.2] with a Hyers-Ulam stability approach.

Theorem 2.9 *Suppose that A is a T -bounded operator with a T -bound b smaller than 1 and constant $a = 0$, and $\mathcal{N}(A) = \mathcal{N}(T)$. Assume that A has the Hyers-Ulam stability and that T is a closed operator. Then $S := T + A$ is a closed operator, S has the Hyers-Ulam stability and*

$$\frac{1}{\gamma(A) + \gamma(T)} \leq K_S \leq \frac{1}{(1 - b)\gamma(T)}.$$

Proof. By Theorem 2.5 the operator T has the Hyers-Ulam stability. Hence it has closed range and so $\gamma(T) > 0$. Since the operator A is T -bounded with a T -bound smaller than 1 and since by [9, Theorem 1.1] T is a closed operator, we deduce that the operator S is closed. In view of $\|Ax\| \leq b\|Tx\|$, we get

$$\|Tx\| - \|Sx\| \leq \|Ax + Tx - Tx\| \leq b\|Tx\| \quad (x \in \mathcal{D}(T)).$$

Hence $(1 - b)\|Tx\| \leq \|Sx\|$. Thus

$$(1 - b) \frac{\|Tx\|}{\|x\|} \leq \frac{\|Sx\|}{\|x\|} \quad x \in (\mathcal{D}(T) - \{0\}).$$

Since $\mathcal{N}(T) = \mathcal{N}(S)$ we have $0 < (1 - b)\gamma(T) \leq \gamma(S)$, therefore S has closed range [9, Theorem 5.2]. Thus S has the Hyers-Ulam stability and $K_S = \gamma(S)^{-1} \leq \frac{1}{(1-b)\gamma(T)}$. Clearly $\gamma(S) \leq \gamma(A) + \gamma(T)$. Therefore $\frac{1}{\gamma(A)+\gamma(T)} \leq K_S$. \square

Recall that a closed operator A from \mathcal{H} into \mathcal{K} is called left semi-Fredholm if $\dim\mathcal{N}(A) < \infty$ and $\mathcal{R}(A)$ is closed. It is called right semi-Fredholm if $\text{codim}\mathcal{R}(A) < \infty$ and $\mathcal{R}(A)$ is closed. We say a closed operator A is semi-Fredholm if it is left or right semi-Fredholm.

Remark 2.10 *Suppose that A is a T -bounded operator with a T -bound b smaller than 1 and constant $a = 0$, and $\mathcal{N}(A) = \mathcal{N}(T)$. If T is a closed operator and has the Hyers-Ulam stability. Then, by Theorem 2.9, the operator $S := A + T$ is closed and has the Hyers-Ulam stability. So that $\mathcal{R}(S)$ is closed.*

The conclusion that S is closed has already obtained in [4, Theorem V.3.6] under the different assumption that the operator T is semi-Fredholm.

Corollary 2.11 *Suppose that A is a left semi-Fredholm and T -bounded operator with constant $a = 0$ and a T -bound b smaller than 1, and T is a closed operator such that $\mathcal{N}(A) = \mathcal{N}(T)$. Then $S := T + A$ is a left semi-Fredholm operator.*

Theorem 2.12 *Suppose that A is a T -bounded operator with a T -bound b smaller than 1 and constant $a = 0$, and $\mathcal{N}(A) = \mathcal{N}(T)$. If $S = T + A$ has the Hyers-Ulam stability then T has the Hyers-Ulam stability. Moreover if S is a closed operator then $\mathcal{R}(S)$ and $\mathcal{R}(T)$ are closed.*

Proof. The operator S has the Hyers-Ulam stability thus there exists a constant $K > 0$ with the following property:

$$\text{For any } x \in \mathcal{D}(S) = \mathcal{D}(T) \text{ there exists a } x_0 \in \mathcal{N}(S) \text{ such that } \|x - x_0\| \leq K\|Sx\|.$$

Since A is a T -bounded operator and, by the proof of Corollary 2.8, $\mathcal{N}(T) = \mathcal{N}(S)$, we have

$$\|x - x_0\| \leq K\|Sx\| \leq K(\|Ax\| + \|Tx\|) \leq K(b + 1)\|Tx\|.$$

Therefore T has the Hyers-Ulam stability.

Now assume that S is a closed operator. Then so is T . In view of S and T having the Hyers-Ulam stability, $\mathcal{R}(S)$ and $\mathcal{R}(T)$ are closed. \square

References

- [1] Christensen, O: *Operators with closed range, pseudo-inverses, and perturbation of frames for a subspace.* Canad. Math. Bull. **42**, no. 1, 37–45 (1999).
- [2] Conway, J.B.: *A Course in Operator Theory*, Graduate Studia in Mathematics Volume 21, Amer. Math. Soc., 1999.
- [3] Czerwik, S.: *Functional equations and inequalities in several variables*, World Scientific, New Jersey, London, Singapore, Hong Kong, 2002.

- [4] Goldberg S.: *Unbounded linear Operators*, McGraw-Hil, New Yourk, 1966.
- [5] Hatori, O., Kobayashi, K., Miura, T. , Takagi, H., Takahasi, S.E.: *On the best constant of Hyers–Ulam stability*, J. Nonlinear Convex Anal. **5**, 387–393 (2004).
- [6] Hirasawa, G., Miura, T. *Hyers–Ulam Stability of a closed operator in a Hilbert space*, Bull. Korean. Math. Soc. **43**, 107–117 (2006).
- [7] Hyers, D.H., Isac, G., Rassias, Th.M.: *Stability of Functional Equations in Several Variables*, Birkhäuser, Basel, 1998.
- [8] Jung, S.-M.: *Hyers–Ulam–Rassias Stability of Functional Equations in Mathematical Analysis*, Hadronic Press, Palm Harbor, 2001.
- [9] Kato, T.: *Perturbation Theory for Linear Operators*, Springer-Verlag, New York, 1966.
- [10] Moslehian, M.S.: *Ternary derivations, stability and physical aspects*, Acta Applicandae Math. **100**, no. 2, 187–199 (2008).
- [11] Moslehian, M.S., Rassias, Th.M.: *Stability of functional equations in non-Archimedean spaces*, Appl. Anal. Disc. Math. **1**, no. 2, 325–334 (2007).
- [12] Rassias, Th.M.: *On the stability of functional equations and a problem of Ulam*, Acta Appl. Math. **62** , no. 1, 23–130 (2000).
- [13] Rassias, Th.M.: *Functional Equations, Inequalities and Applications*, Kluwer Academic Publishers, Dordrecht, Boston and London, 2003.

Mohammad Sal MOSLEHIAN^{1,2}, Ghadir SADEGHI^{1,2}

Received 21.05.2008

¹Department of Pure Mathematics,

Ferdowsi University of Mashhad, P. O. Box

1159, Mashhad 91775, IRAN

e-mail: moslehian@ferdowsi.um.ac.ir and moslehian@ams.org

²Center of Excellence in Analysis

on Algebraic Structures (CEAAS),

Ferdowsi University of Mashhad-IRAN

e-mail: ghadir54@yahoo.com