

1-1-2008

Posner's Second Theorem and an Annihilator Condition with Generalized Derivations

VINCENZO DE FILIPPIS

Follow this and additional works at: <https://journals.tubitak.gov.tr/math>



Part of the [Mathematics Commons](#)

Recommended Citation

FILIPPIS, VINCENZO DE (2008) "Posner's Second Theorem and an Annihilator Condition with Generalized Derivations," *Turkish Journal of Mathematics*: Vol. 32: No. 2, Article 6. Available at: <https://journals.tubitak.gov.tr/math/vol32/iss2/6>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Mathematics by an authorized editor of TÜBİTAK Academic Journals. For more information, please contact academic.publications@tubitak.gov.tr.

Posner's Second Theorem and an Annihilator Condition with Generalized Derivations

Vincenzo De Filippis

Abstract

Let R be a prime ring of characteristic different from 2, with extended centroid C , U its two-sided Utumi quotient ring, $\delta \neq 0$ a non-zero generalized derivation of R , $f(x_1, \dots, x_n)$ a non-central multilinear polynomial over C in n non-commuting variables, $a \in R$ such that $a[\delta(f(r_1, \dots, r_n)), f(r_1, \dots, r_n)] = 0$, for any $r_1, \dots, r_n \in R$. Then one of the following holds:

1. $a = 0$;
2. there exists $\lambda \in C$ such that $\delta(x) = \lambda x$, for all $x \in R$;
3. there exist $q \in U$ and $\lambda \in C$ such that $\delta(x) = (q + \lambda)x + xq$, for all $x \in R$, and $f(x_1, \dots, x_n)^2$ is central valued on R .

Key Words: Prime rings, derivations, left Utumi quotient rings, two-sided Martindale quotient ring, differential identities.

1. Introduction

The well known theorem of Posner in [18] asserts that if R is a prime ring and d a non-zero derivation of R such that $[d(x), x] \in Z(R)$, the center of R , for all $x \in R$, then R must be commutative. Starting from this result, several authors studied the relationship between the structure of a prime ring R and the behaviour of an additive mapping f which satisfies the Engel-type condition $[f(x), x]_k = 0$ which, for $k > 1$, is defined by $[f(x), x]_k = [[f(x), x]_{k-1}, x]$. In [10] Lanski shows that if d is a derivation of R such that $[d(x), x]_k = 0$, for all x in a Lie ideal L of R , then either L is central

2000 AMS Mathematics Subject Classification: 16N60, 16W25.

in R or $\text{char}(R) = 2$ and R satisfies the standard identity $s_4(x_1, \dots, x_4)$ of degree 4. On the other hand, in a prime ring R of characteristic different from 2, any non-central Lie ideal contains the set $\{[x_1, x_2] : x_1, x_2 \in I\}$ of all evaluations of the polynomial $[x_1, x_2]$ in a two-sided ideal I of R . For this reason, many researchers in this area analysed in detail the case when the Lie ideal is replaced by the set of all evaluations of a polynomial $f(x_1, \dots, x_n)$ and $[d(f(x_1, \dots, x_n)), f(x_1, \dots, x_n)]_k$ is a differential identity for some ideal of R . In particular we refer the reader to the results obtained by P. H. Lee and T. K. Lee in [12] and [13]. In case $f(x_1, \dots, x_n)$ is a multilinear polynomial, they prove that it must be central-valued in R unless $\text{char}(R) = 2$ and R satisfies $s_4(x_1, \dots, x_4)$. In a recent paper, we consider another related generalization; more precisely in [4] we describe what happens if the derivation d is replaced by an additive mapping δ defined as follows: for all $x, y \in R$, $\delta(xy) = \delta(x)y + xg(y)$, for some derivation g of R . Such a mapping δ is called a generalized derivation. Obviously any derivation is a generalized derivation. We like to remark that one of the leading roles in the development of the theory of generalized derivations is played by the maps defined as $\delta(x) = bx + xc$ for $b, c \in R$; in this case δ is called inner generalized derivation.

In the light of these definitions, in [4] we prove that if I is a right ideal of R , U the two-sided Utumi quotient ring of R , $C = Z(U)$ the extended centroid of R and $[\delta(f(r_1, \dots, r_n)), f(r_1, \dots, r_n)] = 0$, for any $r_1, \dots, r_n \in I$, then either $\delta(x) = ax$, with $(a - \gamma)I = 0$ and suitable $a \in U$, $\gamma \in C$ or there exists an idempotent element $e \in \text{soc}(RC)$ such that $IC = eRC$ and one of the following holds: (i) $f(x_1, \dots, x_n)$ is central valued in $eRCe$; (ii) $\delta(x) = cx + xb$, where $(c + b + \alpha)e = 0$, for $b, c \in U$, $\alpha \in C$, and $f(x_1, \dots, x_n)^2$ is central valued in $eRCe$; (iii) $\text{char}(R) = 2$ and $s_4(x_1, x_2, x_3, x_4)$ is an identity for $eRCe$.

In particular the following result holds:

Theorem A *Let R be a prime ring with extended centroid C , U its two-sided Utumi quotient ring, $\delta \neq 0$ a non-zero generalized derivation of R , $f(x_1, \dots, x_n)$ a non-central multilinear polynomial over C in n non-commuting variables, such that $[\delta(f(r_1, \dots, r_n)), f(r_1, \dots, r_n)] = 0$, for all $r_1, \dots, r_n \in R$. Then one of the following holds:*

1. $f(x_1, \dots, x_n)$ is central valued in R ;
2. there exists $\lambda \in C$ such that $\delta(x) = \lambda x$, for all $x \in R$;
3. there exist $q \in U$ and $\lambda \in C$ such that $\delta(x) = (q + \lambda)x + xq$, for all $x \in R$, and $f(x_1, \dots, x_n)^2$ is central valued on R ;
4. $\text{char}(R) = 2$ and $s_4(x_1, x_2, x_3, x_4)$ is an identity for R .

Here we will continue the study of the set

$$S = \{[\delta(f(x_1, \dots, x_n)), f(x_1, \dots, x_n)], x_1, \dots, x_n \in R\}$$

for a generalized derivation δ of R . An approach that can be used in studying S is to examine its size and a reasonable criteria for studying the size of S is to examine its left annihilator $Ann_R(S) = \{x \in R, \quad xs = 0, \forall s \in S\}$: if S is large we would expect that $Ann_R(S) = 0$. In fact we will prove this theorem:

Theorem 1 *Let R be a prime ring of characteristic different from 2, with extended centroid C , U its two-sided Utumi quotient ring, $\delta \neq 0$ a non-zero generalized derivation of R , $f(x_1, \dots, x_n)$ a non-central multilinear polynomial over C in n non-commuting variables, $a \in R$ such that $a[\delta(f(r_1, \dots, r_n)), f(r_1, \dots, r_n)] = 0$, for all $r_1, \dots, r_n \in R$. Then one of the following holds:*

1. $a = 0$;
2. there exists $\lambda \in C$ such that $\delta(x) = \lambda x$, for all $x \in R$;
3. there exist $q \in U$ and $\lambda \in C$ such that $\delta(x) = (q + \lambda)x + xq$, for all $x \in R$, and $f(x_1, \dots, x_n)^2$ is central valued on R .

Of course we do not consider the case when R is a domain; in fact, in this case, either $Ann_R(S) = 0$ or $[\delta(f(r_1, \dots, r_n)), f(r_1, \dots, r_n)] = 0$, for all $r_1, \dots, r_n \in R$. In this condition we conclude by Theorem A.

We also would like to remark that in case δ is an usual derivation of R , the conclusion of Theorem 1 follows directly from the following result we proved in [6]:

Theorem B *Let R be a prime ring of characteristic different from 2 with extended centroid C , U its two-sided Utumi quotient ring, $d \neq 0$ a non-zero derivation of R , $f(x_1, \dots, x_n)$ a non-central multilinear polynomial over C in n non-commuting variables, $a \in R$ such that $a[d(f(r_1, \dots, r_n)), f(r_1, \dots, r_n)] = 0$, for any $r_1, \dots, r_n \in R$. Then $a = 0$.*

In [14], T. K. Lee extended the definition of a generalized derivation to the Utumi quotient ring U of R as follows: by a generalized derivation we mean an additive mapping $\delta : I \rightarrow U$ such that $\delta(xy) = \delta(x)y + xd(y)$, for all $x, y \in I$, where I is a dense right ideal of R and d is a derivation from I into U .

In all that follows let U be the two-sided Utumi quotient ring of R and $C = Z(U)$

the center of U , $T = U *_C C\{X\}$ the free product over C of the C -algebra U and the free C -algebra $C\{X\}$, with X the countable set consisting of non-commuting indeterminates $x_1, x_2, \dots, x_n, \dots$. We refer the reader to [1] for the definitions and the related properties of these objects.

Of course U is a prime ring and, by replacing R by U , we may assume, without loss of generality, $R = U$, $C = Z(R)$ and R is a C -algebra centrally closed. Moreover we will use the following notation:

$$f(x_1, \dots, x_n) = x_1 x_2 \dots x_n + \sum_{\sigma \in S_n, \sigma \neq id} \alpha_\sigma x_{\sigma(1)} x_{\sigma(2)} \dots x_{\sigma(n)}$$

for some $\alpha_\sigma \in C$. We also assume $char(R) \neq 2$ and $f(x_1, \dots, x_n)$ non-central valued.

2. The case of inner generalized derivations

In this section we consider the case when δ is an inner generalized derivation induced by the elements $b, c \in R$, that is $\delta(x) = bx + xc$, for all $x \in R$. In this sense, our aim will be to prove the following:

Proposition 1 *Let R be a prime ring of characteristic different from 2, $f(x_1, \dots, x_n)$ a non-central multilinear polynomial over C in n non-commuting variables, $a, b, c \in R$. If $a[bf(r_1, \dots, r_n) + f(r_1, \dots, r_n)c, f(r_1, \dots, r_n)] = 0$, for all $r_1, \dots, r_n \in R$, then one of the following holds:*

1. $a = 0$;
2. $b, c \in C$;
3. there exists $\lambda \in C$ such that $c - b = \lambda$ and $f(x_1, \dots, x_n)^2$ is central valued on R .

2.1. The matrix case

By first we will study the case when $R = M_m(F)$ is the algebra of $m \times m$ matrices over a field F of characteristic different from 2. Notice that the set $f(R) = \{f(r_1, \dots, r_n) : r_1, \dots, r_n \in R\}$ is invariant under the action of all inner automorphisms of R . Hence if denote we $r = (r_1, \dots, r_n) \in R \times R \times R \times \dots \times R = R^n$, then for any inner automorphism φ of $M_m(F)$, we have that $\underline{r} = (\varphi(r_1), \dots, \varphi(r_n)) \in R^n$ and $\varphi(f(r)) = f(\underline{r}) \in f(R)$. Let us denote as usual by e_{ij} the matrix unit with 1 in (i, j) -entry and zero elsewhere.

Since $f(x_1, \dots, x_n)$ is not central then, by [16], there exist $u_1, \dots, u_n \in M_t(F)$ and $\alpha \in F - \{0\}$, such that $f(u_1, \dots, u_n) = \alpha e_{kl}$, with $k \neq l$. Moreover, since the set $\{f(v_1, \dots, v_n) : v_1, \dots, v_n \in M_t(F)\}$ is invariant under the action of all F -automorphisms of $M_t(F)$, then for any $i \neq j$ there exist $r_1, \dots, r_n \in M_t(F)$ such that $f(r_1, \dots, r_n) = \alpha e_{ij}$.

We start by studying the case of 2×2 matrices:

Lemma 1 *Let $R = M_2(F)$ be the algebra of 2×2 matrices over the field F of characteristic different from 2. If there exist $a, b, c \in R$ such that $a[bu + uc, u] = 0$ for any $u \in f(R)$ then either $a = 0$ or $c - b$ is a diagonal matrix.*

Proof. Suppose $a \neq 0$. Let $u = \alpha e_{ij} \in f(R)$, for any $i \neq j$ and $\alpha \neq 0$:

$$0 = a[b\alpha e_{ij} + \alpha e_{ij}c, \alpha e_{ij}] = -\alpha^2 a e_{ij}(c - b)e_{ij}.$$

In other words, either the i -th column of the matrix a is zero or, for all j different from i , the (j, i) -entry γ_{ji} of $(c - b)$ is zero. Suppose that $(c - b)$ is not a diagonal matrix, say $\gamma_{12} \neq 0$. In this case, as we said above, the 2-nd column of a is zero. Of course we may assume $\gamma_{21} = 0$, otherwise the first column of a is zero too, and we are done. In other words we are in the situation:

$$c - b = \begin{bmatrix} \gamma_{11} & \gamma_{12} \\ 0 & \gamma_{22} \end{bmatrix} \quad \gamma_{12} \neq 0, \quad a = \begin{bmatrix} a_{11} & 0 \\ a_{21} & 0 \end{bmatrix}.$$

Let φ be an element of $Aut(M_2(F))$, defined as $\varphi(x) = (1 - e_{12})x(1 + e_{12})$. Now choose $v = (1 - e_{12})u(1 + e_{12}) = \begin{bmatrix} -\alpha & -\alpha \\ \alpha & \alpha \end{bmatrix} \in f(R)$. Hence

$$0 = a[bv + vc, v] = -\alpha^2 \begin{bmatrix} a_{11}(\gamma_{11} - \gamma_{12} - \gamma_{22}) & a_{11}(\gamma_{11} - \gamma_{12} - \gamma_{22}) \\ a_{21}(\gamma_{11} - \gamma_{12} - \gamma_{22}) & a_{21}(\gamma_{11} - \gamma_{12} - \gamma_{22}) \end{bmatrix},$$

which implies

$$\gamma_{11} - \gamma_{12} - \gamma_{22} = 0 \quad (1).$$

Let now χ be another element of $Aut(M_2(F))$, defined as $\chi(x) = (1 + e_{12})x(1 - e_{12})$ and choose $w = (1 + e_{12})u(1 - e_{12}) = \begin{bmatrix} \alpha & -\alpha \\ \alpha & -\alpha \end{bmatrix} \in f(R)$. Hence we have

$$0 = a[bw + wc, w] = \alpha^2 \begin{bmatrix} a_{11}(\gamma_{11} + \gamma_{12} - \gamma_{22}) & a_{11}(-\gamma_{11} - \gamma_{12} + \gamma_{22}) \\ a_{21}(\gamma_{11} + \gamma_{12} - \gamma_{22}) & a_{21}(-\gamma_{11} - \gamma_{12} + \gamma_{22}) \end{bmatrix},$$

that is

$$\gamma_{11} + \gamma_{12} - \gamma_{22} = 0 \quad (2).$$

By equations (1) and (2) and since $\text{char}(R) \neq 2$, it follows $\gamma_{12} = 0$, a contradiction. Therefore, if $a \neq 0$, $(c - b)$ is a diagonal matrix. \square

Lemma 2 *Let $R = M_t(F)$ be the algebra of $t \times t$ matrices over the field F of characteristic different from 2 and $t \geq 3$. If there exist $a, b, c \in R$ such that $a[bu + uc, u] = 0$ for any $u \in f(R)$ then either $a = 0$ or $c - b$ is a diagonal matrix.*

Proof. As above, for any $i \neq j$ there exist $r_1, \dots, r_n \in M_t(F)$ such that $f(r_1, \dots, r_n) = \alpha e_{ij}$, with $\alpha \neq 0$. Hence, for all $i \neq j$,

$$0 = a[bf(r_1, \dots, r_n) + f(r_1, \dots, r_n)c, f(r_1, \dots, r_n)] = 0 = -\alpha^2 a e_{ij} (c - b) e_{ij}.$$

In other words, since $\text{char}(R) \neq 2$ and $\alpha \neq 0$, either the i -th column of the matrix a is zero or, for all j different from i , the (j, i) -entry q_{ji} of $(c - b)$ is zero.

Suppose by contradiction that $(c - b)$ is not a diagonal matrix, then there exists some non-zero entry q_{ji} of $(c - b)$, for $i \neq j$. As we said above, the i -th column of a is zero. Let $m \neq i, j$ and $\varphi_{mi}(x) = (1 + e_{mi})x(1 - e_{mi})$. Consider the following valuations of $f(x_1, \dots, x_n)$:

$$f(\underline{r}) = \gamma e_{ij}, \quad f(\underline{s}) = \varphi_{mi}(f(\underline{r})) = \gamma e_{ij} + \gamma e_{mj}, \quad \gamma \neq 0.$$

Since $f(\underline{s})^2 = 0$ we have $0 = a[bf(\underline{s}) + f(\underline{s})c, f(\underline{s})] = \gamma^2 a(e_{ij} + e_{mj})(c - b)(e_{ij} + e_{mj})$. Moreover, since the i -th column of a is zero, we obtain $\gamma^2 a(q_{ji} + q_{jm})e_{mj} = 0$. Notice that if $q_{ji} + q_{jm} = 0$, then $q_{jm} = -q_{ji} \neq 0$, so the m -th column of a is zero. On the other hand, if $q_{ji} + q_{jm} \neq 0$, it follows again that the m -th column of a is zero. Hence we can say that a has at most one non-zero column, the j -th one.

Let now ψ any F -automorphism of $M_t(F)$, then

$$0 = \psi(a)[\psi(b)\psi(f(r_1, \dots, r_n)) + \psi(f(r_1, \dots, r_n))\psi(c), \psi(f(r_1, \dots, r_n))] = \\ \psi(a)[\psi(b)f(s_1, \dots, s_n) + f(s_1, \dots, s_n)\psi(c), f(s_1, \dots, s_n)]$$

for all $s_1, \dots, s_n \in M_t(F)$. Therefore, as above, we can conclude that, if the (j, i) -entry of $\psi(c - b)$ is non-zero, for some $j \neq i$, then $\psi(a)$ has at most one non-zero column, the j -th one.

Let now $\psi(x) = (1 + e_{jm})x(1 - e_{jm})$, with $m \neq j, i$. Hence $\psi(c - b) = (c - b) + e_{jm}(c - b) - (c - b)e_{jm} - e_{jm}(c - b)e_{jm}$ and so its (j, i) -entry is $q_{ji} + q_{mi}$.

If $q_{ji} + q_{mi} = 0$ then $q_{ji} = -q_{mi} \neq 0$, that is the (m, i) -entry of $(c - b)$ is non-zero. In this case a has at most one non-zero column, the m -th one; but $m \neq j$ and so any column of a is zero.

If $q_{ji} + q_{mi} \neq 0$ then the (j, i) -entry of $\psi(c - b)$ is non-zero, hence $\psi(a)$ has at most one non-zero column, the j -th one.

Since $\psi(a) = (\sum_h a_{hj}e_{hj} + a_{mj}e_{jj}) - (\sum_h a_{hj}e_{hm} + a_{mj}e_{jm})$, then, for any $h \neq j$ must be $a_{hj} = 0$ and also $a_{jj} + a_{mj} = 0$. But in this situation we get $a = 0$. Therefore, if $a \neq 0$, then $q_{ji} = 0$, for all $j \neq i$. \square

Lemma 3 *Let $R = M_t(F)$ be the algebra of $t \times t$ matrices over the field F of characteristic different from 2 and $t \geq 2$. If there exist $a, b, c \in R$ such that $a[bu + uc, u] = 0$ for any $u \in f(R)$ then one of the following holds:*

1. $a = 0$;
2. $b, c \in Z(R)$;
3. $c - b \in Z(R)$ and $u^2 \in Z(R)$, for all $u \in f(R)$.

Proof. Suppose $a \neq 0$. By previous two lemmas, $c - b$ is a diagonal matrix, say $c - b = \sum q_{kk}e_{kk}$. Moreover if φ is an automorphism of $M_t(F)$, the same conclusion holds for $\varphi(c - b)$, since as above

$$0 = \varphi(a)[\varphi(b)\varphi(f(r_1, \dots, r_n)) + \varphi(f(r_1, \dots, r_n))\varphi(c)] = \varphi(a)[\varphi(b)f(s_1, \dots, s_n) + f(s_1, \dots, s_n)\varphi(c), (f(s_1, \dots, s_n))]$$

and $\varphi(a) \neq 0$. Therefore, for any $i \neq j$, $\varphi(c - b) = (1 + e_{ij})(c - b)(1 - e_{ij})$ must be a diagonal matrix. Thus $(q_{jj} - q_{ii})e_{ij} = 0$, that is $q_{jj} = q_{ii}$ and $c - b$ is a central element, that is there exists an element $\lambda \in Z(R)$, such that $b = c + \lambda$. Thus we have that, for all $r_1, \dots, r_n \in R$,

$$0 = a[cf(r_1, \dots, r_n) + f(r_1, \dots, r_n)c, f(r_1, \dots, r_n)] = a[c, f(x_1, \dots, x_n)^2].$$

Let A be the additive subgroup generated by the polynomial $f(x_1, \dots, x_n)^2$. By [3], since $\text{char}(R) \neq 2$, either $f(x_1, \dots, x_n)^2$ is central valued on R , or there exists a non-central Lie

ideal L of R such that $L \subseteq A$. In this last case we get $a[c, u] = 0$ for all $u \in L$. Since R is simple, L is not central and $\text{char}(R) \neq 2$, it is well known that $0 \neq [R, R] \subseteq L$ (see pp 4-5 in [8], Lemma 2 and Proposition 1 in [7], Theorem 4 in [11]). It follows that $a[c, [x_1, x_2]] = 0$ for all $x_1, x_2 \in R$. As a consequence of Theorem 1 in [5], we get $c \in Z(R)$. Therefore either $f(x_1, \dots, x_n)^2$ is central on R , or $b, c \in Z(R)$. \square

2.2. The proof of Proposition 1

In order to continue our investigation, we need to fix some well known facts.

Remark 1 Recall that if B is a basis of U over C , then any element of $T = U *_C C\{x_1, \dots, x_n\}$ can be written in the form $g = \sum_i \alpha_i m_i$, where $\alpha_i \in C$ and m_i are B-monomials, that is $m_i = q_0 y_1 \cdots y_n q_n$, with $q_i \in B$ and $y_i \in \{x_1, \dots, x_n\}$. In [2] it is showed that a generalized polynomial $g = \sum_i \alpha_i m_i$ is the zero element of T if and only if any α_i is zero. As a consequence, if $a_1, a_2 \in U$ are linearly independent over C and $a_1 g_1(x_1, \dots, x_n) + a_2 g_2(x_1, \dots, x_n) = 0 \in T$, for some $g_1, g_2 \in T$, then both $g_1(x_1, \dots, x_n)$ and $g_2(x_1, \dots, x_n)$ are the zero element of T

Lemma 4 *If R does not satisfy any non-trivial generalized polynomial identity, then either $a = 0$ or b and c are elements of C .*

Proof. Suppose that either $b \notin C$ or $c \notin C$, if not we are done. Since R does not satisfy any non-trivial generalized polynomial identity, we have that

$$a[bf(x_1, \dots, x_n) + f(x_1, \dots, x_n)c, f(x_1, \dots, x_n)]$$

is the zero element in the free product $T = U *_C C\{x_1, \dots, x_n\}$, that is

$$a(bf(x_1, \dots, x_n)^2 - f(x_1, \dots, x_n)^2c + f(x_1, \dots, x_n)(c - b)f(x_1, \dots, x_n)) = 0 \in T.$$

Suppose a and ab are linearly independent over C . We have

$$abf(x_1, \dots, x_n)^2 - a(f(x_1, \dots, x_n)^2c + f(x_1, \dots, x_n)(c - b)f(x_1, \dots, x_n)) = 0 \in T.$$

By Remark 1, $abf(x_1, \dots, x_n)^2 = 0 \in T$. Since R does not satisfy any non-trivial generalized polynomial identity, this forces $ab = 0$, which is a contradiction.

Thus we assume a and ab are linearly C -dependent, that is $ab = \beta a$, with $\beta \in C$. In this case we have:

- if $\beta \neq 0$:

$$\beta af(x_1, \dots, x_n)^2 - af(x_1, \dots, x_n)^2c + af(x_1, \dots, x_n)(c - b)f(x_1, \dots, x_n) = 0 \in T;$$

- if $\beta = 0$:

$$-af(x_1, \dots, x_n)^2c + af(x_1, \dots, x_n)(c - b)f(x_1, \dots, x_n) = 0 \in T.$$

In any case, if a and $(c - b)$ are linearly independent over C , by Remark 1 we obtain that in particular $af(x_1, \dots, x_n)(c - b)f(x_1, \dots, x_n)$ is the zero element in T . Since $(c - b) \neq 0$, we get the required conclusion $a = 0$.

Finally, let a and $(c - b)$ be linearly dependent over C . If $(c - b) = 0$, it follows that

$$\beta af(x_1, \dots, x_n)^2 - af(x_1, \dots, x_n)^2c = 0 \in T,$$

in particular $af(x_1, \dots, x_n)^2c$ is the zero element in T . If $a \neq 0$, then $c = 0$, that is also $b = 0$, a contradiction.

In case $(c - b) \neq 0$ then, for some $0 \neq \gamma \in C$, we have

$$\begin{aligned} &\beta\gamma(c - b)f(x_1, \dots, x_n)^2 - \gamma(c - b)(f(x_1, \dots, x_n))^2c \\ &- \gamma(c - b)f(x_1, \dots, x_n)(c - b)f(x_1, \dots, x_n) = 0 \in T \end{aligned}$$

in particular $\gamma(c - b)(f(x_1, \dots, x_n))^2c = 0 \in T$, which implies $c = 0$. By our assumption it follows that $b \notin C$, which implies the contradiction that

$$-\beta\gamma bf(x_1, \dots, x_n)^2 - \gamma bf(x_1, \dots, x_n)bf(x_1, \dots, x_n)$$

is a non-trivial generalized polynomial identity for R . □

In order to prove the next Lemma, here we premit the following:

Fact 1 *Let R be a dense ring of linear transformations over an infinite dimensional right vector space V over a division ring D . Then for any linearly D -independent subset $\{v_1, \dots, v_k\}$ of V , and for any subset $\{w_1, \dots, w_k\}$ of V , there exist $r_1, \dots, r_n \in R$ such that $f(r_1, \dots, r_n)v_i = w_i$, for all $i = 1, \dots, k$.*

Proof. Let $U_1 = \{v_1, \dots, v_k\} \subseteq V$ be a subset of linearly D -independent vectors. Since $\dim_D V = \infty$, there exists a set

$$U_2 = \{v_{ij}; \quad i = 1, \dots, k; \quad j = 1, \dots, n - 1\}$$

of linearly D -independent vectors such that $U_1 \cup U_2$ is also a linearly D -independent subset of V . In other words, if denote $v_i = v_{in}$, for all $i = 1, \dots, k$, we have that

$$\{v_{ij}; \quad i = 1, \dots, k; \quad j = 1, \dots, n\}$$

is a subset of independent vectors in V . By the density of R , there exist $r_1, \dots, r_n \in R$ such that

$$\begin{aligned} r_1 v_{i1} &= w_i, & r_i v_{ji} &= v_{j,i-1}, & i &= 2, \dots, n; & j &= 1, \dots, k \\ r_i v_{jl} &= 0, & & & & \text{for } l \neq i. \end{aligned}$$

Easy calculations show that $f(r_1, \dots, r_n)v_i = w_i$, for all $i = 1, \dots, k$. □

Lemma 5 *Let R be a dense ring of linear transformations over an infinite dimensional right vector space V over a division ring D . Then either $a = 0$ or b and c are central elements.*

Proof. Suppose that a is non-zero.

Our first aim is to show that for any $v \in V$, if $av \neq 0$ then v, cv are linearly D -dependent.

By contradiction let v, cv be D -independent. There exist $w, w_1, \dots, w_{n-1}, v_1, \dots, v_{n-1} \in V$ such that $v, cv = u, w, w_1, \dots, w_{n-1}, v_1, \dots, v_{n-1}$ are linearly independent. By Fact 1, there exist $r_1, \dots, r_n \in R$ such that

$$f(r_1, \dots, r_n)v = 0, \quad f(r_1, \dots, r_n)u = w, \quad f(r_1, \dots, r_n)w = v.$$

Hence, if av is non-zero, then we get the contradiction

$$\begin{aligned} 0 &= a[bf(r_1, \dots, r_n) + f(r_1, \dots, r_n)c, f(r_1, \dots, r_n)]v = \\ &= -af(r_1, \dots, r_n)^2u = -av \neq 0. \end{aligned}$$

Now suppose $av = 0$.

Since $a \neq 0$, there exists $w \in V$ such that $aw \neq 0$. Hence $a(w - v) = aw \neq 0$. By the previous argument we have that w, cw are linearly D -dependent and $(w - v), c(w - v)$ is as well.

Thus there exist $\alpha, \beta \in D$ such that $cw = w\alpha$ and $c(w - v) = (w - v)\beta$. Moreover v, w are linearly independent.

Suppose first that there exist $\lambda, \mu \in D$ such that $b(w - v) = w\lambda + v\mu$. Since $\dim_D V = \infty$, there exist $w_3, \dots, w_{n-1} \in V$ such that $v, w, w_3, \dots, w_{n-1}$ are linearly independent and by Fact 1 there exist $s_1, \dots, s_n \in R$ such that

$$f(s_1, \dots, s_n)v = 0, \quad f(s_1, \dots, s_n)w = w - v, \quad f(s_1, \dots, s_n)^2w = w - v$$

and

$$0 = a(bf(s_1, \dots, s_n)^2 - f(s_1, \dots, s_n)^2c + f(s_1, \dots, s_n)(c - b)f(s_1, \dots, s_n))w =$$

$$a(w\lambda + v\mu + (w - v)(\beta - \alpha) - (w - v)\lambda) = aw(\beta - \alpha).$$

Suppose now that $\{b(w - v), w, v\}$ are linearly D-independent. Hence there exist $w_3, \dots, w_{n-1}, u_1, \dots, u_{n-1} \in V$ such that $b(w - v), v, w, w_3, \dots, w_{n-1}, u_1, \dots, u_{n-1}$ are linearly independent and $t_1, \dots, t_n \in R$ such that

$$f(t_1, \dots, t_n)v = 0, \quad f(t_1, \dots, t_n)w = w - v, \quad f(t_1, \dots, t_n)^2w = w - v$$

and in this case we also have

$$f(t_1, \dots, t_n)b(w - v) = b(w - v)$$

which implies that

$$0 = a(bf(t_1, \dots, t_n)^2 - f(t_1, \dots, t_n)^2c + f(t_1, \dots, t_n)(c - b)f(t_1, \dots, t_n))w =$$

$$a(b(w - v) + (w - v)(\beta - \alpha) - b(w - v)) = aw(\beta - \alpha).$$

In any case, we have that $aw(\beta - \alpha) = 0$. Because $aw \neq 0$ then $\alpha = \beta$ and $cv = v\alpha$. This means that for any choice of $v \in V$, v, cv are linearly D-dependent. Standard arguments prove that there exists $\beta \in D$ such that $cv = v\beta$, for all $v \in V$ and also, by using this fact, that $c \in Z(R)$. Therefore for all $r_1, \dots, r_n \in R$

$$abf(r_1, \dots, r_n)^2 - af(r_1, \dots, r_n)bf(r_1, \dots, r_n) = 0.$$

Our final aim is to prove that $b \in Z(R)$. To do this we repeat the same above argument: we want to show that for any $v \in V$ then v, bv are linearly D-dependent.

Suppose by contradiction that v, bv are D-independent. There exist $w, w_1, \dots, w_{n-1}, v_1, \dots, v_{n-1} \in V$ such that $v, bv, w, w_1, \dots, w_{n-1}, v_1, \dots, v_{n-1}$ are linearly independent. By Fact 1, there exist $r_1, \dots, r_n \in R$ such that

$$f(r_1, \dots, r_n)v = 0, \quad f(r_1, \dots, r_n)u = v, \quad f(r_1, \dots, r_n)w = v.$$

Hence, for $av \neq 0$, we get the contradiction

$$0 = a[bf(r_1, \dots, r_n), f(r_1, \dots, r_n)]w =$$

$$abf(r_1, \dots, r_n)^2w - af(r_1, \dots, r_n)bf(r_1, \dots, r_n)w = -av \neq 0.$$

Let now $av = 0$. As above there exists $w \in V$ such that $aw \neq 0$. Hence $a(w - v) = aw \neq 0$. Again it follows that w, bw are linearly D -dependent and is also the case for $(w - v), b(w - v)$.

Thus there exist $\alpha', \beta' \in D$ such that $bw = w\alpha'$ and $b(w - v) = (w - v)\beta'$. Moreover v, w are linearly independent. Moreover there exist $w_3, \dots, w_{n-1} \in V$ such that $v, w, w_3, \dots, w_{n-1}$ are linearly independent and again by Fact 1 there exist $s_1, \dots, s_n \in R$ such that

$$f(s_1, \dots, s_n)v = 0, \quad f(s_1, \dots, s_n)w = w - v, \quad f(s_1, \dots, s_n)^2w = w - v$$

and

$$0 = a(bf(s_1, \dots, s_n)^2 - f(s_1, \dots, s_n)bf(s_1, \dots, s_n))w = a(w\lambda + v\mu + (w - v)(\beta - \alpha)).$$

Since $aw \neq 0$, one has $\alpha = \beta$ and $qv = v\alpha$.

Therefore in any case v, bv are linearly D -dependent, and $b \in Z(R)$. □

2.3. Proof of Proposition 1

As we said above, we assume $C = Z(R)$ and R is a C -algebra centrally closed, that is $R = RC$. If R does not satisfy any non-trivial generalized polynomial identity then, by lemma 4, we are done. Thus we may suppose that R satisfies a non-trivial generalized polynomial identity. By Martindale's theorem in [17], R is a primitive ring which is isomorphic to a dense ring of linear transformations of a vector space V over a division ring D . If $\dim_D V = \infty$, then, by lemma 5, we get the required conclusion.

Therefore we consider the case $\dim_D(V) = k$, with k finite positive integer. Of course $k \geq 2$, because R is not a domain. In this condition R is a simple ring which satisfies a non-trivial generalized polynomial identity. By lemma 2 in [10] (see also theorem 2.3.29 in [19]), $R \subseteq M_t(F)$, for a suitable field F and $t \geq 2$, moreover $M_t(F)$ satisfies the same generalized identity of R , hence $a[bf(r_1, \dots, r_n) + f(r_1, \dots, r_n)c, f(r_1, \dots, r_n)] = 0$, for all $r_1, \dots, r_n \in M_t(F)$ and moreover $f(x_1, \dots, x_n)$ is a non-central polynomial for $M_t(F)$. In this case we are done by lemma 3. □

3. The proof of the Theorem

In [14] Lee proved that every generalized derivation can be uniquely extended to a generalized derivation of U and thus all generalized derivations of R will be implicitly assumed to be defined on the whole U and obtained the following result: (Theorem 3 in

[14]) Every generalized derivation δ on a dense right ideal of R can be uniquely extended to U and assumes the form $\delta(x) = bx + d(x)$, for some $b \in U$ and a derivation d on U .

In this section we denote by $f^d(x_1, \dots, x_n)$ the polynomial obtained from $f(x_1, \dots, x_n)$ by replacing each coefficient α_σ with $d(\alpha_\sigma)$. Thus we write $d(f(r_1, \dots, r_n)) = f^d(r_1, \dots, r_n) + \sum_i f(r_1, \dots, d(r_i), \dots, r_n)$, for all r_1, r_2, \dots, r_n in R .

In light of this, we finally prove the main result:

Theorem 1 *Let R be a prime ring of characteristic different from 2, C its extended centroid, U its two-sided Utumi quotient ring, $\delta \neq 0$ a non-zero generalized derivation of R , $f(x_1, \dots, x_n)$ a non-central multilinear polynomial over C in n non-commuting variables, $a \in R$ such that*

$$a[\delta(f(r_1, \dots, r_n)), f(r_1, \dots, r_n)] = 0$$

for any $r_1, \dots, r_n \in R$. Then one of the following holds:

1. $a = 0$;
2. there exists $\lambda \in C$ such that $\delta(x) = \lambda x$, for all $x \in R$;
3. there exist $q \in U$ and $\lambda \in C$ such that $\delta(x) = (q + \lambda)x + xq$, for all $x \in R$, and $f(x_1, \dots, x_n)^2$ is central valued on R .

Proof. Suppose by contradiction that $a \neq 0$. Since R satisfies the generalized differential identity

$$a[\delta(f(x_1, \dots, x_n)), f(x_1, \dots, x_n)],$$

the above cited Lee's result says that R satisfies

$$a[bf(x_1, \dots, x_n) + d(f(x_1, \dots, x_n)), f(x_1, \dots, x_n)] \tag{3}$$

If d is an inner derivation induced by an element $c \in U$, then R satisfies the generalized polynomial identity

$$a([bf(x_1, \dots, x_n) + cf(x_1, \dots, x_n) - f(x_1, \dots, x_n)c, f(x_1, \dots, x_n)])$$

which is

$$a([(b+c)f(x_1, \dots, x_n) - f(x_1, \dots, x_n)c, f(x_1, \dots, x_n)]).$$

In this case we are done by proposition 1.

Hence let d be an outer derivation of R . In this case R satisfies the differential identity

$$a[\delta(f(x_1, \dots, x_n)), f(x_1, \dots, x_n)] =$$

$$a \left([bf(x_1, \dots, x_n) + f^d(x_1, \dots, x_n) + \sum_i f(x_1, \dots, d(x_i), \dots, x_n), f(x_1, \dots, x_n)] \right).$$

By Kharchenko's theorem (see [9] and [15]), R satisfies the generalized polynomial identity

$$a[bf(x_1, \dots, x_n) + f^d(x_1, \dots, x_n) + \sum_i f(x_1, \dots, y_i, \dots, x_n), f(x_1, \dots, x_n)],$$

and in particular R satisfies the blended component

$$a[bf(x_1, \dots, x_n), f(x_1, \dots, x_n)].$$

Applying Proposition 1, since $a \neq 0$ and $f(x_1, \dots, x_n)$ is not central, one has that $b \in C$. Therefore, by (3), R satisfies the differential identity

$$a[d(f(x_1, \dots, x_n)), f(x_1, \dots, x_n)].$$

In this situation, by Theorem B, we conclude that $d = 0$ and $\delta(x) = bx$, with $b \in C$, for all $x \in R$. \square

Acknowledgement

The author wishes to thank the referee; by following her/his valuable suggestions, he has been able to shorten the proof of Lemma 5.

References

- [1] Beidar, K.I., Martindale, III W.S. and Mikhalev, A.V.: Rings with generalized identities, Pure and Applied Math., Dekker, New York, 1996.
- [2] Chuang, C.L.: GPIs having coefficients in Utumi quotient rings, Proc. Amer. Mat. Soc. vol.103 No.3, 723-728, 1988.
- [3] Chuang, C.L.: The additive subgroup generated by a polynomial, Israel J. Math. 59(1), 98-106, 1987.
- [4] De Filippis, V.: Engel conditions with generalized derivations on multilinear polynomials, Israel J. Math., to appear.
- [5] De Filippis, V.: Prime rings with annihilator conditions on power values of derivations on multilinear polynomials, Taiwanese Journal Math., vol.5 (4), 725-730, 2001.

- [6] De Filippis, V. and Di Vincenzo O.M.: Posner's second theorem and an annihilator condition, *Math. Pannonica* vol.12/1, 2001, 69-81.
- [7] Di Vincenzo, O.M.: On the n-th centralizer of a Lie ideal, *Boll. UMI (7) 3-A*, 77-85, 1989.
- [8] Herstein, I.N.: *Topics in ring theory*, Univ. of Chicago Press, Chicago 1969.
- [9] Kharchenko, V.K.: Differential identities of prime rings, *Algebra and Logic* 17, 155-168, 1978.
- [10] Lanski C.: An Engel condition with derivation, *Proc. Amer. Math. Soc.*, vol.118 No.3, 731-734, 1993.
- [11] Lanski, C. and Montgomery, S.: Lie structure of prime rings of characteristic 2, *Pacific Journal of Math.*, vol. 42, n.1, 117-135, 1972.
- [12] Lee, P.H. and Lee, T.K.: Derivations with Engel conditions on multilinear polynomials, *Proc. Amer. Math. Soc.* 124, No.9, 2625-2629, 1996.
- [13] Lee, T.K.: Derivations with Engel conditions on polynomials, *Algebra Colloq.* 5:1, 13-24, 1998.
- [14] Lee, T.K.: Generalized derivations of left faithful rings, *Comm. Algebra*, 27(8), 4057-4073, 1999.
- [15] Lee, T.K.: Semiprime ring with differential identities, *Bull. Inst. Math. Acad. Sinica*, vol.20 No.1, 27-38, 1992.
- [16] Leron, U.: Nil and power central polynomials in rings, *Trans. Amer. Math. Soc.* 202, 97-103, 1975.
- [17] Martindale, III W.S.: Prime rings satisfying a generalized polynomial identity, *J. Algebra* 12, 576-584, 1969.
- [18] Posner, E.C.: Derivations in prime rings, *Proc. Amer. Math. Soc.* 8, 1093-1100, 1957.
- [19] Rowen, L.: *Polynomial identities in Ring Theory*, Pure and Applied Math. vol. 84, Academic Press, New York 1980.

Vincenzo DE FILIPPIS
 DI.S.I.A. Faculty of Engineering
 University of Messina
 Contrada Di Dio, 98166 Messina-ITALY
 e-mail: defilippis@unime.it

Received 28.02.2007