

1-1-2005

Excellent and Convenient Procedures for Reduction of Benzene and Its Derivatives

ALİYE ALTUNDAŞ

ABDULLAH MENZEK

DEMET DEMİRCİ GÜLTEKİN

MELEK KARAKAYA

Follow this and additional works at: <https://journals.tubitak.gov.tr/chem>

 Part of the [Chemistry Commons](#)

Recommended Citation

ALTUNDAŞ, ALİYE; MENZEK, ABDULLAH; GÜLTEKİN, DEMET DEMİRCİ; and KARAKAYA, MELEK (2005) "Excellent and Convenient Procedures for Reduction of Benzene and Its Derivatives," *Turkish Journal of Chemistry*. Vol. 29: No. 5, Article 7. Available at: <https://journals.tubitak.gov.tr/chem/vol29/iss5/7>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Chemistry by an authorized editor of TÜBİTAK Academic Journals. For more information, please contact academic.publications@tubitak.gov.tr.

Excellent and Convenient Procedures for Reduction of Benzene and Its Derivatives

Aliye ALTUNDAŞ^{1*}, Abdullah MENZEK^{2*}, Demet DEMİRCİ GÜLTEKİN² and Melek KARAKAYA²

¹*Department of Chemistry, Faculty of Arts and Sciences, Gazi University, 06500 Ankara-TURKEY*

²*Department of Chemistry, Faculty of Arts and Sciences, Atatürk University, 25240 Erzurum-TURKEY*

e-mail: aaltundas@gazi.edu.tr; amenzek@atauni.edu.tr

Received 08.05.2005

Reductions of benzene, toluene, ortho-xylene, cumene, anisole and benzoic acid to corresponding nonconjugated dienes were realized at room temperature by gas NH₃ and Li. The reduced products were easily isolated because THF and dioxane were dissolved in the water. The yields and conversions of the reactions were 70-89% and 80-100%, respectively.

Key Words: Arenes, lithium, reductions, alcohols, protonations, ethers.

Introduction


Reduction reactions of aromatic compounds such as the Birch reduction and related reactions are important in synthetic organic chemistry and therefore these reactions have been investigated by many chemists¹⁻⁶. The reduction products, especially 1,4-cyclohexadiene or intermediates formed in these reactions, serve as starting materials for many target molecules⁷⁻¹³.

The reduction of aromatic rings by solutions of alkali metals in liquid ammonia was discovered by Wooster and Godfrey¹⁴, who reacted toluene with sodium in ammonia followed by the addition of water. They reported a “highly unsaturated liquid product”, which was not identified further. However, the real development of this reaction was to follow in the work by Birch¹⁵. This reaction is generally referred to as the Birch reduction, although in some cases it is simply called metal-ammonia reduction. Wild and Nelson¹⁶ found adding alcohol last to be advantageous, as opposed to having it present when the metal is added, and it was subsequently discovered that it should be avoided altogether with polynuclear compounds.

Among all aromatic compounds, benzene is the first to come to mind. The reduction of benzene to 1,4-cyclohexadiene is important. The first observation of crucial importance in this field was Wooster's observation^{5,17,18}. Wooster showed that the presence of alcohol in liquid ammonia permitted the reduction of benzene to 1,4-cyclohexadiene. Other reduction reactions of benzene by alkali metal in liquid ammonia

*Corresponding authors

were realized in different conditions^{3,6,19–22}. Dumanski and Swerew²³ reduced benzene with metallic calcium into 1,4-cyclohexadiene at room temperature and gas ammonia. Preparation of 1,4-cyclohexadiene by this method does not seem suitable as a preparative method because its yield is very low¹⁸. Benzene and its derivatives were reduced in low molecular weight amines and ethylenediamine by Li^{24–27}.



Scheme 1

We realized the reduction of naphthalene and anthracene with sodium and tert-butyl alcohol (t-BuOH) to 1,4-dihydronaphthalene and 9,10-dihydroanthracene respectively in high yields at room temperature (r.t.) as the sole products (Scheme 1)²⁸. We investigated in this study a convenient Birch reduction procedure with mild conditions and high yields for reductions of benzene and its derivatives because reduction products are important in organic synthesis^{7–13}.

Experimental

General Method: All chemicals and solvents are commercially available and were used after distillation or treatment with drying agents. The ¹H- and ¹³C- NMR spectra were recorded on a 200 (50)-MHz Varian spectrometer; δ in ppm, with Me₄Si as the internal standard.

Procedure 1: Reduction of benzene to 1,4-cyclohexadiene.

In a 500-mL, 2-necked, round-bottomed flask fitted with a reflux condenser and a stirring bar were placed tert-butanol (72 g, 0.963 mol, 2.85 equivalents), dry benzene (27.3 g, 0.35 mol, 1 equivalent) and dry THF (120 mL). The flask was attached to gas ammonia (NH₃) whose pressure was approximately 1 atmosphere (atm) and the resulting solution was stirred. The reaction mixture was cooled in an ice-water bath and then freshly cut lithium (7.35 g, 1.05 g-atom, 3 equivalents) was added over 3-5 min. The temperature of the bath was allowed to rise gradually to room temperature. After the addition of lithium was completed, the reaction mixture was stirred for 5 h. Two phases appeared in the reaction mixture. The top and bottom phases were brown and gray, respectively. The mixture was cooled in an ice-water bath again. Cold water was added slowly and carefully to the flask until all the lithium was consumed, as evidenced by the conversion of the colors to white. The reaction mixture was poured into a mixture (250 g) of water and ice and was acidified with the addition of 2 N cold hydrochloric acid. The organic layer was separated and washed with cold water (100 mL), a solution of NaHCO₃ (5%, 50 mL) and water (75 mL), in order that. The reduction product, 1,4-cyclohexadiene, was dried over CaCl₂ and filtered. The yield (23.5 g) and conversion of the reaction were 84% and 100%, respectively.

Procedure 2: Reduction of benzene to 1,4-cyclohexadiene. (This experimental procedure may also be used if gas NH₃ is not continuously present in any laboratory.)

In a 500-mL, 2-necked, round-bottomed flask fitted with a reflux condenser, a stirrer bar and dropping funnel was placed dry dioxane (100 mL). The flask was attached to a balloon, of gas ammonia (NH₃), at 1 atmosphere (atm) and then washed with NH₃ while the dioxane was stirred. After stirring in an atmosphere of NH₃ for 15-20 min, freshly cut lithium (8.0 g, 1.152 g-atom, 3 equivalents) was added in quantities for

5 min, followed by stirring for 15-20 min. Dry benzene (30 g, 385 mmol, 1 equivalent) was added and the mixture was stirred for 30 min. The reaction mixture was cooled in an ice-water bath and then a solution of *t*-BuOH (85 g, 1.152 g-atom, 3 equivalents) in dioxane (10 mL) was added slowly by dropping funnel. To drop the *t*-BuOH well, the stopcock of NH₃ may be switched on and off. The heat of the reaction mixture has to be checked. NH₃ was put in the balloon again and attached to the flask. When the addition of *t*-BuOH was complete, the reaction mixture was stirred for 2 h, and then the cold bath was removed. This mixture was stirred for an additional 20 h and then cooled by an ice-water bath. A mixture of water and ice was then added slowly until all the lithium was consumed, as evidenced by the conversion of the brown color to white-gray. A saturated solution (100 mL) of NH₄Cl was added, and the organic phase was separated and then washed with a saturated solution (2 x 30 mL) of NH₄Cl and water (2 x 30 mL), in order that. The reduction product, 1,4-cyclohexadiene, was dried over CaCl₂ and filtered. The yields and conversions of the reactions were 80-90% and 100%, respectively.

Reduction of toluene to 1-methyl-cyclohexa-1,4-diene: (Entry 2),

The reaction was realized according to procedure 1. Toluene (27.3 g, 0.3 mol), HOBu^t (63 g, 0.85 mol), THF (100 mL) and Li (6.3 g, 0.9 mol) were used in the reaction. The reaction lasted 7 h. Pure 1-methyl-cyclohexa-1,4-diene (24.2 g, 87%) was obtained.

Reduction of cumene to 1-isopropyl-cyclohexa-1,4-diene: (Entry 3),

The reaction was realized according to procedure 1. Cumene (16.4 g, 0.24 mol), HOBu^t (60 g, 0.81 mol), THF (100 mL) and Li (6.3 g, 0.9 mol) were used in the reaction. The reaction lasted 20 h. 1-Isopropyl-cyclohexa-1,4-diene and cumene were obtained in a mixture (25.3 g). Percentages of 1-isopropyl-cyclohexa-1,4-diene and cumene in the mixture were 80 and 20, respectively.

Reduction of ortho-xylene to 1,2-dimethyl-cyclohexa-1,4-diene: (Entry 4),

The reaction was realized according to procedure 1. Ortho-xylene (21.2 g, 0.2 mol), HOBu^t (45 g, 0.6 mol), THF (80 mL) and Li (5.6 g, 0.8 mol) were used in the reaction. The reaction lasted 24 h. Pure 1,2-dimethyl-cyclohexa-1,4-diene (17.06 g, 79%) was obtained.

Reduction of anisole to 1-methoxy-cyclohexa-1,4-diene: (Entry 5),


The reaction was realized according to procedure 1. Anisole (16.2 g, 0.15 mol), HOBu^t (33.3 g, 0.45 mol), THF (60 mL) and Li (3.5 g, 0.5 mol) were used in the reaction. The reaction lasted 24 h. The solution of cold hydrochloride acid was not added to the reaction mixture. The mixture was extracted with ether (3 x 75 mL). The combined organic layer was dried over Na₂SO₄ and the solvent was evaporated. Pure 1-methoxy-cyclohexa-1,4-diene (14 g, 85%) was obtained.

Reduction of benzoic acid to cyclohexa-2,5-dienecarboxylic acid: (Entry 6),

The reaction was realized according to procedure 1. Benzoic acid (4.0 g, 32.78 mmol), THF (50 mL) and Li (1.15 g, 164.29 mmol) were used in the reaction. As mentioned in the literature²⁹, alcohol was not used for this procedure as a proton source. The reaction lasted 8 h. The mixture was extracted with CHCl₃ (3 x 25 mL). The combined organic layer was dried over Na₂SO₄ and the solvent was evaporated. Benzoic acid and cyclohexa-2,5-dienecarboxylic acid were obtained in a mixture (3.75 g). Percentages of benzoic acid and cyclohexa-2,5-dienecarboxylic acid in the mixture were 17 and 83, respectively.














Results and Discussion

We continued our investigation of the reduction of aromatic compounds in different conditions and reduced benzene and its derivatives with lithium (Li) and alcohol (ROH) in tetrahydrofuran (THF) or dioxane at room temperature and under ammonia atmosphere (Scheme 2 and Table). The pressure of gas ammonia is approximately atmospheric pressure. In these procedures, aromatic compounds in solvent are reacted with Li in the presence of gas ammonia, and then the reaction mixture is quenched by HOEt, water or a saturated solution of NH_4Cl . The reaction mixtures may be cooled to 0 °C. Both procedures were used for the reduction of benzene. Only procedure 1 was used for the reduction of benzene derivatives. As shown, procedure 1 is better than procedure 2.



Scheme 2

Table. Reactants, products, yields and conversions in the reductions (procedure 1).

Entry	Reactant	Product	Yield (%)	Conversion (%)
1			84	100
2			87	100
3			70	80
4			79	100
5			85	100
6			76	83
7		-	0	0

Phenol (Table) was not reduced by procedure 1, while the others were easily reduced. Relative rates of reduction of aromatic rings depend on the electron density of rings. Electron donating groups on the aromatic rings decrease the rate of the reaction. The electron density of ring in phenol is much greater than the others because phenol converts into phenoxide anion in the reaction mixture. However, regiochemistry in reduction reactions is affected by the type of groups present on the ring because they tend to stabilize or destabilize the radical anion². Known regiochemistry was observed in these reactions (Table).

Compared to the other reduction procedures of converting benzene and its derivatives to corresponding nonconjugated dienes, the new reduction procedures have certain advantages. These include the following: 1) These reduction reactions are carried out at room temperature, avoiding the low temperatures, under -33°C, needed to obtain liquid NH₃. 2) The procedures are environmentally friendly³⁰⁻³⁴. Much more NH₃ is needed when liquid NH₃ is used as solvent. Evaporation of liquid NH₃ may damage the environment. 3) Control of moisture is easier in our method, and the reaction may go on for longer periods. When the reaction is conducted with liquid ammonia, it may be quenched by the developing moisture. 4) Evaporation of liquid NH₃ may take a long time and some side reactions such as isomerization and reoxidation may be observed. 5) It is unnecessary for the researcher to observe the reaction carefully and continuously in the present method because temperature control is not necessary, whereas the temperature must be checked in the reaction with liquid NH₃.

Conclusion

We have found new, excellent and convenient procedures for the reduction of benzene and its derivatives to corresponding nonconjugated dienes. The structure and purities of the reduced products were determined by NMR spectroscopy. The procedure is environmentally friendly. The reduced products were easily isolated because dioxane and THF were dissolved in the water. The yields and conversions of the reactions were 70-89% and 80-100%, respectively. Further reactions of aromatic compounds with metals (Na, Li etc.) are under investigation.

Acknowledgments

The authors are indebted to the Department of Chemistry (Atatürk University) and TÜBİTAK (Project No: 104T354) for financial support and to Dr. Ebru Mete for technical assistance.

References

1. H.O. House, **Modern Synthetic Reactions**; 2nd ed., W. A Benjamin, Los Angeles, CA, pp 1-227, 1972.
2. P.W. Rabideau, **Tetrahedron** **45**, 1579-1603 (1989).
3. M. Hudlicky, **Reduction in Organic Chemistry**; 2nd ed., ACS Monograph, 188, pp 61-79, 1996.
4. A. Menzek, **J. Chem. Educ.** **79**, 700-702 (2002).
5. A.J. Birch and D. Nasipuri, **Tetrahedron** **6**, 148-153, (1959).
6. R.G. Harvey, **Synthesis** 161-172 (1970).

7. J.M. Hook and L.N. Mander, **Natural Prod. Rep.** **3**, 35-85 (1986).
8. M. Balcı, Y. Sütbeyaz, and H. Seçen, **Tetrahedron** **46**, 3715-3742 (1990).
9. K.A. Powell, A.L. Hughes, H. Katchian, J.F. Jerauld and H.Z. Sable, **Tetrahedron** **28**, 2019-2027 (1972).
10. M. Mittelbach, N. Poklukar and H. Junek, **Liebigs Ann. Chem.** **2**, 185-188 (1990).
11. S. Spange, **Prog. Polym. Sci.** **25**, 781-849 (2000).
12. R.J. Giguere, A.M. Namen, B.O. Lopez, A. Arepally, D.E. Ramos, G. Majetich and J. Defauw, **Tetrahedron Lett.** **28**, 6553-6556 (1987).
13. M.S. Gültekin, E. Salamcı and M. Balcı, **Carbohydr. Res.** **338**, 1615-1619 (2003).
14. C.B. Wooster and K.L. Godfrey, **J. Am. Chem. Soc.** **59**, 596-597 (1937).
15. A.J. Birch, **J. Chem. Soc.**, 430-446 (1944).
16. A.L. Wilds and N.A. Nelson, **J. Am. Chem. Soc.** **75**, 5360-5365 (1953).
17. C.B. Wooster, **U. S. Patent**, 2182242 (1937).
18. J.P. Wibaut and F.A. Haak, **Recueil** **67**, 85-90 (1986).
19. K.N. Campbell and J.P. McDermott, **J. Am. Chem. Soc.** **67**, 282-283 (1945).
20. A.P. Krapcho and A.A. Bothner, **J. Am. Chem. Soc.** **81**, 3658-3666 (1959).
21. L.H. Slaugh and J.H. Raley, **J. Org. Chem.** **32**, 369-371 (1967).
22. P.W. Rabideau and D.L. Huser, **J. Org. Chem.** **48**, 4266-4271 (1983).
23. A.W. Dumanski and A.W. Swerew, **Chem. Zentr.** **III**, 746 (1923).
24. R.A. Benkeser, R.E. Robinson, D.M. Sauve and O.H. Thomas, **J. Am. Chem. Soc.** **77**, 3230-3233 (1959).
25. A.J. Birch, H.Q. Smith, **Rev. Chem. Soc.** **7**, 17-33 (1958).
26. R.A. Benkeser, J.A. Laugal and A. Rappa, **Tetrahedron Lett.** **25**, 2089-2092 (1984).
27. M.E. Grast, L.J. Dolby, S. Esfandiari, N.A. Fedoruk, N.C. Chamberlain and A.A. Avey, **J. Org. Chem.** **65**, 7098-7104 (2000).
28. A. Menzek, A. Altundaş and D. Gültekin, **J. Chem. Res-S.**, 752-753 (2003).
29. M.E. Kuehne and B.F. Lambert, **Org. Synth. Coll. Vol.** **5**, 400-403 (1973).
30. W.E. Luttrell, **Chem. Health Safe.** **9(3)**, 30-31 (2002).
31. W. Bajonowski, **Int J Refrigeration** **8(1)**, 56-57 (1985).
32. G.J. O’Kane, **Anaesthesia** **38(2)**, 1208-1213 (1983).
33. P.J. Temple, D.S. Harper, R.G. Pearson and S.N. Linzon, **Environmental Pollution** (1970-1979), **20(4)**, 297-302 (1979).
34. P.G. Urben, M.J. Pitt and L.A. Battle, **Bretherick’s Handbook of Reactive Chemical Hazards**, 5nd ed.– Vol. 1, Butterworth Heimann Ltd, Oxford OX2 8DP, 1553-1554, 1995.