

1-1-2007

Higher Order Generalization of Positive Linear Operators Defined by a Class of Borel Measures

OKTAY DUMAN

Follow this and additional works at: <https://journals.tubitak.gov.tr/math>



Part of the [Mathematics Commons](#)

Recommended Citation

DUMAN, OKTAY (2007) "Higher Order Generalization of Positive Linear Operators Defined by a Class of Borel Measures," *Turkish Journal of Mathematics*: Vol. 31: No. 3, Article 7. Available at: <https://journals.tubitak.gov.tr/math/vol31/iss3/7>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Mathematics by an authorized editor of TÜBİTAK Academic Journals. For more information, please contact academic.publications@tubitak.gov.tr.

Higher Order Generalization of Positive Linear Operators Defined by a Class of Borel Measures

Oktaç Duman

Abstract

In the present paper, we introduce a sequence of linear operators, which is a higher order generalization of positive linear operators defined by a class of Borel measures studied in [2]. Then, using the concept of A -statistical convergence we obtain some approximation results which are stronger than the aspects of the classical approximation theory.

Key Words: Statistical convergence, A -statistical convergence, positive linear operators, regular matrices, the elements of the Lipschitz class, Korovkin-type approximation theorem.

1. Introduction

Let I be an arbitrary interval of the real line, and let $C(I)$ denote the linear space of all real-valued continuous functions on I . Assume that g is a non-negative increasing function on $[0, \infty)$ with $g(0) = 1$. If I is an unbounded interval, then we consider the following function space

$$C_g(I) = \left\{ f \in C(I) : \lim_{|y| \rightarrow \infty; (y \in I)} \frac{|f(y)|}{(g(|y|))^c} = 0 \text{ for any } c > 0 \right\}, \quad (1)$$

which was examined in [2], [3]. Here, we should remark that if $I = [a, +\infty)$ (or $I = (a, +\infty)$), then the item “ $|y| \rightarrow \infty; (y \in I)$ ” in the definition (1) reduces to “ $y \rightarrow +\infty$ ”; however, if $I = (-\infty, a]$ (or $I = (-\infty, a)$), then we have “ $y \rightarrow -\infty$ ”. On the other hand, if I is an bounded interval, then we will use the space $C(I)$ instead of $C_g(I)$.

2000 Mathematics Subject Classification: 41A25, 41A36.

Now, for each fixed $x \in I$, let $\{\mu_{n,x} : n \in \mathbb{N}\}$ be a collection of measures defined on (I, \mathcal{B}) , where \mathcal{B} is the sigma field of Borel measurable subsets of I . Assume that, for any $\delta > 0$, the condition

$$\sup_{n \in \mathbb{N}} \int_{I \setminus I_\delta} g(|y|) d\mu_{n,x}(y) < \infty \tag{2}$$

holds, where $I_\delta := [x - \delta, x + \delta] \cap I$. In the condition (2), the boundedness is pointwise with respect to x ; that is, it is bounded for each fixed $x \in I$. With this terminology, in [2], some approximation properties of the following positive linear operators defined on $C_g(I)$ were investigated:

$$L_n(f; x) = \int_I f(y) d\mu_{n,x}(y) \quad , \quad n \in \mathbb{N} \text{ and } f \in C_g(I). \tag{3}$$

Define the space $C_g^{[r]}(I)$ by

$$C_g^{[r]}(I) = \left\{ f : f^{(r)} \in C_g(I) \right\}, \quad (r = 0, 1, 2, \dots).$$

If $r = 0$, then observe that $C_g^{[0]}(I) = C_g(I)$. We now consider the r -th order generalization of the operators L_n defined by (3) as follows

$$L_n^{[r]}(f; x) = \sum_{k=0}^r \int_I f^{(k)}(y) \frac{(x-y)^k}{k!} d\mu_{n,x}(y), \tag{4}$$

where $f \in C_g^{[r]}(I)$, $(r = 0, 1, 2, \dots)$, $n \in \mathbb{N}$, and also the function g satisfy the condition (2). We note that this kind of generalization was also considered in [11]. It is easy to see that if $r = 0$, then we have

$$L_n^{[0]}(f; x) = L_n(f; x).$$

The main goal of the present paper is to investigate various approximation properties of the linear operators $L_n^{[r]}$ defined by (4) with the help of the concept of A -statistical convergence. Recently, it has been shown that regular (non-matrix) summability transformations are also quite effective on the approximation of positive linear operators (see [2], [3], [4], [5], [9]). Especially, using the concept of the A -statistical convergence, where

A is a non-negative regular matrix, instead of the ordinary convergence in the approximation theory gives us many advantages, since A statistical convergence is stronger than the usual convergence.

Before proceeding further, we recall the concept of A -statistical convergence.

Let $A := (a_{jn}), j, n \in \mathbb{N}$, be a non-negative regular summability, i.e. $\lim Ax = L$ whenever $\lim x = L$, where $Ax := ((Ax)_j)$ is called an A -transform of $x := (x_n)$ and is given by

$$(Ax)_j := \sum_{n=1}^{\infty} a_{jn}x_n,$$

provided that the series convergence for each $j \in \mathbb{N}$ (see [10]). Then a sequence $x := (x_n)$ is called A -statistical convergent to a number L if, for every $\varepsilon > 0$,

$$\lim_j \sum_{n:|x_n-L|\geq\varepsilon} a_{jn} = 0.$$

We denote this limit by $st_A - \lim x = L$ [7] (see also [12], [14]). If we take $A = C_1$, the Cesàro matrix of order one, then C_1 -statistical convergence is equivalent to statistical convergence [6], [8]. Also replacing the matrix A by the identity matrix, A -statistical convergence coincides with the ordinary convergence. Kolk [12] proved that A -statistical convergence is stronger than ordinary convergence in the case of which $\lim_j \max_n |a_{jn}| = 0$.

2. A -Statistical Approximation Properties

In this section using A -statistical convergence we investigate some approximation properties of the operators $L_n^{[r]}$ defined by (4).

We note that a function $f \in C(I)$ belongs to $Lip_M(\alpha)$, $0 < \alpha \leq 1$, provided

$$|f(y) - f(x)| \leq M |y - x|^\alpha \quad (x, y \in I \text{ and } M > 0). \tag{5}$$

Then we obtain the following result.

Theorem 2.1 *Let I be an arbitrary interval of the real line, and let r be a non-negative integer. Assume that*

$$\int_I d\mu_{n,x}(y) = 1 \quad (\text{for each } x \in I \text{ and } n \in \mathbb{N}). \tag{6}$$

Then for all $f \in C_g^{[r]}(I)$ such that $f^{(r)} \in Lip_M(\alpha)$, $0 < \alpha \leq 1$, and for each $x \in I$, we have

$$\left| L_n^{[r]}(f; x) - f(x) \right| \leq CL_n(|x - y|^{\alpha+r}; x)$$

where

$$C = \frac{M\alpha}{\alpha + r} \frac{B(\alpha, r)}{(r - 1)!}, \tag{7}$$

and $B(\alpha, r)$ is the beta function.

Proof. By (4) and (6), we get

$$f(x) - L_n^{[r]}(f; x) = \int_I \left\{ f(x) - \sum_{k=0}^r f^{(k)}(y) \frac{(x - y)^k}{k!} \right\} d\mu_{n,x}(y). \tag{8}$$

Applying the Taylor's formula (see [11]) we may write that

$$f(x) - \sum_{k=0}^r f^{(k)}(y) \frac{(x - y)^k}{k!} = \frac{(x - y)^r}{(r - 1)!} \int_0^1 (1 - t)^{r-1} [f^{(r)}(y + t(x - y)) - f^{(r)}(y)] dt. \tag{9}$$

Since $f^{(r)} \in Lip_M(\alpha)$, we get from (5) that

$$\left| f^{(r)}(y + t(x - y)) - f^{(r)}(y) \right| \leq Mt^\alpha |x - y|^\alpha. \tag{10}$$

Considering (10) in (9), and using the beta integral, we conclude that

$$\left| f(x) - \sum_{k=0}^r f^{(k)}(y) \frac{(x - y)^k}{k!} \right| \leq |x - y|^{\alpha+r} \frac{M\alpha}{\alpha + r} \frac{B(\alpha, r)}{(r - 1)!}. \tag{11}$$

So combining (11) with (8), we have

$$\begin{aligned} \left| f(x) - L_n^{[r]}(f; x) \right| &\leq \frac{M\alpha}{\alpha + r} \frac{B(\alpha, r)}{(r - 1)!} \int_I |x - y|^{\alpha+r} d\mu_{n,x}(y) \\ &= \frac{M\alpha}{\alpha + r} \frac{B(\alpha, r)}{(r - 1)!} L_n(|x - y|^{\alpha+r}; x), \end{aligned}$$

which gives the desired result. □

Theorem 2.2 *Let $A = (a_{jn})$ be a non-negative regular summability matrix, and let I be an arbitrary interval of the real line. Let $0 < \alpha \leq 1$ and let r be a non-negative integer. Assume that the condition (6) is satisfied. Assume further that $g : [0, \infty) \rightarrow \mathbb{R}$, $g(y) = e^y$ and $h_x : I \rightarrow \mathbb{R}$, $h_x(y) = |x - y|^{\alpha+r}$ for each fixed $x \in I$. If the condition*

$$st_A - \lim_n L_n(h_x, x) = 0 \tag{12}$$

holds, then for all $f \in C_g^{[r]}(I)$ such that $f^{(r)} \in Lip_M(\alpha)$, we have

$$st_A - \lim_n \left| L_n^{[r]}(f; x) - f(x) \right| = 0.$$

Proof. Let $x \in I$ be fixed. By the definitions of the functions g and h_x , observe that $h_x \in C_g(I)$. Now, for a given $\varepsilon > 0$, define the following sets:

$$U := \left\{ n \in \mathbb{N} : \left| L_n^{[r]}(f; x) - f(x) \right| \geq \varepsilon \right\}$$

and

$$V := \left\{ n \in \mathbb{N} : L_n(h_x; x) \geq \frac{\varepsilon}{C} \right\},$$

where the constant C is given by (7). So it follows from Theorem 2.1 that $U \subseteq V$. Therefore, we get, for all $j \in \mathbb{N}$, that

$$\sum_{n \in U} a_{jn} \leq \sum_{n \in V} a_{jn}. \tag{13}$$

Note the condition (12) implies $\lim_j \sum_{n \in V} a_{jn} = 0$. So, we conclude from (13) that

$\lim_j \sum_{n \in U} a_{jn} = 0$, whence the result. □

If we use the test functions $e_i(y) = y^i$, ($i = 0, 1, 2$), instead of (12) in Theorem 2.1, then we have the following approximation result via A -statistical convergence.

Theorem 2.3 *Under the conditions of Theorem 2.2, if*

$$st_A - \lim_n |L_n(e_i, x) - e_i(x)| = 0, \quad (i = 0, 1, 2), \tag{14}$$

then for all $f \in C_g^{[r]}(I)$ such that $f^{(r)} \in Lip_M(\alpha)$, we have

$$st_A - \lim_n \left| L_n^{[r]}(f; x) - f(x) \right| = 0.$$

Proof. We first note that, by the definition of g , the test functions e_i , ($i = 0, 1, 2$) belong to $C_g(I)$. So, by Theorem 1 in [2], the condition (14) yields that for all $h \in C_g(I)$

$$st_A - \lim_n |L_n(h, x) - h(x)| = 0. \tag{15}$$

In particular, take $h := h_x$, which is defined in Theorem 2.2. Since $h_x(x) = 0$, it follows from (15) that

$$st_A - \lim_n |L_n(h_x, x)| = 0,$$

which gives (12). Therefore the proof follows from Theorem 2.2. □

Corollary 2.4 *If I is closed and bounded interval of the real line, say $I = [a, b]$, and also*

$$st_A - \lim_n \|L_n(e_i, \cdot) - e_i\|_{C[a,b]} = 0, \quad (i = 0, 1, 2),$$

then for all $f \in C^{[r]}[a, b]$ such that $f^{(r)} \in Lip_M(\alpha)$ we have

$$st_A - \lim_n \left\| L_n^{[r]}(f, \cdot) - f \right\|_{C[a,b]} = 0,$$

where $\|\cdot\|_{C[a,b]}$ denotes the usual sup norm on $[a, b]$.

Special Cases

- (a) Choosing $r = 0$ in Theorem 2.3, we get Theorem 1 in [2].
- (b) The choice of $r = 0$ in Corollary 2.4 reduces to Corollary 2 in [2].
- (c) If we replace the matrix A by the Cesàro matrix of order one and choose $r = 0$ in Corollary 2.4, then we get the statistical approximation theorem introduced by Gadjiev and Orhan (see Theorem 1 in [9]).
- (d) If we replace the matrix A by the identity matrix and also choose $r = 0$ in Corollary 2.4, then we get the classical Korovkin-type approximation theorem (see, for instance, [1], [13]).

Acknowledgment

The author would like to thank to the referee for his/her valuable suggestions which improved the paper considerably.

References

- [1] Devore, R.A.: The Approximation of Continuous Functions by Positive Linear Operators, Lecture Notes in Math. 293, Springer, Berlin, 1972.
- [2] Duman, O., Khan, M.K., Orhan, C.: *A*-statistical convergence of approximating operators. Math. Inequal. Appl. 6, 689-699 (2003).
- [3] Duman, O., Özarıslan, M.A., Doğru, O.: On integral type generalizations of positive linear operators. Studia Math. 174, 1-12 (2006).
- [4] Duman O., Orhan, C.: Statistical approximation by positive linear operators. Studia Math. 161, 187-197 (2004).
- [5] Erkuş, E., Duman, O.: *A*-statistical extension of the Korovkin type approximation theorem. Proc. Indian Acad. Sci. (Math. Sci.) 115, 499-508 (2005).
- [6] Fast, H.: Sur la convergence statistique. Colloq. Math. 2, 241-244 (1951).
- [7] Freedman, A.R., Sember, J.J.: Densities and summability. Pacific J. Math. 95, 293-305 (1981).
- [8] Fridy, J.A.: On statistical convergence. Analysis 5, 301-313 (1981).
- [9] Gadjiev, A.D., Orhan, C.: Some approximation theorems via statistical convergence. Rocky Mountain J. Math. 32, 129-138 (2002).
- [10] Hardy, G.H.: Divergent Series, Oxford Univ. Press, London, 1949.
- [11] Kirov, G., Popova, L.: A generalization of the linear positive operators. Math. Balkanica 7, 149-162 (1993).
- [12] Kolk, E.: Matrix summability of statistically convergent sequences. Analysis 13, 77-83 (1993).
- [13] Korovkin, P.P.: Linear Operators and Theory of Approximation, Hindustan Publ. Co., Delhi, 1960.
- [14] Miller, H.I.: A measure theoretical subsequence characterization of statistical convergence. Trans. Amer. Math. Soc. 347, 1811-1819 (1995).

Oktay DUMAN
TOBB Economics and Technology University,
Faculty of Arts and Sciences,
Department of Mathematics,
Söğütözü 06530, Ankara-TURKEY
e-mail: oduman@etu.edu.tr

Received 12.04.2006