

1-1-2007

## Multipliers and the Relative Completion in $L_w^p(G)$

CENAP DUYAR

A. TURAN GÜRKANLI

Follow this and additional works at: <https://journals.tubitak.gov.tr/math>



Part of the [Mathematics Commons](#)

---

### Recommended Citation

DUYAR, CENAP and GÜRKANLI, A. TURAN (2007) "Multipliers and the Relative Completion in  $L_w^p(G)$ ," *Turkish Journal of Mathematics*: Vol. 31: No. 2, Article 6. Available at: <https://journals.tubitak.gov.tr/math/vol31/iss2/6>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Mathematics by an authorized editor of TÜBİTAK Academic Journals. For more information, please contact [academic.publications@tubitak.gov.tr](mailto:academic.publications@tubitak.gov.tr).

## Multipliers and the Relative Completion in $L_w^p(G)$

*C. Duyar, A. T. Gürkanlı*

### Abstract

Quek and Yap defined a relative completion  $\tilde{A}$  for a linear subspace  $A$  of  $L^p(G)$ ,  $1 \leq p < \infty$ ; and proved that there is an isometric isomorphism, between  $\text{Hom}_{L^1(G)}(L^1(G), A)$  and  $\tilde{A}$ , where  $\text{Hom}_{L^1(G)}(L^1(G), A)$  is the space of the module homomorphisms (or multipliers) from  $L^1(G)$  to  $A$ . In the present, we defined a relative completion  $\tilde{A}$  for a linear subspace  $A$  of  $L_w^p(G)$ , where  $w$  is a Beurling's weighted function and  $L_w^p(G)$  is the weighted  $L^p(G)$  space, ([14]). Also, we proved that there is an algebraic isomorphism and homeomorphism, between  $\text{Hom}_{L_w^1(G)}(L_w^1(G), A)$  and  $\tilde{A}$ . At the end of this work we gave some applications and examples.

**Key words and phrases:** Module homomorphism (or multiplier), relative completion, essential module, weighted  $L^p(G)$  space. 1991 AMS subject classification codes 43.

### 1. Introduction and Preliminaries

Throughout this work,  $G$  will denote a non-compact and non-discrete locally compact Abelian group with Haar measure  $d\mu$ . The left and right translation operators are denoted by  $L_y f(x) = f(x - y)$ ,  $R_y f(x) = f(x + y)$  for all  $x, y \in G$ . The Fourier transform for any  $f \in L^1(G)$  is denoted by  $\hat{f}$ . A real-valued, measurable function  $w$  on  $G$  is said to be a weight function (Beurling's weight function) if  $1 \leq w(x)$  and  $w(x + y) \leq w(x)w(y)$  for  $x, y \in G$ . We say that  $w_1 \prec w_2$  if and only if there exists  $c > 0$  such that  $w_1(x) \leq cw_2(x)$

*AMS Mathematics Subject Classification:* Primary 55S10, 55S05

for all  $x \in G$ . Two weight functions are called equivalent and written  $w_1 \approx w_2$ , if  $w_1 \prec w_2$  and  $w_2 \prec w_1$ . For  $1 \leq p < \infty$ , we set

$$L_w^p(G) = \{f : fw \in L^p(G)\},$$

where  $L^p(G)$  denotes the usual Lebesgue space. It is a Banach space under the norm

$$\|f\|_{p,w} = \|fw\|_p = \left( \int_G |f(x)|^p w^p(x) d\mu(x) \right)^{\frac{1}{p}}.$$

If  $p = 1$ , then  $L_w^1(G)$  becomes an algebra under convolution, called Beurling algebra [14]. A weight function  $w$  is said to satisfy the Beurling Domar condition (shortly (BD)) if one has

$$\sum_{n \geq 1} \frac{\log w(nx)}{n^2} < \infty$$

for all  $x \in G$  [4]. Let  $(A, \|\cdot\|_A)$  be a Banach algebra and let  $(X, \|\cdot\|_X)$  a left Banach  $A$ -module, i. e.  $X$  is module over  $A$  in algebraic sense and satisfies  $\|a.b\|_X \leq \|a\|_A \|b\|_X$  for all  $a \in A$  and  $b \in X$ . Then the closed linear subspace of  $X$  spanned by

$$AX = \{ax : a \in A, x \in X\}$$

is called the essential part of  $X$  and is denoted by  $X_e$ . If  $X = X_e$ , then  $X$  is said to be an essential left Banach  $A$ -module [5].

If  $V$  and  $W$  are left(right) Banach  $A$ -modules, then a multiplier (or module homomorphism) from  $V$  to  $W$  is a bounded linear operator  $T$  from  $V$  to  $W$ , which commutes with module multiplication i. e.  $T(av) = aT(v)$  for  $a \in A$  and  $v \in V$ . We denote by  $\text{Hom}_A(V, W)$  or  $M(V, W)$  the space of multipliers from  $V$  to  $W$ .

## 2. The Main Results

**Definition 2.1** Let  $A$  be a linear subspace of  $L_w^p(G)$ ,  $1 \leq p < \infty$ , with the following properties:

(i)  $(A, \|\cdot\|_A)$  is a Banach  $L_w^1(G)$ -module under convolution with respect to norm  $\|\cdot\|_A$ , satisfying  $\|\cdot\|_{p,w} \leq \|\cdot\|_A$ .

(ii) There is a bounded approximate identity  $(e_\alpha)_{\alpha \in I}$  of  $L_w^1(G)$  such that  $\|f * e_\alpha - f\|_A \rightarrow 0$  for all  $f \in A$ , and  $\|e_\alpha\|_{1,w} < M$  for a number  $M > 0$  and all  $\alpha \in I$ .

The relative completion  $\tilde{A}$  of  $A$  is defined in the following way:

$$\tilde{A} = \left\{ f \in L_w^p(G) : f * e_\alpha \in A \text{ for all } \alpha \in I, \sup_{\alpha \in I} \|f * e_\alpha\|_A < \infty \right\}.$$

It is easy to see that  $\tilde{A}$  is a normed space with the norm  $\|\cdot\|_{\tilde{A}}$ , defined by

$$\|f\|_{\tilde{A}} = \sup_{\alpha \in I} \|f * e_\alpha\|_A.$$

**Proposition 2.2** Let  $A$  be as in Definition 2.1. The following properties are satisfied:

(i) If  $f \in \tilde{A}$  and  $g \in L_w^1(G)$ , then  $f * g \in \tilde{A}$  and

$$\|f * g\|_{\tilde{A}} \leq \|f\|_{\tilde{A}} \|g\|_{1,w}.$$

(ii) If  $f \in A$ , then

$$\|f\|_{\tilde{A}} \leq M \|f\|_A \quad \text{and} \quad \|f\|_A \leq \|f\|_{\tilde{A}},$$

i.e. the norms  $\|\cdot\|_A$  and  $\|\cdot\|_{\tilde{A}}$  are equivalent on  $A$ .

(iii)  $A$  is a closed subspace of  $\tilde{A}$ .

**Proof.** (i) Let  $f \in \tilde{A}$  and  $g \in L_w^1(G)$ . Then  $f * e_\alpha \in A$  for all  $\alpha \in I$  and  $\sup_{\alpha \in I} \|f * e_\alpha\|_A < \infty$ . Also, we can write  $(f * g) * e_\alpha = (f * e_\alpha) * g \in A$ , because  $A$  is a

Banach  $L_w^1(G)$ -module. Hence

$$\begin{aligned} \|f * g\|_{\tilde{A}} &= \sup_{\alpha \in I} \|(f * g) * e_\alpha\|_A \leq \sup_{\alpha \in I} (\|f * e_\alpha\|_A \|g\|_{1,w}) \\ &= \|g\|_{1,w} \sup_{\alpha \in I} \|f * e_\alpha\|_A = \|f\|_{\tilde{A}} \|g\|_{1,w}. \end{aligned}$$

(ii) Let  $f \in A$ . We have  $f * e_\alpha \in A$  for all  $\alpha \in I$ , because  $A$  is a Banach  $L_w^1(G)$ -module. By Definition 2.1, we have

$$\|f\|_{\tilde{A}} = \sup_{\alpha \in I} \|f * e_\alpha\|_A \leq \sup_{\alpha \in I} (\|f\|_A \|e_\alpha\|_{1,w}) \leq M \|f\|_A.$$

Now let any  $\varepsilon > 0$  be given. Then we can choose  $\alpha_0 \in I$  such that

$$\|f - f * e_\alpha\|_A < \varepsilon$$

for all  $\alpha \geq \alpha_0$ . Hence, we obtain

$$\|f\|_A \leq \|f - f * e_{\alpha_0}\|_A + \|f * e_{\alpha_0}\|_A < \varepsilon + \sup_{\alpha \in I} \|f * e_\alpha\|_A = \varepsilon + \|f\|_{\tilde{A}}.$$

This gives the inequality  $\|f\|_A \leq \|f\|_{\tilde{A}}$ .

(iii) It is easy to see that  $A \subset \tilde{A}$ . We will show that  $A$  is a complete space with respect to the norm  $\|\cdot\|_{\tilde{A}}$ . Let  $(f_n)$  be a Cauchy sequence in  $(A, \|\cdot\|_{\tilde{A}})$ . Then it is a Cauchy sequence in  $(A, \|\cdot\|_A)$  by (ii). Since  $(A, \|\cdot\|_A)$  is a Banach space, there is a function  $f \in A$  such that  $(f_n)$  converges to  $f$ . Also, since the norms  $\|\cdot\|_A$  and  $\|\cdot\|_{\tilde{A}}$  are equivalent on  $A$ , this implies that  $(f_n)$  converges to  $f$  with respect to the norm  $\|\cdot\|_{\tilde{A}}$ . Hence,  $(A, \|\cdot\|_{\tilde{A}})$  is a complete space. This shows that  $A$  is closed in  $\tilde{A}$ .  $\square$

**Proposition 2.3** *The definition of the space  $\tilde{A}$  does not depend on the approximate identity chosen.*

**Proof.** Let  $(e_\alpha)_{\alpha \in I}$  be as in Definition 2.1. Suppose that  $(v_\beta)_{\beta \in J}$  is an other bounded approximate identity in  $L_w^1(G)$  such that  $\|f * v_\beta - f\|_A \rightarrow 0$  for all  $f \in A$  and  $\|v_\beta\|_{1,w} < M_1$  for some  $M_1 > 0$ . Now, we define

$$X = \left\{ f \in L_w^p(G) : f * v_\beta \in A \text{ for all } \beta \in J, \sup_{\beta \in J} \|f * v_\beta\|_A < \infty \right\}$$

with the norm  $\|\cdot\|_X$ , defined by

$$\|f\|_X = \sup_{\beta \in J} \|f * v_\beta\|_A.$$

Assume that  $f \in \tilde{A}$ . Then we write  $\|f\|_{\tilde{A}} < \infty$  and  $f * e_\alpha \in A$  for all  $\alpha \in I$ . Since  $A$  is a Banach  $L_w^1(G)$ -module, we have  $f * e_\alpha * v_{\beta_0} = f * v_{\beta_0} * e_\alpha \in A$  for a fix  $\beta_0 \in J$ . So, we write

$$\|f * e_\alpha * v_{\beta_0} - f * v_{\beta_0}\|_{\tilde{A}} \leq \|f\|_{\tilde{A}} \|e_\alpha * v_{\beta_0} - v_{\beta_0}\|_{1,w}$$

by Proposition 2.2. Also, since  $(e_\alpha)_{\alpha \in I}$  is an approximate identity in  $L_w^1(G)$ , the net  $(f * e_\alpha * v_{\beta_0})_{\alpha \in I}$  converges to  $f * v_{\beta_0}$ . This shows that  $f * v_{\beta_0} \in A$ , because  $A$  is a closed subspace of  $\tilde{A}$  by Proposition 2.2. Therefore, we have  $f * v_\beta \in A$  for all  $\beta \in J$ . By (i) and (ii) in Proposition 2.2, we have

$$\begin{aligned} \|f * v_\beta\|_A &\leq \|f * v_\beta\|_{\tilde{A}} \leq \|f * v_\beta - f * v_\beta * e_\alpha\|_{\tilde{A}} + \|f * v_\beta * e_\alpha\|_{\tilde{A}} \\ &\leq \|f\|_{\tilde{A}} \|v_\beta - v_\beta * e_\alpha\|_{1,w} + \|f * e_\alpha\|_{\tilde{A}} \|v_\beta\|_{1,w} \\ &\leq \|f\|_{\tilde{A}} \|v_\beta - v_\beta * e_\alpha\|_{1,w} + MM_1 \|f * e_\alpha\|_A \\ &\leq \|f\|_{\tilde{A}} \|v_\beta - v_\beta * e_\alpha\|_{1,w} + MM_1 \|f\|_{\tilde{A}}. \end{aligned}$$

Hence

$$\begin{aligned} \|f\|_X &= \sup_{\beta \in J} \|f * v_\beta\|_A \leq \|f\|_{\tilde{A}} \sup_{\beta \in J} \|v_\beta - v_\beta * e_\alpha\|_{1,w} + MM_1 \|f\|_{\tilde{A}} \\ &\leq \|f\|_{\tilde{A}} \sup_{\beta \in J} (\|v_\beta\|_{1,w} + \|v_\beta\|_{1,w} \|e_\alpha\|_{1,w}) + MM_1 \|f\|_{\tilde{A}} \\ &\leq \|f\|_{\tilde{A}} (M_1 + MM_1) + MM_1 \|f\|_{\tilde{A}} = \|f\|_{\tilde{A}} (M_1 + 2MM_1). \end{aligned}$$

Thus, since  $\|f\|_{\tilde{A}} < \infty$ , we have  $\|f\|_X < \infty$ . That means  $f \in X$ . This shows that  $\tilde{A} \subset X$ .

Similary, we can show that  $X \subset \tilde{A}$ . This gives  $X = \tilde{A}$ .  $\square$

It is easy to prove the following Lemma 2.4.

**Lemma 2.4**  $L_w^p(G)$  is an essential Banach module over  $L_w^1(G)$ .

Now, we offer the statement of Corollary 8.10 in [13], used in the proof of the following lemma:

If  $A$  is a Banach algebra with approximate identity, and if  $W$  is an essential  $A$ -module which is reflexive as a Banach space, then

$$\text{Hom}_A(A, W) \cong W.$$

**Lemma 2.5** If  $p > 1$ , then there exists an isometric module isomorphism, between the spaces  $\text{Hom}_{L_w^1(G)}(L_w^1(G), L_w^p(G))$  and  $L_w^p(G)$ , i. e.

$$\text{Hom}_{L_w^1(G)}(L_w^1(G), L_w^p(G)) \cong L_w^p(G).$$

**Proof.** It is known that  $L_w^1(G)$  is a Banach algebra with approximate identity. By Lemma 2.4,  $L_w^p(G)$  is an essential Banach  $L_w^1(G)$ -module. If  $p > 1$ , it is also known that  $L_w^p(G)$  is reflexive as a Banach space [10]. Then, by Corollary 8.10 in [13], we have

$$\text{Hom}_{L_w^1(G)}(L_w^1(G), L_w^p(G)) \cong L_w^p(G).$$

□

**Theorem 2.6** *Let  $A$  be as in Definition 2.1. Then the spaces  $\text{Hom}_{L_w^1(G)}(L_w^1(G), A)$  and  $\tilde{A}$  are algebraically isomorphic and homeomorphic (i.e.  $\text{Hom}_{L_w^1(G)}(L_w^1(G), A) \cong \tilde{A}$ ), if  $p > 1$ , or if  $p = 1$  and  $\text{Hom}_{L_w^1(G)}(L_w^1(G), A) \subset L_w^1(G)$ .*

**Proof.** Assume that  $f \in \tilde{A}$ . We now define a mapping  $T_f$  such that  $T_f(g) = f * g$  for all  $g \in L_w^1(G)$ . By Definition of  $\tilde{A}$ , we have  $f * e_i - f * e_j \in A$ . Hence, by (i) and (ii) of Proposition 2.2,

$$\begin{aligned} \|f * g * e_i - f * g * e_j\|_A &\leq \|f * g * e_i - f * g * e_j\|_{\tilde{A}} \\ &\leq \|f\|_{\tilde{A}} \|g * e_i - g * e_j\|_{1,w} \\ &\leq \|f\|_{\tilde{A}} (\|g * e_i - g\|_{1,w} + \|g - g * e_j\|_{1,w}). \end{aligned}$$

Then, since  $(e_\alpha)_{\alpha \in I}$  is an approximate identity in  $L_w^1(G)$ ,  $(f * g * e_\alpha)_{\alpha \in I}$  is a Cauchy net in the space  $A$ . (It is known that, for a normed space, the definitions of completeness in the mean of topological vector space and in the mean of metric space are equivalent. Hence one can use net instead of sequence in the normed space). Since  $A$  is a Banach space, there is a function  $h \in A$  such that  $f * g * e_\alpha \rightarrow h$  in  $A$ . In addition, by (i) of Proposition 2.2,  $f * g$  is an element of  $\tilde{A}$ , where  $f \in \tilde{A}$  and  $g \in L_w^1(G)$ . Hence

$$\begin{aligned} \|f * g - h\|_{\tilde{A}} &\leq \|f * g - f * g * e_\alpha\|_{\tilde{A}} + \|f * g * e_\alpha - h\|_{\tilde{A}} \\ &\leq \|f\|_{\tilde{A}} \|g - g * e_\alpha\|_{1,w} + M \|f * g * e_\alpha - h\|_A. \end{aligned}$$

Since the net  $(f * g * e_\alpha)_{\alpha \in I}$  converges to  $h \in A$  and  $(e_\alpha)_{\alpha \in I}$  is an approximate identity in  $L_w^1(G)$ , this shows that  $T_f(g) = f * g = h \in A$ . Again, by Lemma 2.2, we obtain

$$\begin{aligned} \|T_f\| &= \sup_{g \neq 0} \frac{\|T_f(g)\|_A}{\|g\|_{1,w}} = \sup_{g \neq 0} \frac{\|f * g\|_A}{\|g\|_{1,w}} \\ &\leq \sup_{g \neq 0} \frac{\|f * g\|_{\tilde{A}}}{\|g\|_{1,w}} \leq \sup_{g \neq 0} \frac{\|f\|_{\tilde{A}} \|g\|_{1,w}}{\|g\|_{1,w}} = \|f\|_{\tilde{A}} \end{aligned}$$

for each  $g \in L_w^1(G)$ . This shows that  $T_f$  is a continuous mapping from  $L_w^1(G)$  to  $A$ . Then, it is easy to verify that  $T_f$  is a continuous module homomorphism from  $L_w^1(G)$  to  $A$ . By Definition 2.1, we have

$$\begin{aligned} \|T_f\| &= \sup_{g \neq 0} \frac{\|T_f(g)\|_A}{\|g\|_{1,w}} = \sup_{g \neq 0} \frac{\|f * g\|_A}{\|g\|_{1,w}} \geq \sup_{\alpha \in I} \frac{\|f * e_\alpha\|_A}{\|e_\alpha\|_{1,w}} \\ &\geq \sup_{\alpha \in I} \frac{\|f * e_\alpha\|_A}{M} = \frac{1}{M} \sup_{\alpha \in I} \|f * e_\alpha\|_A = \frac{1}{M} \|f\|_{\tilde{A}}. \end{aligned}$$

Conversely, assume that  $T \in \text{Hom}_{L_w^1(G)}(L_w^1(G), A)$ . By Definition 2.1, since  $A \subset L_w^p(G)$  and  $\|\cdot\|_{p,w} \leq \|\cdot\|_A$ , we obtain

$$\|T(g)\|_{p,w} \leq \|T(g)\|_A \leq \|T\| \|g\|_{1,w},$$

for all  $g \in L_w^1(G)$ . Then  $T \in \text{Hom}_{L_w^1(G)}(L_w^1(G), L_w^p(G))$ . Hence, by Lemma 2.5 for the case  $p > 1$  and by our hypothesis for the case  $p = 1$ , there is a function  $f \in L_w^p(G)$  such that  $T(g) = f * g$  for all  $g \in L_w^1(G)$ . Also, by Definition 2.1, we have

$$\begin{aligned} \|f\|_{\tilde{A}} &= \sup_{\alpha \in I} \|f * e_\alpha\|_A = M \sup_{\alpha \in I} \frac{\|f * e_\alpha\|_A}{M} \leq M \sup_{\alpha \in I} \frac{\|f * e_\alpha\|_A}{\|e_\alpha\|_{1,w}} \\ &\leq M \sup_{g \neq 0} \frac{\|f * g\|_A}{\|g\|_{1,w}} = M \sup_{g \neq 0} \frac{\|T(g)\|_A}{\|g\|_{1,w}} = M \|T\| < \infty. \end{aligned}$$

This gives  $f \in \tilde{A}$ . Thus, every  $T \in \text{Hom}_{L_w^1(G)}(L_w^1(G), A)$  is of the form  $T_f$  for some  $f \in \tilde{A}$  with  $\|T_f\| \leq \|f\|_{\tilde{A}}$  and  $\|f\|_{\tilde{A}} \leq M \|T_f\|$ . Hence, the mapping  $f \rightarrow T_f$  is an algebraic isomorphism and homeomorphism.  $\square$

## Applications

Let  $w$  be a weight function on a locally compact abelian group  $G$  and  $S_w(G)$  be the subalgebra of the Beurling algebra  $L_w^1(G)$  satisfying the following conditions:

- 1)  $S_w(G)$  is dense in  $L_w^1(G)$ .
- 2)  $S_w(G)$  is a Banach algebra under some norm  $\|\cdot\|_{S_w}$  and invariant under translations.



3) For each  $f \in S_w(G)$ ,  $\|L_y f\|_{S_w} \leq w(y) \|f\|_{S_w}$  for all  $y \in G$ .

4) Given any  $f \in S_w(G)$  and  $\varepsilon > 0$  there exists a neighbourhood  $U$  of the unit element  $e \in G$  such that  $\|L_y f - f\|_{S_w} < \varepsilon$  for all  $y \in U$ .

5) For all  $f \in S_w(G)$ ,  $\|f\|_{1,w} \leq \|f\|_{S_w}$ .

Some properties and multipliers of the space  $S_w(G)$  have been discussed in [1], [2], [3].

**Proposition 2.7**  $S_w(G)$  is an essential Banach convolution ideal in  $L_w^1(G)$ .

**Proof.** By definition of  $S_w(G)$ , we have  $\|f\|_{1,w} \leq \|f\|_{S_w}$  for all  $f \in S_w(G)$ . Hence, for any compact subset  $K \subset G$ , we write the inequality

$$\|f \cdot \chi_K\|_1 \leq \int_K |f(x)| w(x) dx \leq \|f\|_{1,w} \leq \|f\|_{S_w}. \quad (3.1)$$

Hence  $S_w(G)$  is a Banach function space (shortly BF-Space). Also since  $S_w(G)$  is translation invariant and the translation operator is continuous from  $G$  into  $S_w(G)$ , then  $S_w(G)$  is an essential Banach convolution ideal in a Beurling algebra  $L_{w_0}^1(G)$ , where  $w_0(x) = \max\{1, \|L_x\|\}$  [7]. A simple calculation shows that  $\|L_x\|_{1,w} \leq w(x)$ . Thus we have  $w_0(x) \leq w(x)$  for all  $x \in G$  and  $L_w^1(G) \subset L_{w_0}^1(G)$ . Now let  $f \in S_w(G)$  and  $g \in L_w^1(G)$ . Then we write

$$\|f * g\|_{S_w} \leq \|f\|_{S_w} \|g\|_{1,w_0} \leq \|f\|_{S_w} \|g\|_{1,w}. \quad (3.2)$$

Let  $(e_\alpha)_{\alpha \in I}$  be a bounded approximate identity in  $L_w^1(G)$ . Since  $(e_\alpha)_{\alpha \in I}$  is also a bounded approximate identity in  $L_{w_0}^1(G)$ , using (3.2) we have

$$\lim_\alpha \|f * e_\alpha - f\|_{S_w} = 0$$

for all  $f \in S_w(G)$ . Thus  $S_w(G)$  is an essential Banach convolution ideal in  $L_w^1(G)$  by Corollary 15.3 in [5]. Since the conditions of Definition 2.1 are satisfied, we have

$$\text{Hom}_{L_w^1(G)}(L_w^1(G), S_w(G)) \cong \tilde{S}_w(G)$$

by Theorem 2.7. □

**Examples** 1) Let  $1 \leq p, q < \infty$  and let  $w, \omega$  be weight functions on a locally compact abelian group  $G$  and its dual group  $\hat{G}$  respectively. We set

$$A_{w,\omega}^{p,q}(G) = \left\{ f : f \in L_w^p(G), \hat{f} \in L_{\omega}^q(G) \right\}$$

and equip this space with the norm

$$\|f\|_{w,\omega}^{p,q} = \|f\|_{p,w} + \left\| \hat{f} \right\|_{q,\omega},$$

where  $(\wedge)$  is the generalized Fourier transform. It has been proved in [8] that  $A_{w,\omega}^{p,q}(G)$  is a Banach space. Also if  $w$  satisfies Beurling-Domar condition (BD), then  $A_{w,\omega}^{p,q}(G)$  admits an approximate identity bounded in  $L_w^1(G)$ . Furthermore,  $A_{w,\omega}^{p,q}(G)$  is an essential Banach convolution module over  $L_w^1(G)$  by Proposition 1.13 (a) in [8]. Hence

$$\text{Hom}_{L_w^1(G)}(L_w^1(G), A_{w,\omega}^{p,q}(G)) \cong \tilde{A}_{w,\omega}^{p,q}(G)$$

by Theorem. 2.7.

2) Let  $G$  be a locally compact Abelian group. Suppose that  $A_w(G) = L_w^1(G) \cap L_w^p(G)$ ,  $1 \leq p < \infty$ . In [11], Oztop and Gürkanlı proved that  $A_w(G)$  is a Banach convolution algebra with the norm  $\|f\|_{A_w} = \|f\|_{1,w} + \|f\|_{p,w}$ , and investigated some properties of this space. It is easy to see that  $A_w(G) \subset L_w^p(G)$  and  $\|\cdot\|_{p,w} \leq \|\cdot\|_{A_w}$ . Also,  $A_w(G)$  is a Banach  $L_w^1(G)$ -module [11]. It is known that  $L_w^1(G)$  has a bounded approximate identity [14]. It is also an approximate identity for  $A_w(G)$  [10]. Hence the condition of Definiton 2.1 are satisfied. Thus we have

$$\text{Hom}_{L_w^1(G)}(L_w^1(G), A_w(G)) \cong \tilde{A}_w(G).$$

3) Let  $G$  be a locally compact Abelian group. In [2], Feichtinger and Gürkanlı defined the linear space

$$A_w^p(G) = \left\{ f : f \in L_w^1(G), \hat{f} \in L_w^p(G) \right\}, \quad 1 \leq p < \infty.$$

by Theorem 2.7. In the mentioned paper, it is shown that  $A_w^p(G)$  is a Banach convolution algebra with the norm  $\|\cdot\|_w^p$ , defined by  $\|f\|_w^p = \|f\|_{1,w} + \left\| \hat{f} \right\|_{p,w}$ . Also, they have

investigated some properties of this space. It is easy to see that  $A_w^p(G) \subset L_w^1(G)$  and  $\|\cdot\|_{1,w} \leq \|\cdot\|_w^p$ . Also,  $(A_w^p(G), \|\cdot\|_w^p)$  is a Banach ideal over  $L_w^1(G)$ . Since  $w$  satisfies (BD), then  $L_w^1(G)$  has a bounded approximate identity  $(e_\alpha)_{\alpha \in I}$ , whose Fourier transforms have compact support and  $\|f * e_\alpha - f\|_w^p \rightarrow 0$  for all  $f \in A_w^p(G)$  by Theorem 4.2 in [2]. Hence

$$\text{Hom}_{L_w^1(G)}(L_w^1(G), A_w^p(G)) \cong \tilde{A}_w^p(G)$$

by Theorem 2.7.

### Acknowledgement

The authors want to thank the referee for his significant suggestions regarding this paper.

### References

- [1] Cigler, J.: Normed ideals in  $L_1(G)$ , Indag.Math. 31, 272-282 (1969).
- [2] Dogan, M., Gürkanlı, A.T.: Multipliers of the space  $S_w(G)$ , Mathematica Balkanica, New Series 15, Fasc.3-4. 199-212 (2001).
- [3] Dogan, M., Gürkanlı, A.T.: On functions with Fourier transform in  $S_w(G)$ , Bull. Cal. Math. Soc. 92(2). 111-120 (2000).
- [4] Domar, Y.: Harmonic analysis based on certain commutative Banach algebras, Acta Math. 96, 1-66 (1956).
- [5] Doran, R., S.Wichmann, J.: Approximate Identities and Factorization in Banach Modules, Lecture Notes in Mathematics, Springer Verlag, 305 (1979).
- [6] Feichtinger, H.G., Gürkanlı, A. T.: 1990. On a family of weighted convolution algebras, Internat. J. Math. Sci. 3, 13, 517-525 (1990).
- [7] Feichtinger, H.G.: A compactness criterion for translation invariant Banach spaces of functions, Analysis Mathematica. 8, 165-172 (1982).
- [8] Fischer, R.H., Gürkanlı, A.T.: Liu, T.S., On a family of weighted spaces, Math.Slovaca. 46, 1, 71-82 (1996).

DUYAR, GÜRKANLI

- [9] Larsen,R.: Introduction to the theory of multipliers , Springer –Verlag, Berlin-Heidelberg-NewYork 1971.
- [10] Murty, G. N. K.-Umni, K. R.: Multipliers on weighted spaces, Functional Analysis and its applications, International Conference Madras, Lecture Notes in Mathematics, Springer Verlag, 273-291 (1973).
- [11] Oztop, S.-Gürkanlı, A. T.: Weighted  $L^1(G) \cap L^p(G)$  spaces and some properties, Sixth National Mathematics Symposium, 196-211 (1993).
- [12] Quek, T. S.-Yap, L., H.: Multipliers From  $L^1(G)$  to a Lipshitz space, Journal of Mathematical Analysis on Applications. 69, 531-579 (1979).
- [13] Rieffel, H.: Induced Banach Representation of Banach Algebras and Locally compact groups, Locally compact groups, Journal of Functional Analysis. 1, 443-491 (1967).
- [14] Rieter, H.: Classical Harmonic Analysis and Locally compact groups, Oxford at the Claderon Press 1968.

C. DUYAR, A. T. GÜRKANLI  
Ondokuz Mayıs University  
Faculty of Arts and Sciences  
Department of Mathematics  
55139, Kurupelit, Samsun-TURKEY  
e-mail: gurkanli@omu.edu.tr  
e-mail: cenapd@omu.edu.tr

Received 18.10.2005