

1-1-2007

## Uniqueness of Coprimary Decompositions

M. MAANI-SHIRAZI

P. F. SMITH

Follow this and additional works at: <https://journals.tubitak.gov.tr/math>



Part of the [Mathematics Commons](#)

---

### Recommended Citation

MAANI-SHIRAZI, M. and SMITH, P. F. (2007) "Uniqueness of Coprimary Decompositions," *Turkish Journal of Mathematics*: Vol. 31: No. 1, Article 4. Available at: <https://journals.tubitak.gov.tr/math/vol31/iss1/4>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Mathematics by an authorized editor of TÜBİTAK Academic Journals. For more information, please contact [academic.publications@tubitak.gov.tr](mailto:academic.publications@tubitak.gov.tr).

## Uniqueness of Coprimary Decompositions

*M. Maani-Shirazi and P. F. Smith*

### Abstract

Uniqueness properties of coprimary decompositions of modules over non-commutative rings are presented.

**Key Words:** Coprimary, decomposition, normal decomposition, prime ideal, left Noetherian ring, right Noetherian ring.

### 1. Introduction

Throughout this paper,  $R$  is a ring (not necessarily commutative) with an identity element  $1 \neq 0$  and  $M$  is a non-zero unital left  $R$ -module. For any submodules  $N, L$  of  $M$ , we define  $(N : L) = \{r \in R : rL \subseteq N\}$ . Note that  $(N : L)$  is an ideal of  $R$ . Moreover,  $(N : L) = R$  if and only if  $L \subseteq N$ . Let  $N$  be a submodule of  $M$  and let  $A$  be an ideal of  $R$ ; we set  $(N :_M A) = \{m \in M : Am \subseteq N\}$ . Note that  $(N :_M A)$  is a submodule of  $M$ .

In this paper, by making use of the technique employed in [7], we shall prove uniqueness properties of coprimary decompositions.

Note that, when  $R$  is a commutative Noetherian ring,  $M$  is coprimary if and only if  $M$  is secondary. It is well known that every non-zero injective module over a commutative Noetherian ring has a secondary representation (see [6]). By a similar method to that used in [6], we obtain the following result. For  $R$  non-commutative left and right Noetherian we show that if  $M$  is injective and if the zero ideal of  $R$  is a finite intersection of strongly primary ideals, then  $M$  has a coprimary decomposition.

## 2. Coprimary Decompositions

**Definition.** Given a prime ideal  $P$  of  $R$ , a non-zero module  $M$  is called  $P$ -coprimary if

- (i)  $(N : M) \subseteq P$  for every proper submodule  $N$  of  $M$ , and
- (ii)  $P^h \subseteq (0 : M)$  for some positive integer  $h$ .

Note that if  $M$  is  $P$ -coprimary, then  $P^h \subseteq (0 : M) \subseteq P$  for some positive integer  $h$ .  
 $M$  is called coprimary if it is  $P$ -coprimary for some prime ideal  $P$  of  $R$ .

A non-zero module  $M$  has a coprimary decomposition if there exist a positive integer  $n$  and submodules  $M_i (1 \leq i \leq n)$  of  $M$  such that

- (i)  $M = M_1 + \cdots + M_n$ , and
- (ii)  $M_i$  is coprimary for each  $1 \leq i \leq n$ .

If  $M$  has a coprimary decomposition, then we say that  $M$  has a normal coprimary decomposition if there exist a positive integer  $n$ , distinct prime ideals  $P_i (1 \leq i \leq n)$  of  $R$ , and  $P_i$ -coprimary submodules  $M_i (1 \leq i \leq n)$  of  $M$  such that

- (i)  $M = M_1 + \cdots + M_n$ , and
- (ii)  $M \neq M_1 + \cdots + M_{i-1} + M_{i+1} + \cdots + M_n$  for all  $1 \leq i \leq n$ .

**Lemma 2.1.** Let  $P$  be a prime ideal of  $R$  and let  $M$  be a  $P$ -coprimary module. Then  $M/K$  is a  $P$ -coprimary  $R$ -module for each proper submodule  $K$  of  $M$ .

**Proof.** This is clear. □

**Corollary 2.2.** If  $M$  has a coprimary decomposition, then  $M/K$  has a coprimary decomposition for every proper submodule  $K$  of  $M$ .

**Proof.** There exist a positive integer  $n$  and coprimary submodules  $M_i (1 \leq i \leq n)$  of  $M$  such that  $M = M_1 + \cdots + M_n$ . Then  $M/K = ((M_1 + K)/K) + \cdots + ((M_n + K)/K)$ . Then, for each  $1 \leq i \leq n$ ,  $(M_i + K)/K \cong M_i/(M_i \cap K)$  so that  $(M_i + K)/K = 0$  or  $(M_i + K)/K$  is coprimary by Lemma 2.1. □

**Lemma 2.3.** Let  $P$  be a prime ideal of  $R$ , let  $n$  be a positive integer, and let  $M_i (1 \leq i \leq n)$  be non-zero left  $R$ -modules. Then the  $R$ -module  $M_1 \oplus \cdots \oplus M_n$  is  $P$ -coprimary if and only if  $M_i$  is  $P$ -coprimary for each  $1 \leq i \leq n$ .

**Proof.** ( $\Rightarrow$ ) This follows from Lemma 2.1.

( $\Leftarrow$ ) Let  $N$  be a proper submodule of the module  $M = M_1 \oplus \cdots \oplus M_n$ . There exists  $1 \leq i \leq n$  such that  $M_i \not\subseteq N$ . Then  $(M_i + N)/N \cong M_i/(M_i \cap N)$  and  $M_i \cap N$  is a proper

submodule of  $M_i$  so that  $(N : M) \subseteq (N : M_i + N) \subseteq P$ . There exists a positive integer  $h$  such that  $P^h \subseteq (0 : M_i)$  for each  $1 \leq i \leq n$ . Then  $P^h \subseteq \bigcap_{i=1}^n (0 : M_i) = (0 : M)$ . Thus  $M$  is  $P$ -coprimary.  $\square$

**Corollary 2.4.** *Let  $P$  be a prime ideal of  $R$ , let  $n$  be a positive integer, and let  $M_i (1 \leq i \leq n)$  be  $P$ -coprimary submodules of  $M$ . Then the submodule  $M_1 + \cdots + M_n$  of  $M$  is a  $P$ -coprimary  $R$ -module.*

**Proof.** This follows from Lemmas 2.1 and 2.3.  $\square$

**Corollary 2.5.** *If  $M$  has a coprimary decomposition, then  $M$  has a normal coprimary decomposition.*

**Proof.** This follows from Corollary 2.4.  $\square$

One can easily prove the following result.

**Lemma 2.6.** *Let  $P$  be a prime ideal of  $R$  and let  $M$  be a semisimple module. Then the following statements are equivalent.*

- (i)  $M$  is  $P$ -coprimary.
- (ii) Every non-zero submodule of  $M$  is  $P$ -coprimary.
- (iii) Every simple submodule of  $M$  is  $P$ -coprimary.

**Corollary 2.7.** *Let  $M$  be a semisimple module. Then  $M$  has a coprimary decomposition if and only if the set  $\{(0 : N) : N \text{ is a simple submodule of } M\}$  is finite.*

**Proof.** This follows from Lemma 2.6.  $\square$

**Lemma 2.8.** *Let  $P$  be a prime ideal of  $R$ . Then  $M$  is  $P$ -coprimary if and only if, for every ideal  $A$  of  $R$ ,  $M = AM$  if  $A \not\subseteq P$  and there exists a positive integer  $h$  such that  $A^h M = 0$  if  $A \subseteq P$ .*

**Proof.** This is straightforward.  $\square$

**Lemma 2.9.** *If  $M$  has a coprimary decomposition, then for each ideal  $A$  of  $R$  there exists a positive integer  $h$  such that  $M = AM + (0 :_M A^h)$ .*

**Proof.** There exist a positive integer  $n$ , prime ideals  $P_i (1 \leq i \leq n)$  of  $R$ , and  $P_i$ -

coprimary submodules  $M_i (1 \leq i \leq n)$  of  $M$  such that  $M = M_1 + \cdots + M_n$ . Let  $A$  be an ideal of  $R$ . For each  $1 \leq i \leq n$ , Lemma 2.8 gives  $M_i = AM_i$  or  $M_i \subseteq (0 :_M A^{h_i})$  for some positive integer  $h_i$ . Let  $h = \max_{1 \leq i \leq n} h_i$ . Then  $M_i \subseteq AM + (0 :_M A^h)$  for all  $1 \leq i \leq n$ . It follows that  $M = AM + (0 :_M A^h)$ .  $\square$

We shall be interested in the following property of a ring  $R$ .

(P) For every proper ideal  $A$  of  $R$  there exists a positive integer  $n$  such that  $B^n \subseteq A$  for every ideal  $B$  of  $R$  with  $B^h \subseteq A$  for some positive integer  $h$ .

Note that any ring which has the ascending chain condition on two-sided ideals or any ring in which prime ideals are finitely generated left ideals satisfies the property (P) (see [3, Lemma 3.1]).

**Lemma 2.10.**  *$R$  satisfies (P) if and only if for every proper ideal  $A$  of  $R$ , the sum of all nilpotent ideals of the ring  $R/A$  is also a nilpotent ideal of  $R/A$ .*

**Proof.** ( $\Leftarrow$ ) This is clear.

( $\Rightarrow$ ) Let  $C$  be the ideal of  $R$  containing  $A$  such that  $C/A$  is the sum of all nilpotent ideals of the ring  $R/A$ . Let  $n$  be the positive integer in the property (P). Let  $c_i \in C (1 \leq i \leq n)$ . There exist a positive integer  $h$  and ideals  $B_j (1 \leq j \leq h)$  of  $R$  such that  $B_j^n \subseteq A \subseteq B_j (1 \leq j \leq h)$  and  $c_i \in B_1 + \cdots + B_h (1 \leq i \leq n)$ . Note that  $(B_1 + \cdots + B_h)^{hn} \subseteq A$  and hence  $(B_1 + \cdots + B_h)^n \subseteq A$ . This implies that  $c_1 \cdots c_n \in A$ . Thus  $C^n \subseteq A$ .  $\square$

**Lemma 2.11.** *Let  $R$  satisfy the property (P). Then  $M$  is coprimary if and only if for every ideal  $A$  of  $R$  either  $M = AM$  or  $A^h M = 0$  for some positive integer  $h$ .*

**Proof.** ( $\Rightarrow$ ) This follows from Lemma 2.8.

( $\Leftarrow$ ) Let  $P$  be the ideal of  $R$  containing  $A = (0 : M)$  such that  $P/A$  is the sum of all nilpotent ideals of the ring  $R/A$ . By Lemma 2.10, there exists a positive integer  $n$  such that  $P^n \subseteq A$ . Let  $B, C$  be ideals of  $R$  such that  $BC \subseteq P$ . If  $M = BM$  and  $M = CM$ , then  $M = BM = BCM \subseteq PM$  so that  $M = PM = P^2M = \cdots = P^n M = 0$ , a contradiction. Thus  $M \neq BM$  or  $M \neq CM$ . By the hypothesis,  $B \subseteq P$  or  $C \subseteq P$ . It follows that  $P$  is a prime ideal of  $R$  and hence  $M$  is  $P$ -coprimary by Lemma 2.8.

Next we give an example to show that in Lemma 2.11 the condition on  $R$  is necessary.

**Example 2.12.** Let  $p$  be any prime number, let  $F$  be a field of characteristic  $p$ , let  $G$  be the Prüfer  $p$ -group, and let  $R$  be the group algebra  $F[G]$ . (See [2, p.37] for the definition of Prüfer groups). Then  $R$  is a commutative ring with unique maximal ideal  $J = \sum_{g \in G} R(g - 1)$  and  $J$  is a nil ideal of  $R$  such that  $J = J^2$ . If  $A$  is any ideal of  $R$  then  $A$  is nilpotent unless  $A = J$  or  $A = R$ . Now let  $M$  denote the  $R$ -module  $J$ . Then, for any ideal  $A$  of  $R$ ,  $M = AM$  or  $A^k M = 0$  for some positive integer  $k$ . However,  $M$  is not coprimary because  $J$  is the only prime ideal of  $R$  and  $M = JM$ .

**Theorem 2.13.** Let  $M$  have a coprimary decomposition. Let  $M = K_1 + \cdots + K_s$  and  $M = L_1 + \cdots + L_t$  be normal coprimary decompositions of  $M$  where  $K_i$  is  $P_i$ -coprimary for some prime ideal  $P_i$  ( $1 \leq i \leq s$ ) and  $L_j$  is  $Q_j$ -coprimary for some prime ideal  $Q_j$  ( $1 \leq j \leq t$ ). Then  $s = t$  and  $\{P_1, \dots, P_s\} = \{Q_1, \dots, Q_t\}$ .

**Proof.** Without loss of generality, we can suppose that  $P_1$  is maximal in the set  $\{P_1, \dots, P_s\} \cup \{Q_1, \dots, Q_t\}$ . There exists a positive integer  $n$  such that  $P_1^n K_1 = 0$ . Thus

$$P_1^n M = P_1^n K_1 + \cdots + P_1^n K_s \subseteq K_2 + \cdots + K_s,$$

also

$$P_1^n M = P_1^n L_1 + \cdots + P_1^n L_t.$$

Because  $M \neq P_1^n M$ , there exists a positive integer  $j$  such that  $1 \leq j \leq t$  and  $L_j \neq P_1^n L_j$  and hence  $P_1^n \subseteq Q_j$  by Lemma 2.8. This implies that  $P_1 \subseteq Q_j$ . Without loss of generality, we can suppose that  $j = 1$  and hence  $P_1 = Q_1$ . We can suppose that  $P_1^n K_1 = Q_1^n L_1 = 0$ . Then Lemma 2.8 gives

$$P_1^n M = P_1^n K_1 + \cdots + P_1^n K_s = K_2 + \cdots + K_s,$$

and

$$P_1^n M = P_1^n L_1 + \cdots + P_1^n L_t = L_2 + \cdots + L_t.$$

By induction,  $s = t$  and  $\{P_i : 2 \leq i \leq s\} = \{Q_j : 2 \leq j \leq s\}$ . The result follows.  $\square$

In view of Theorem 2.13, we call prime ideals  $P_i$  ( $1 \leq i \leq s$ ) of  $R$  the coassociated prime ideals of  $M$  provided there exists a normal coprimary decomposition  $M = K_1 + \cdots + K_s$ ,

where  $K_i$  is a  $P_i$ -coprimary submodule of  $M$  for each  $1 \leq i \leq s$ .

**Theorem 2.14.** *Let  $M$  have a coprimary decomposition and let  $P_i (1 \leq i \leq n)$  be the coassociated prime ideals of  $M$ , for some positive integer  $n$ . Suppose that there exists  $1 \leq k \leq n$  such that for all  $1 \leq i \leq k$  and all  $k+1 \leq j \leq n$ ,  $P_j \not\subseteq P_i$ . Let  $M = M_1 + \cdots + M_n$  and  $M = L_1 + \cdots + L_n$  be normal coprimary decompositions of  $M$  in terms of  $P_i$ -coprimary submodules  $M_i$  and  $L_i (1 \leq i \leq n)$ . Then  $M_1 + \cdots + M_k = L_1 + \cdots + L_k$ .*

**Proof.** There exists a positive integer  $s$  such that  $P_j^s M_j = P_j^s L_j = 0 (k+1 \leq j \leq n)$ . Let  $A = P_{k+1}^s \cdots P_n^s$ . Then for all  $1 \leq i \leq k$ ,  $A \not\subseteq P_i$  so that  $M_i = AM_i$  and  $L_i = AL_i$ . Now we have

$$AM = AM_1 + \cdots + AM_k + AM_{k+1} + \cdots + AM_n = M_1 + \cdots + M_k,$$

and

$$AM = AL_1 + \cdots + AL_k + AL_{k+1} + \cdots + AL_n = L_1 + \cdots + L_k.$$

Thus  $M_1 + \cdots + M_k = L_1 + \cdots + L_k$ .  $\square$

Let  $P$  be a prime ideal of  $R$ .  $M^P$  is defined to be the intersection  $\cap AM$ , where  $A$  runs over the ideals of  $R$  not contained in  $P$ .

**Remark 2.15.** *Let  $M = M_1 + \cdots + M_n$  and  $M = L_1 + \cdots + L_n$  be normal coprimary decompositions of  $M$  where  $n$  is a positive integer and, for each  $1 \leq i \leq n$ ,  $M_i$  and  $L_i$  are  $P_i$ -coprimary submodules of  $M$  for some prime ideal  $P_i$  of  $R$ . If  $P_j$  is minimal in the set  $\{P_1, \dots, P_n\}$ , then  $M_j = L_j$  by Theorem 2.14. Moreover, we have also  $M_j = L_j = M^P$  (see [5]).*

Next, we give a characterization of the coassociated prime ideals of  $M$  with coprimary decomposition.

**Theorem 2.16.** *Let  $P$  be a prime ideal of  $R$  and let  $M$  have a coprimary decomposition. Then  $P$  is a coassociated prime ideal of  $M$  if and only if  $P = (K : M)$  for some proper submodule  $K$  of  $M$ .*

**Proof.** Let  $M = M_1 + \cdots + M_n$  be a normal coprimary decomposition of  $M$  where  $n$  is a positive integer and, for each  $1 \leq i \leq n$ ,  $M_i$  is a  $P_i$ -coprimary submodule of  $M$  for some prime ideal  $P_i$  of  $R$ . Let  $P$  be a coassociated prime ideal of  $M$ . Without loss of generality, we can suppose that  $P = P_1$ . There exists a positive integer  $k$  such that  $P^k M_1 = 0$ . Thus

$M = M_1 + M_2 + \cdots + M_n$  but  $M \neq P^k M_1 + M_2 + \cdots + M_n$ . There exists  $1 \leq j \leq k$  such that  $M = P^{j-1} M_1 + M_2 + \cdots + M_n$  but  $M \neq P^j M_1 + M_2 + \cdots + M_n$ . Let  $K$  denote the proper submodule  $P^j M_1 + M_2 + \cdots + M_n$ .

Let  $A = (K : M)$ . Clearly  $PM \subseteq K$  gives  $P \subseteq A$ . If  $P \neq A$ , then  $M_1 = AM_1$  and hence  $M_1 \subseteq AM \subseteq K$  so that  $K = M$ , a contradiction. Thus  $P = A$ .

Conversely, let  $Q$  be a prime ideal of  $R$  such that  $Q = (N : M)$  for some proper submodule  $N$  of  $M$ . There exists  $1 \leq i \leq n$  such that  $M_i \not\subseteq N$ . Without loss of generality, we can suppose that there exists  $1 \leq t \leq n$  such that  $M_i \not\subseteq N$  for all  $1 \leq i \leq t$  but  $M_i \subseteq N$  for all  $t+1 \leq i \leq n$ . For each  $1 \leq i \leq t$ ,  $M_i \cap N$  is a proper submodule of  $M_i$  and  $QM_i \subseteq M_i \cap N$  so that  $Q \subseteq P_i$ . There exists a positive integer  $s$  such that  $P_i^s M_i = 0$  ( $1 \leq i \leq t$ ). Now  $M = M_1 + \cdots + M_n = M_1 + \cdots + M_t + N$  and hence  $(P_1^s \cdots P_t^s)M \subseteq N$  so that  $P_1^s \cdots P_t^s \subseteq Q$ . It follows that there exists  $1 \leq j \leq t$  such that  $P_j \subseteq Q$  and hence  $P_j = Q$ . Therefore  $Q$  is a coassociated prime ideal of  $R$ .  $\square$

**Lemma 2.17.** *If  $M$  has a coprimary decomposition, then every minimal prime ideal over  $A = (0 : M)$  is a coassociated prime ideal of  $M$ .*

**Proof.** Let  $M = M_1 + \cdots + M_n$  be a normal coprimary decomposition of  $M$  where  $n$  is a positive integer and, for each  $1 \leq i \leq n$ ,  $M_i$  is a  $P_i$ -coprimary submodule of  $M$  for some prime ideal  $P_i$  of  $R$ . Then  $A = \bigcap_{i=1}^n (0 : M_i)$ . Suppose  $Q$  is a minimal prime ideal of  $A$ . There exists  $1 \leq i \leq n$  such that  $A \subseteq (0 : M_i) \subseteq Q$ . So  $Q = P_i$ .  $\square$

**Lemma 2.18.** *Let  $R$  be a prime left or right Noetherian ring and let  $M = PM$  for all non-zero prime ideals  $P$  of left  $R$ -module. Then  $M$  is 0-coprimary.*

**Proof.** Let  $A$  be a non-zero ideal of  $R$ . There exist a positive integer  $n$ , prime ideals  $P_i$  ( $1 \leq i \leq n$ ) of  $R$  such that  $P_1 \cdots P_n \subseteq A \subseteq P_1 \cap \cdots \cap P_n$ . But  $M = P_i M$  for all  $1 \leq i \leq n$ . So  $M = P_n M = \cdots = P_1 \cdots P_n M \subseteq AM$  and hence  $M = AM$ . Lemma 2.8 yields that  $M$  is 0-coprimary.  $\square$

**Remark 2.19.** *Let  $R$  be left or right Noetherian. Then there exists a prime ideal  $P$  of  $R$  such that  $PM \neq M$ . For, suppose that  $QM = M$  for all prime ideals  $Q$  of  $R$ . There exist a positive integer  $n$  and prime ideals  $P_i$  ( $1 \leq i \leq n$ ) of  $R$  such that  $0 = P_1 \cdots P_n$ .*



Then we have  $M = P_n M = \cdots = P_1 \cdots P_n M = 0$ , a contradiction.

**Corollary 2.20.** *Let  $R$  be left or right Noetherian. Then  $M$  has a coprimary quotient  $R$ -module.*

**Proof.** Since  $R$  is left or right Noetherian and by Remark 2.19, there exists a prime ideal  $P$  of  $R$  such that  $PM \neq M$  but  $QM = M$  for all prime ideals  $Q$  of  $R$  properly containing  $P$ . Then  $M/PM$  is a 0-coprimary  $(R/P)$ -module by Lemma 2.18 which implies that  $M/PM$  is a  $P$ -coprimary  $R$ -module.  $\square$

For any non-empty set  $I$ ,  $M^{(I)}$  is the direct sum  $\bigoplus_{i \in I} M_i$ , where  $M_i = M(i \in I)$ .

The prime radical,  $\sqrt{A}$ , of an ideal  $A$  of  $R$  is defined to be the intersection of all prime ideals which contain  $A$ .

**Lemma 2.21.** *Let  $P$  be a prime ideal of  $R$  and let  $M$  be  $P$ -coprimary. Then  $M^{(I)}$  is a  $P$ -coprimary  $R$ -module for every non-empty set  $I$ .*

**Proof.** We have  $\sqrt{(0 : M^{(I)})} = \sqrt{(0 : M)} = P$ . There exists a positive integer  $n$  such that  $P^n M = 0$ . Let  $A$  be an ideal of  $R$ . If  $A \subseteq P$  then  $A^n M^{(I)} = (A^n M)^{(I)} \subseteq (P^n M)^{(I)} = 0$ . Now suppose that  $A \not\subseteq P$ . Then  $AM = M$  and so  $AM^{(I)} = (AM)^{(I)} = M^{(I)}$ . By Lemma 2.8,  $M^{(I)}$  is  $P$ -coprimary.  $\square$

Recall that any left  $R$ -module is  $M$ -generated if it is a quotient module of  $M^{(I)}$  for some non-empty set  $I$ .

**Corollary 2.22.** *If  $M$  has a coprimary decomposition, then any non-zero  $M$ -generated  $R$ -module has a coprimary decomposition.*

**Proof.** Let  $M = M_1 + \cdots + M_n$  be a coprimary decomposition of  $M$  where  $n$  is a positive integer and, for each  $1 \leq i \leq n$ ,  $M_i$  is a  $P_i$ -coprimary submodule of  $M$  for some prime ideal  $P_i$  of  $R$ . Let  $I$  be a non-empty set. Then we have  $M^{(I)} = M_1^{(I)} + \cdots + M_n^{(I)}$ . Lemma 2.21 yields that  $M_i^{(I)}$  is  $P_i$ -coprimary for each  $1 \leq i \leq n$ . Hence  $M^{(I)}$  has a coprimary decomposition. Corollary 2.2 completes the proof.  $\square$

**Remark 2.23.** *Let  $P$  be a maximal ideal of  $R$ . Then  $M$  is  $P$ -coprimary if and only if  $P^n M = 0$  for some positive integer  $n$ . In this case, every non-zero submodule of  $M$  is*

also  $P$ -coprimary.

**Theorem 2.24.** *The following statements are equivalent.*

- (i) *Every non-zero left  $R$ -module has a coprimary decomposition.*
- (ii) *The left  $R$ -module  $R$  has a coprimary decomposition.*
- (iii) *There exist positive integers  $n, h$  and maximal ideals  $P_i (1 \leq i \leq n)$  of  $R$  such that  $P_1^h \cap \cdots \cap P_n^h = 0$ .*

**Proof.** (i)  $\Rightarrow$  (ii) This is clear.

(ii)  $\Rightarrow$  (i) This follows from Corollary 2.22.

(ii)  $\Rightarrow$  (iii) Let  $R = A_1 + \cdots + A_n$  be a coprimary decomposition of the left  $R$ -module  $R$  where  $n$  is a positive integer and, for each  $1 \leq i \leq n$ ,  $A_i$  is a  $P_i$ -coprimary submodule of the left  $R$ -module  $R$  for some prime ideal  $P_i$  of  $R$ . There exists a positive integer  $h$  such that  $P_i^h A_i = 0$  for each  $1 \leq i \leq n$  and so  $P_1^h \cap \cdots \cap P_n^h = 0$ . Note that any prime ring which has a coprimary decomposition as a left module over itself has only two ideals. Let  $P$  be a prime ideal of  $R$ . Then the ring  $R/P$  is prime with coprimary decomposition as a module over itself. So  $R/P$  has only two ideals, i.e.,  $P$  is maximal. We have shown that every prime ideal of  $R$  is maximal. The result follows.

(iii)  $\Rightarrow$  (ii) We have  $R \cong R/P_1^h \oplus \cdots \oplus R/P_n^h$  as left  $R$ -modules. It follows that  $R$  has a coprimary decomposition as a left  $R$ -module by Remark 2.23.  $\square$

There are modules in which every non-zero submodule has a coprimary decomposition (see [5]). In certain situations it is possible to write down explicitly a normal coprimary decomposition for a non-zero module once its coassociated prime ideals are known, as we show next.

**Theorem 2.25.** *Suppose that every non-zero submodule of  $M$  has a coprimary decomposition. Let  $N$  be a non-zero submodule of  $M$  and let  $P_i (1 \leq i \leq k)$  be the coassociated prime ideals of  $N$ . Then there exists a positive integer  $h$  such that  $N = (0 :_N P_1^h)^{P_1} + \cdots + (0 :_N P_k^h)^{P_k}$  is a normal coprimary decomposition of  $N$ .*

**Proof.** Let  $N = N_1 + \cdots + N_k$  be a normal coprimary decomposition of  $N$  where  $k$  is a positive integer and, for each  $1 \leq i \leq k$ ,  $N_i$  is a  $P_i$ -coprimary submodule of  $N$  for some prime ideal  $P_i$  of  $R (1 \leq i \leq k)$ . There exists a positive integer  $h$  such that  $P_i^h N_i = 0 (1 \leq i \leq k)$ . Then, for each  $1 \leq i \leq k$ , we have

$$N_i = N_i^{P_i} \subseteq (0 :_N P_i^h)^{P_i} \subseteq (0 :_N P_i^h) \subseteq N$$

and so

$$P_i^h \subseteq (0 : (0 :_N P_i^h)) \subseteq (0 : (0 :_N P_i^h)^{P_i}) \subseteq (0 : N_i) \subseteq P_i.$$

For each  $1 \leq i \leq k$ ,  $\sqrt{(0 : (0 :_N P_i^h)^{P_i})} = \sqrt{(0 : (0 :_N P_i^h))} = P_i$  and by the hypothesis  $(0 :_N P_i^h)$  has a coprimary decomposition so that  $P_i$  is the only minimal member in the set of coassociated prime ideals of  $(0 :_N P_i^h)$ . Hence by Remark 2.15,  $(0 :_N P_i^h)^{P_i}$  is  $P_i$ -coprimary for each  $1 \leq i \leq n$  so that  $N = (0 :_N P_1^h)^{P_1} + \cdots + (0 :_N P_k^h)^{P_k}$  is a normal coprimary decomposition of  $N$ .  $\square$

Let  $A$  be an ideal of  $R$ . Then  $A$  is said to be left primary if, given any two ideals  $B$  and  $C$  of  $R$  such that  $BC \subseteq A$ , then either  $C \subseteq A$  or  $B^n \subseteq A$  for some positive integer  $n$ . In a similar way we can define right primary.  $A$  is said to be primary if it is both left and right primary. If  $R$  is left and right Noetherian and if  $A$  is a proper ideal of  $R$ , which is primary, then  $P = \sqrt{A}$  is prime such that  $P^n \subseteq A$  for some positive integer  $n$ . In this case,  $A$  is called  $P$ -primary. If  $A$  is a proper ideal of  $R$ , then  $C(A)$  will denote the set of elements  $c$  in  $R$  such that  $c + A$  is a non-zero-divisor in  $R/A$ . Clearly,  $c \in C(A)$  if and only if, for any  $r \in R$ ,  $cr \in A$  or  $rc \in A$  implies  $r \in A$ . The ideal  $A$  of  $R$  is said to be strongly primary if  $A$  is primary and  $C(A) = C(\sqrt{A})$ .

In [6], it is proved that every non-zero injective module over a commutative Noetherian ring has a secondary representation. By a similar method, we prove Theorem 2.28.

**Lemma 2.26.** *Let  $R$  be left and right Noetherian, let  $A$  be a strongly  $P$ -primary ideal of  $R$ , and let  $M$  be an injective  $R$ -module. Then  $N = (0 :_M A)$  is zero or coprimary.*

**Proof.** Suppose  $N$  is a non-zero submodule of  $M$ . Let  $B$  be an ideal of  $R$ . If  $B \subseteq P$ , then  $B^h N \subseteq P^h N \subseteq AN = 0$  for some positive integer  $h$ . Now suppose  $B \not\subseteq P$ . Clearly  $BN \subseteq N$ . Let  $n \in N$ . There is a left  $R$ -module homomorphism  $\varphi : R/A \rightarrow M$  for which  $\varphi(r + A) = rn$  for all  $r \in R$ . Since  $B \not\subseteq P$ ,  $(B + P)/P$  is a non-zero ideal of the prime left and right Noetherian ring  $R/P$ . By Goldie's Theorem, there exists an element  $b \in B \cap C(P)$ . Define a mapping  $\theta : R/A \rightarrow R/A$  by  $\theta(r + A) = rb + A$  for all  $r \in R$ . Since  $b \in B \cap C(P)$  and  $A$  is strongly  $P$ -primary,  $\theta$  is a left  $R$ -module monomorphism. As the diagram

$$\begin{array}{ccc} 0 \rightarrow & R/A & \xrightarrow{\theta} R/A \\ & \varphi \downarrow & \\ & & M \end{array}$$

has exact row, it can be completed with a left  $R$ -module homomorphism  $\psi : R/A \rightarrow M$  which makes the extended diagram commute. Thus  $n = \varphi(1 + A) = \psi\theta(1 + A) = \psi(b + A) = b\psi(1 + A)$ . Since  $\psi(1 + A) \in N, n \in bN \subseteq BN$ . We have shown that  $N \subseteq BN$ . The result follows.  $\square$

For the proof of the next result see [6, Lemma 2.2].

**Lemma 2.27.** *Let  $M$  be injective, let  $n$  be a positive integer, and let  $A_i(1 \leq i \leq n)$  be ideals of  $R$ . Then  $\sum_{i=1}^n (0 :_M A_i) = (0 :_M \bigcap_{i=1}^n A_i)$ .*

**Theorem 2.28.** *Let  $R$  be left and right Noetherian such that the zero ideal of  $R$  is a finite intersection of strongly primary ideals. Then every non-zero injective  $R$ -module has a coprimary decomposition.*

**Proof.** Let  $n$  be a positive integer and let  $A_i(1 \leq i \leq n)$  be strongly primary ideals of  $R$  such that  $0 = \bigcap_{i=1}^n A_i$ . Let  $M$  be a non-zero injective  $R$ -module. Then  $M = (0 :_M 0) = (0 :_M \bigcap_{i=1}^n A_i) = \sum_{i=1}^n (0 :_M A_i)$ , where  $(0 :_M A_i)$  is zero or coprimary for all  $1 \leq i \leq n$ .  $\square$

Note that the condition on  $R$  in Theorem 2.28 is satisfied if  $R$  is the universal enveloping algebra of a finite-dimensional nilpotent Lie algebra (see [1, p.78] and [4]).

## References

- [1] Chatters, A. W. and Hajarnavis, C. R.: Non-commutative rings with primary decomposition, Quart. J. Math. Oxford (2) 22, 73-83 (1971).
- [2] Hungerford, T. W.: *Algebra*, Springer-Verlag, New York, 1974.
- [3] Krause, G.: On fully left bounded left Noetherian rings, J. Algebra 23 (1), 88-99 (1972).
- [4] McConnell, J. C.: The intersection theorem for a class of non-commutative rings, Proc. London Math. Soc. 17 (3), 487-498 (1967).
- [5] Mogami, I. and Tominaga, H.: On coprimary decomposition theory for modules, Math. J. Okayama Univ. 17 (2), 125-130 (1975).

MAANI SHIRAZI, SMITH

- [6] Sharp, R. Y.: Secondary representations for injective modules over commutative Noetherian rings, Proc. Edinburgh Math. Soc. (2) 20 (2), 143-151 (1976).
- [7] Smith, P. F.: Uniqueness of primary decompositions, Turkish J. Math. 27 (3), 425-434 (2003).

M. MAANI-SHIRAZI

Department of Mathematics,  
College of Sciences, Shiraz University,  
Shiraz 71454, IRAN  
e-mail: liliumsor7@yahoo.com

P. F. SMITH

Department of Mathematics,  
University of Glasgow,  
Glasgow, G12 8QW, Scotland, UK  
e-mail: pfs@maths.gla.ac.uk

Received 07.07.2005