

1-1-2005

Pyranocoumarins from *Seseli gummiferum* subsp. *corymbosum* Growing in Turkey

ALEV TOSUN

NAZİRE ÖZKAL

MASAKI BABA

TORU OKUYAMA

Follow this and additional works at: <https://journals.tubitak.gov.tr/chem>

 Part of the [Chemistry Commons](#)

Recommended Citation

TOSUN, ALEV; ÖZKAL, NAZİRE; BABA, MASAKI; and OKUYAMA, TORU (2005) "Pyranocoumarins from *Seseli gummiferum* subsp. *corymbosum* Growing in Turkey," *Turkish Journal of Chemistry*. Vol. 29: No. 3, Article 13. Available at: <https://journals.tubitak.gov.tr/chem/vol29/iss3/13>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Chemistry by an authorized editor of TÜBİTAK Academic Journals. For more information, please contact academic.publications@tubitak.gov.tr.

Pyranocoumarins from *Seseli gummiferum* subsp. *corymbosum* Growing in Turkey

Alev TOSUN¹, Nazire ÖZKAL¹, Masaki BABA², Toru OKUYAMA^{2*}

¹Ankara University, Faculty of Pharmacy, Department of Pharmacognosy,
06100 Tandoğan, Ankara-TURKEY

²Meiji Pharmaceutical University, Department of Natural Medicine and Phytochemistry
2-522-1, Noshio, Kiyose-shi, Tokyo, 204-8588-JAPAN
e-mail: okuyama@my-pharm.ac.jp

Received 25.11.2004

Seseli gummiferum Pall. ex Sm. subsp. *corymbosum* (Boiss. & Heldr.) P.H. Davis (Syn: *S. corymbosum* Boiss. & Heldr.) (Umbelliferae) collected in southern Anatolia was investigated for the presence of coumarins. A new angular-type pyranocoumarin, corymbocoumarin (**1**), along with 5 known coumarins (**2-6**) were isolated from the aerial parts of this plant. Corymbocoumarin (**1**) was established to be (-)-(3',4'S)-3'-acetoxy-4'-isovaleryloxy-3',4'-dihydroseselin (**1**) and coumarins **2-6** were identified as (-)-(3',4'S)-3'-acetoxy-4'-angeloyloxy-3',4'-dihydroseselin (**2**), (+)-(3',4'S)-3'-hydroxy-4'-angeloyloxy-3',4'-dihydroseselin (d-laserpitin) (**3**), (-)-(3',4'S)-3'-angeloyloxy-4'-hydroxy-3',4'-dihydroseselin (**4**), 3'-acetoxy-4'-isobutyloxy-3',4'-dihydroseselin (**5**) and osthole (**6**), respectively, by spectroscopic methods. The structural elucidation and absolute configurations were determined by chemical correlations with known compounds.

Key Words: Umbelliferae, corymbocoumarin, angular-type pyranocoumarin, *Seseli gummiferum* Pall. ex Sm. subsp. *corymbosum* (Boiss. & Heldr.) P.H. Davis

Introduction

The genus *Seseli* L. is represented by 12 taxa (11 species and 1 subspecies) in the *Flora of Turkey*, of which 4, including the title species, are native to the region. *S. gummiferum* subsp. *corymbosum* is a perennial or monocarpic plant growing in southern Anatolia, Turkey¹⁻⁴. There is no record of the chemical constituents of this plant.

This paper describes the isolation of a new angular-type dihydropyranocoumarin, corymbocoumarin (**1**), along with 4 known angular-type pyranocoumarins (**2-5**) and a simple coumarin (**6**), all isolated from the *n*-hexane extract of the aerial parts of *S. gummiferum* subsp. *corymbosum*.

*Corresponding author

Experimental

General experimental procedures

Chromatographic separations were carried out on a silica gel open column (0.063-0.200 mm, 70-230 mesh Merck 1.07734). TLC (Silica gel 60 F₂₅₄, Merck 1.05554) was observed under UV at 366 nm and 254 nm, and then sprayed with sulfuric acid and 5% KOH solution. Prep. HPLC was performed on a TOSOH-Prep-HPLC equipped with an HPLC packed column of Senshu-Pak silica 4251-N (250 mm x 10 mm) and TOSOH UV-8010, 320 nm. Melting points were measured on a BUCHI SMP-20, electrothermal melting point apparatus and a YANACO micro melting point apparatus. Optical rotations were determined on a JASCO DIP-140 automatic polarimeter at 20 °C. ¹H and ¹³C NMR spectra were measured in CDCl₃, on a JEOL JNM-EX 270 FT-NMR spectrometer. Chemical shifts are expressed in δ units relative to TMS ($\delta = 0$) as internal standard. Mass spectra were recorded on a JEOL-JMS DX 302 MS spectrometer operating at 70 eV in electron impact mode.

Plant material

Aerial parts of *Seseli gummiferum* Pall. ex Sm. subsp. *corymbosum* (Boiss. & Heldr.) P.H. Davis (Umbelliferae) were collected from the Village of Pınarbaşı, Kadife Mountains, Akseki-Antalya, at an altitude of 1650-1900 m, during the flowering season of August, 2000. A voucher specimen identified by Prof. Hayri Duman is deposited in the Herbarium of Ankara University's Faculty of Pharmacy (AEF 21701).

Extraction and Isolation

Air-dried aerial parts (1 kg) of *S. gummiferum* subsp. *corymbosum* were successively extracted with *n*-hexane, Et₂O, EtOAc and MeOH in a Soxhlet apparatus over 8 h. The organic solvents were evaporated to dryness in vacuo to yield the corresponding *n*-hexane (85 g), Et₂O (31.66 g), EtOAc (13.14 g) and MeOH extracts (110 g), respectively. The *n*-hexane extract was chromatographed on a silica gel column using an *n*-hexane:EtOAc gradient elution system.

The new coumarin, corymbocoumarin (**1**) (1.5 g), and coumarin **6** (3.2 g) were isolated from the fractions eluted with *n*-hexane:EtOAc (9:1). The fractions obtained by elution with *n*-hexane:EtOAc (8:2) were purified by HPLC eluted with *n*-hexane:EtOAc (3:1), to give coumarins **2** (20.5 mg), **3** (77.8 mg), **4** (48.3 mg) and **5** (10.2 mg).

Coumarin 2 (2): white oil, $[\alpha]_D - 36.45^\circ$ (CHCl₃); EIMS: C₂₁H₂₂O₇ (M⁺, m/z: 386; 13.07%), 326 (1.80), 311 (5.45), 303 (1.70), 286 (9.25), 244 (40.24), 229 (89.47), 191 (19.72), 83 (base peak, 100), 55 (50.31), 43 (37.11); ¹H NMR (270 MHz, in CDCl₃, δ ppm; see Table 1); ¹³C NMR (see Table 2).

Coumarin 3 (3): white oil, $[\alpha]_D + 84.61^\circ$ (CHCl₃). EIMS: C₁₉H₂₀O₆ (M⁺, m/z: 344; 18.99%), 326 (7.64), 311 (20.60), 245 (26.58), 244 (21.16), 229 (35.17), 203 (40.57), 83 (base peak, 100), 55 (39.37); ¹H NMR (270 MHz, in CDCl₃, δ ppm; see Table 1); ¹³C NMR (see Table 2).

Coumarin 4 (4): glass-like substance, m.p. 71 °C, $[\alpha]_D - 38.84^\circ$ (CHCl₃). EIMS: C₁₉H₂₀O₆ (M⁺, m/z: 344; 10.91%), 229 (31.22), 83 (base peak, 100), 71 (18.34), 55 (34.62), 43 (27.26); ¹H NMR (270 MHz, in CDCl₃, δ ppm; see Table 1); ¹³C NMR (see Table 2).

Coumarin 5 (5): glass - like substance. EIMS: C₁₉H₂₂O₇ (M⁺, m/z: 374; 12.18%), 314 (8.62), 299 (13.83), 229 (base peak, 100), 71 (16.68), 43 (16.14); ¹H NMR (270 MHz, in CDCl₃, δ ppm; see Table 1); ¹³C NMR (see Table 2).

Coumarin 6 (6): colorless needles, m.p. 83 - 84 °C, C₁₅H₁₆O₃ (M⁺, m/z: 244; 100%), 229 (72.7), 213 (25.94), 201 (32.50), 189 (44.53); ¹H NMR (270 MHz, in CDCl₃, δ ppm): 6.23 (1H, d, *J* = 9.6 Hz, H - 3), 7.61 (1H, d, *J* = 9.6 Hz, H - 4), 7.29 (1H, d, *J* = 8.6 Hz, H - 5), 6.83 (1H, d, *J* = 8.6 Hz, H - 6), 3.53 (2H, d, *J* = 7.3 Hz, H - 1'), 5.22 (1H, t - like, *J* = 7.3 Hz, H - 2'), 1.66 (3H, s, H - 4'), 1.84 (3H, s, H - 5'), 3.92 (3H, s, - OCH₃).

Table 1. ¹H NMR spectral data of *Seseli* pyranocoumarins **1-5** (ppm from TMS, in CDCl₃).



Proton	1	2	3	4	5
H-3	6.23 (d, <i>J</i> = 9.6)	6.24 (d, <i>J</i> = 9.6)	6.20 (d, <i>J</i> = 9.2)	6.25 (d, <i>J</i> = 9.6)	6.23 (d, <i>J</i> = 9.6)
H-4	7.59 (d, <i>J</i> = 9.6)	7.61 (d, <i>J</i> = 9.6)	7.60 (d, <i>J</i> = 9.2)	7.65 (d, <i>J</i> = 9.6)	7.60 (d, <i>J</i> = 9.6)
H-5	7.36 (d, <i>J</i> = 8.6)	7.36 (d, <i>J</i> = 8.6)	7.35 (d, <i>J</i> = 8.6)	7.34 (d, <i>J</i> = 8.9)	7.36 (d, <i>J</i> = 8.6)
H-6	6.80 (d, <i>J</i> = 8.6)	6.81 (d, <i>J</i> = 8.6)	6.80 (d, <i>J</i> = 8.6)	6.80 (d, <i>J</i> = 8.6)	6.81 (d, <i>J</i> = 8.6)
H-3'	5.31 (d, <i>J</i> = 5.0)	5.41 (d, <i>J</i> = 4.6)	4.10 (d, <i>J</i> = 3.0)	5.47 (br. m)	5.32 (d, <i>J</i> = 5.0)
H-4'	6.55 (d, <i>J</i> = 4.6)	6.60 (d, <i>J</i> = 5.0)	6.51 (d, <i>J</i> = 5.0)	5.22 (d, <i>J</i> = 5.0)	6.53 (d, <i>J</i> = 5.0)
gem-Me	1.45, 1.42	1.48, 1.43	1.50, 1.44	1.51, 1.44	1.45, 1.42
Ester part	2.10 (3H s, OCOCH ₃)	2.11 (3H s, OCOCH ₃)	3.14 (1H m, OH)	3.41 (1H br.s, OH)	2.09 (3H s, OCOCH ₃)
	2.24-2.30 (2H m)	6.14 (1H m)	6.08 (1H m)	6.15 (1H m)	2.61 (1H m)
	2.15 (1H t-like, <i>J</i> = 6.6)	1.96 (3H dd, <i>J</i> = 7.3, 1.7)	1.99 (3H d, <i>J</i> = 7.3)	1.99 (3H d, <i>J</i> = 7.3)	1.20 (3H d, <i>J</i> = 6.9)
	1.00, 0.98 (each 3H d, <i>J</i> = 6.3)	1.87 (3H t-like, <i>J</i> = 1.7)	1.89 (3H t-like, <i>J</i> = 1.7)	1.94 (3H d-like)	1.23 (3H d, <i>J</i> = 6.9)

Table 2. ^{13}C NMR spectral data of *Seseli* pyranocoumarins **1-5** (ppm from TMS, in CDCl_3).

Carbon	1	2	3	4	5
2	159.71	159.87	159.87	160.61	159.72
3	113.26	113.10	112.85	112.51	113.30
4	143.14	143.27	143.27	143.90	143.20
5	129.24	129.11	129.20	128.68	129.23
6	114.38	114.30	114.47	114.52	114.43
7	156.57	156.69	156.89	155.96	156.64
8	107.19	106.99	107.19	110.82	107.22
4a	112.49	112.49	112.17	112.36	112.54
8a	153.96	153.94	154.12	154.30	154.09
2'	77.84	77.65	78.53	77.58	77.31
3'	70.55	69.74	71.43	72.45	70.55
4'	60.40	60.97	63.20	59.86	60.54
<i>gem</i> -Me	25.34	24.85	25.68	25.70	25.34
	22.14	22.90	20.79	22.52	22.32
Ester part					
CO	169.81	169.76	—	—	169.86
OCOCH₃	20.69	20.60	—	—	20.74
others	171.93	166.41	168.09	166.90	175.85
	43.25	139.71	138.72	139.43	34.14
	25.50	126.92	127.37	127.19	18.94
	22.43	20.43	20.34	20.54	18.83
	22.39	15.71	15.65	15.78	—

Results and Discussion

The coumarins **2-6** were identified as (–)-(*3'S,4'S*)-3'-acetoxy-4'-angeloyloxy-3',4'-dihydroseselin (**2**), (+)-(*3'S,4'S*)-3'-hydroxy-4'-angeloyloxy-3',4'-dihydroseselin (d-laserpitin) (**3**), (–)-(*3'S,4'S*)-3'-angeloyloxy-4'-hydroxy-3',4'-dihydroseselin (**4**), 3'-acetoxy-4'-isobutyloxy-3',4'-dihydroseselin (**5**), and osthole (**6**), by analysis of physical and spectroscopic data and comparison with previously published values^{5–11}.



	R₁	R₂
1	COCH ₃	COCH ₂ CH(CH ₃) ₂
2	COCH ₃	COC(CH ₃)=CHCH ₃
3	H	COC(CH ₃)=CHCH ₃
4	COC(CH ₃)=CHCH ₃	H
5	COCH ₃	COCH(CH ₃) ₂

6 Osthole

Structure determination of corymbocoumarin (1)

Corymbocoumarin (**1**), colorless needles; m.p. 105.0-108.0 °C, $[\alpha]_D -6.502^\circ$ (CHCl₃), showed a molecular ion peak at m/z 388 (EIMS: C₂₁H₂₄O₇). The mass spectrum of **1** exhibits a fragmentation pattern almost identical with that of **2**. The ¹H NMR spectrum of **1** shows 2 methyl doublets at δ 0.98 and 1.00 ($J=6.3$), *gem* dimethyl singlets at δ 1.42 and 1.45, and an acetoxy methyl signal at δ 2.10. The complex signals appearing at δ 2.15 (1H, triplet like) and 2.24-2.30 (2H, multiplet) were assigned to the methine and methylene protons.

In the aromatic proton region, 2 pairs of doublets at δ 6.23 and 7.59 ($J=9.6$ Hz) and at δ 6.80 and 7.36 ($J=8.6$ Hz) were attributed to the protons at C₃, C₄, C₆ and C₅, respectively. The doublets at δ 5.31 (1H, $J=5.0$ Hz) and 6.55 (1H, $J=4.6$ Hz) were assigned to the methine protons at C_{3'} and C_{4'} of the *cis*-khellactone diesters, which showed a characteristic splitting pattern. This spectrum is almost identical with that of **2**.

Assigning the ester group positions at C_{3'} and C_{4'} was achieved with mass spectrometry. Based on mass fragmentograms on seselin-type coumarins proposed by Bohlmann et al.¹², Takata et al.¹³ were able to analyze the fragmentation patterns of peucedanocoumarins I and II isolated from the crude Chinese drug, "Baihua Qianhu". Similarly, the mass spectral fragmentation pattern of **1** was seen at m/z 328 (7.5%) and 313 (4.3%) for the loss of an acetoxy group, and m/z 287 (8.3%) for loss of an isovaleryloxy group (Figure 1). Hence, we conclude that **1** is a khellactone diester with an acetoxy and an isovaleryloxy group.

As previously described¹⁴, **1** was subjected to alkaline hydrolysis with 1N KOH to yield (-) *cis* and (+) *trans*-khellactones. Alkaline hydrolysis of **1**, with 0.5 N KOH at reflux for 5 min, yielded *cis*-3'-acetoxy-4'-hydroxy-3',4'-dihydro-seselin. This product is also obtained from **2** under the same conditions (Figure 2).

The configuration at C_{3'} and C_{4'} of **1** was determined by the difference in the separation pattern of the 2 geminal methyl carbon signals in the ¹³C NMR spectrum. Gonzalez et al. determined that the 2'-*gem*-dimethyl groups of dihydropyran give a broad singlet in 3',4'-*trans* compounds and 2 close singlets in 3',4'-*cis* compounds⁵. Moreover, previously, the ¹H NMR signals appearing at δ 1.42 and 1.45 (Δ 0.03) in **1** and at δ 1.43 and 1.48 (Δ 0.05) in **2** indicated *cis* configurations at C_{3'} and C_{4'}, respectively¹⁵.




Figure 1. The MS fragmentation pattern of **1**.




Figure 2. Alkaline hydrolysis of **1** and **2**.

Thus, a new angular-type dihydropyranocoumarin, corymbocoumarin, was determined as $(-)$ - $(3'S,4'S)$ -3'-acetoxy-4'-isovaleryloxy-3', 4'-dihydroreseselin (**1**).

Acknowledgment

We would like to thank "Meiji Pharmaceutical University-Onda Scholarship" for its support of this study.

References

1. I.C. Hedge and J.M. Lamond, "*Seseli* L." in Flora of Turkey and the East Aegean Islands, Vol. 4, ed. P.H. Davis, Edinburgh University Press, Edinburgh, 1972.
2. P.H. Davis, R.R. Mill and K. Tan, "*Flora of Turkey and the East Aegean Islands*", Vol. 10, Edinburgh University Press, Edinburgh, 1988.
3. H. Duman, "*Seseli* L." in Flora of Turkey and the East Aegean Islands, Vol. 11, eds. A. Güner, N. Özhatay, T. Ekim, K.H.C. Başer, Edinburgh University Press, Edinburgh, 2000.
4. G. Parolly and B. Nordt, *Willdenowia* **31**, 87-93 (2001).
5. A.G. Gonzales, J.T. Barroso, H.L. Dorta, J.R. Luis and F. Rodriguez-Luis, *Phytochemistry* **18**, 1021-1023 (1979).
6. R.D.H. Murray, J. Mendez and S.A. Brown, "*The Natural Coumarins*", John Wiley & Sons Ltd., New York, 1982.
7. T.M. Swager and J.H. Cardellina, II., *Phytochemistry* **24**, 805-813 (1985).
8. A. Bellino, P. Venturella, M.L. Marino, O. Servettaz and G. Venturella, *Phytochemistry* **25**, 1195-1199 (1986).
9. S. Yamaguchi, "*Studies on the Component of Turkish Umbelliferous Plants and Japanese Sea Weeds*" Master Thesis of Meiji Pharmaceutical University, No. 431, Tokyo, Japan, 1993.

Pyranocoumarins from *Seseli gummiferum* subsp. *corymbosum* Growing in Turkey..., A. TOSUN, et al.,

10. Y. Ikeshiro, I. Mase and Y. Tomita, **Phytochemistry** **33**, 1543-1545 (1993).
11. B. Fan, M. Baba, A. Mizuno, Y. Okada, J. Xu and T. Okuyama, **J. Japanese Botany** **75**, 257-261 (2000).
12. F. Bohlmann, V.S. Bhaskar Rao and M. Grenz, **Tetrahedron Lett.** **36**, 3947 - 3950 (1968).
13. M. Takata, S. Shibata and T. Okuyama, **Planta Med.** **56**, 307-311 (1990).
14. T. Okuyama and S. Shibata, 1981. **Planta Med.** **42**, 89-86 (1981).
15. S. Shibata and T. Okuyama, **Abstracts of Chinese Medicines** **3**, 214-230 (1989).