

1-1-2006

Response of the Pineal Gland in the Rats Exposed to Three Different Light Spectra of Short Periods

ERİNÇ ARAL

SEMA USLU

EMİNE SUNAL

AYLA EKER SARIBOYACI

İMER OKAR

See next page for additional authors

Follow this and additional works at: <https://journals.tubitak.gov.tr/veterinary>



Part of the [Animal Sciences Commons](#), and the [Veterinary Medicine Commons](#)

Recommended Citation

ARAL, ERİNÇ; USLU, SEMA; SUNAL, EMİNE; SARIBOYACI, AYLAKER; OKAR, İMER; and ARAL, ERTUNÇ (2006) "Response of the Pineal Gland in the Rats Exposed to Three Different Light Spectra of Short Periods," *Turkish Journal of Veterinary & Animal Sciences*: Vol. 30: No. 1, Article 4. Available at: <https://journals.tubitak.gov.tr/veterinary/vol30/iss1/4>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Veterinary & Animal Sciences by an authorized editor of TÜBİTAK Academic Journals. For more information, please contact academic.publications@tubitak.gov.tr.

Response of the Pineal Gland in the Rats Exposed to Three Different Light Spectra of Short Periods

Authors

ERİNÇ ARAL, SEMA USLU, EMİNE SUNAL, AYLA EKER SARIBOYACI, İMER OKAR, and ERTUNÇ ARAL

Response of the Pineal Gland in Rats Exposed to Three Different Light Spectra of Short Periods

Erinç ARAL^{1*}, Sema USLU², Emine SUNAL², Ayla EKER SARIBOYACI¹, İmer OKAR³, Ertunç ARAL⁴

¹Department of Histology-Embryology, Faculty of Medicine, Osmangazi University, Eskisehir - TURKEY

²Department of Biochemistry, Faculty of Medicine, Osmangazi University, Eskisehir - TURKEY

³Department of Histology-Embryology, Faculty of Medicine, Marmara University, İstanbul - TURKEY

⁴Department of Physics, Faculty of Arts and Sciences, Osmangazi University, Eskisehir - TURKEY

Received: 19.06.2003

Abstract: The pineal gland is a part of the photo-endocrine system. Photon energy is important for the function of this system, and affects the level of nocturnal melatonin. In mammals light-induced suppression of melatonin production is dependent on the intensity and wavelength of the light used. We studied the chronic effects of light wavelengths on the pineal gland in rats after exposure to a short photoperiod using 3 different light spectra with the same irradiance. Male Sprague-Dawley rats were used. The animals were divided into 5 groups. They were exposed 8/16 L:D periods in rooms under automatically regulated light and climate. Lee filters were used for the light spectra application. From animals at the age of 6 months blood samples were taken over a week at the third hour of the dark period under dim red light. The pineal glands were processed for electron microscopy. Melatonin levels in the blood were analyzed by ELISA. Melatonin levels were the higher in control group and group 5 (blue light spectra) compared to the others. We demonstrated that blue light spectra have a greater effect on melatonin production in rats.

Key Words: Light spectra, pineal gland, melatonin, short photoperiod

Kısa Fotoperiyotta Üç Farklı Işık Dalga Boyuna Maruz Bırakılmış Sıçanlarda Epifiz Yanıtı

Özet: Epifiz, foto-endokrin sistemin bir parçasıdır. Foton enerjisi bu sistemin fonksiyonu üzerinde çok önemlidir ve günlük melatonin üretimini etkiler. Memelilerde melatonin üretimi ışığın şiddeti ve dalga boyu etkisindedir. Yapılan bu çalışmada erkek Sprague-Dawley türü sıçanlarda kısa fotoperiyotta (kış dönemi) ve sabit ışık şiddetinde üç farklı dalga boyunun, uzun süre kullanımda epifiz üzerindeki etkisi araştırılmıştır. Hayvanlar beş gruba ayrılarak 8 saat aydınlık ve 16 saat karanlık olmak üzere ışığı ve ısıyı otomatik olarak ayarlanan odalarda büyütülmüşlerdir. Dalga boyları için Lee filtreler kullanılmıştır. Altı aylık yaşa ulaşan hayvanlardan karanlık periyodun 3. saatinde 3 lükslük kırmızı ışık altında bir hafta süre ile kan örnekleri alınmıştır. Kandaki melatonin miktarları ELISA metoduyla ölçülmüştür. Epifizler elektron mikroskopik gözlemler için hazırlanmıştır. Sonuçlarımızda kandaki melatonin değerleri, mavi dalga boyu uygulanan grupta ve kontrol grubunda diğer gruplara oranla yüksek bulunmuştur. Bu nedenle sıçanlarda melatonin üretimi üzerine mavi dalga boyu etkisinin diğer dalga boylarına göre daha fazla olduğunu düşünmekteyiz.

Anahtar Sözcükler: Işık spektrumu, epifiz, melatonin, kısa fotoperiyod

Introduction

The pineal gland is a circumventricular organ situated outside the blood-brain barrier. It is innervated by the fibers of the superior cervical ganglion (1). The gland synthesized melatonin hormone has very different effects on the organism (2). The pineal gland is a part of the photo-endocrine system. Photon energy is important for

the function of this system, and affects the level of nocturnal melatonin (3,4). The most important effect of melatonin synthesized by the pineal gland is to have a diurnal rhythm influenced by light and dark periods (5,6). The blood concentration of melatonin increases in the dark period of the circadian rhythm and is inhibited by the light (7,8). Dark and light environments are

* E-mail: erinca@ogu.edu.tr

differentiated through the eyes. Color sensation is determined by the wavelength and frequency of visible light, which is defined as an interval, not as color. All the wavelengths defining the color spectrum are present in daylight and their preference for the animal experiments have been determined for albino animal species with photopic (black/white) and scotopic (colored vision) electro-retinograms (9). The investigated animals perceive the light with their eyes and it reaches the hypothalamic suprachiasmatic nucleus via the retino-hypothalamic nerves. The signals are then conducted through the paraventricular nucleus, anterior median cerebral cluster, and reticular formation and medulla spinalis to the superior cervical ganglion. Impulses reaching the ganglion increase or decrease in response to the activity of the pineal gland. The synthesis of serotonin or specific methyl indoles begins or stops via the above mechanism (10,11).

All animal species living in the regions from the equator to the poles have fertile phases where light is the critical environmental factor regulating the seasonal model of coupling. The differences are regulated according to the seasons of summer and winter, and defined long and short photoperiods of days are initiated by setting up constant light arrangements. The synthesis and secretion of melatonin increase in winter months, i.e. in the short photoperiod, and decrease in summer months, i.e. in the long photoperiod (3).

The effect of photoperiod on the metabolism of melatonin and effects of light intensity and wavelengths on photo-periodical changes have been investigated for many species (12-14). The results determining the pineal gland activity are influenced by the light intensity and wavelengths in different ways with respect to the species but at the same level (11,15,16). Melatonin levels in various body fluids have been measured with different techniques (17). In recent years the morphology of the mammalian pineal gland has been extensively studied (1,2). Some investigators demonstrated that pineocytes showed ultrastructural features, indicating an increased metabolic and synthetic activity of these cells in some species (18-21). The aim of this study was to investigate the long-time effect of light wavelengths on melatonin production and pineocytes ultrastructure using 3 different light spectra (I max = 670 nm red, I max = 570 nm yellow, I max = 450 nm blue bandwidths).

Materials and Methods

Three generation male Sprague-Dawley rats weighing about 230 ± 250 g were housed in clear polycarbonate cages in a windowless room. The cages were screened by metal lids, which caused no selective interference with the light transmission from overhead luminaries. The rats were kept in an air-conditioned room. Room temperature (22 ± 2 °C) and humidity were automatically regulated. Food and water were supplied ad libitum (12,22). The animals were treated in accordance with the Guidelines for Animal Care and Use of the National Institutes of Health.

The animals were divided into 5 groups. The beginning of the gestation period was marked as day 0. The first born male rats were used in the experiment.

All new-born rats were isolated from their parents at the age of 1 month and subjected to the above protocol of illumination until the age of 6 months.

Group 1 (n = 10): Control (8 h of light: 16 of darkness)

Group 2 (n = 10): Dark (continuous dark period)

Group 3 (n = 10): Red light (Lee filter no: 182)

Group 4 (n = 10): Yellow light (Lee filter no: 101)

Group 5 (n = 10): Blue light (Lee filter no: 183)

Experimental design

The rats were exposed to fluorescent light (36 W) sources of 2500 lux during the experiment. Intensity was measured with a luxmeter (Foot Candle/ Extech instruments) inside the cages. Each cage was well ventilated with fans and coated with special filters. The distribution of the spectral power of the light sources was determined with a Lee filter (Figure 1). Ten animals were housed within each cage and adapted to an 8:16 h light:dark cycle (lights on from 08:00 to 16:00 daily) (15).

Determination of melatonin

When the animals had reached the age of 6 months, blood samples were taken over a week at the third hour of the dark period. At the end of each week, the animals were anaesthetized and intracardiac blood samples and tissues were taken under dim light during the dark period. Plasma samples were stored frozen until the time of assays (12). Melatonin levels were analyzed with ELISA (IBL-Hamburg, Germany) (17).




Figure 1. Lee filters used in the experiments.

The pineal glands were processed for electron microscopy (23). The examination was performed in a JEOL JEM 1220 electron microscope.

Statistical analysis

Melatonin levels of animals were compared by one-way ANOVA.

Results

Biochemical Results

We found some differences between the groups. The melatonin blood level was higher in the control group and group 5 than in the others (Table).

There were significant differences between the groups ($F_{4; 43} = 7.98 P < 0.01$). There were differences between group 1 and groups 2, 3 and 4 and between group 2 and group 5 with respect to melatonin level measurements ($P < 0.01$), groups of 3, 4 and 5 showed no differences. ($P > 0.05$).

Histology

We observed a homogeneous population of pinealocytes in the pineal gland of this species. The Golgi apparatus, rough endoplasmic reticulum, mitochondria, lysosomes, clear vesicles, vacuoles containing flocculent material, microtubules and lipid droplets were consistent with the components of the pinealocytes cytoplasm: infrequently observed organelles including "synaptic ribbons". Pinealocytes contain a large number of mitochondria of generally moderate electron density in groups.

Figure 2 shows lipid droplets and mitochondria in the pinealocyte cytoplasm in group 3 (under growing red light spectra). Figure 3 shows in group 4 (under growing yellow light spectra) 2 pinealocyte irregularly shaped nuclei with prominent nucleoluses and small, peripheral clumps of chromatine. Figure 4 shows in group 5 (under growing blue light spectra) numerous mitochondria in pinealocyte cytoplasm.

Table. Melatonin values (µg/ml) in the blood.

Groups	Number of Rats	Mean and Standard Error
1	7	48.00 ± 6.29*
2	12	22.92 ± 3.04
3	11	27.91 ± 2.97
4	8	21.88 ± 2.42
5	10	39.30 ± 4.17*
Total	48	30.95 ± 2.10

Significance level * = $P < 0.01$




Figure 2. In group 3 (under growing red light spectra) lipid droplets and mitochondria (→) in pinealocyte cytoplasm. X 12.000




Figure 3. In group 4 (under growing yellow light spectra) two pinealocytes irregularly shaped nuclei have prominent nucleoluses and small, peripheral clumps of chromatine. X 5.000




Figure 4. In group 5 (under growing blue light spectra) numerous mitochondria in pinealocytes cytoplasm. X 5.000

Discussion

In most mammalian species, the synthesis and release of melatonin increase at night and decrease with light. Light is the most important environmental factor regulating the melatonin generating system in the gland (3). In earlier studies light of wavelength has been shortly used at wavelengths 510-550 nm, 558-636 nm, 653-668 nm during the period of peak pineal melatonin production, (15) and at 435 nm, 530 nm, 590 nm, 660 nm for 96 hours of exposure to continuous light (22) at different times of light or dark period. The same methods have also been used for studying the effects of light stimuli (7,23,24,). It is reported that the intensity and

wavelengths of light have the same effects the activity of the pineal gland (15,25). Furthermore, it is described that the wavelength of light has various effects that vary with the application of wavelengths and circadian phase (11,16,26). In generally, short and long day photoperiods have been used in experimental studies (3,11). It has been demonstrated that a long day photoperiod leads to dynamic procedures while a short day has contrasting effects. Therefore, in this study the method and duration of wavelengths were different.

In our study, in groups a homogeneous population of pinealocytes was present in the pineal gland of this species. The Golgi apparatus, rough endoplasmic

reticulum, numerous mitochondria, lysosomes, clear vesicles, vacuoles containing flocculent material, microtubules and lipid droplets were consistent with the components of the pinealocytes cytoplasm: infrequently observed organelles including "synaptic ribbons". These data correlated with some studies (27). The mammalian pinealocytes contain a large number of mitochondria, the matrix of which is generally of moderate electron density. Some changes in the mitochondria have been described in mammalian pinealocytes after various experimental procedures. According to some authors mitochondria are somehow involved in serotonin synthesis (3). In our study pinealocytes contain a large number of mitochondria which generally of moderate electron density.

Some studies reported that both serum and pineal melatonin concentrations were significantly higher at night than during the daytime and melatonin level was at its maximum in the third hour of the dark period in circulation. Melatonin has annual, seasonal and monthly rhythms as well as a daily rhythm (3). For this reason in this study all blood samples were taken at the same hour and at the third hour of the dark period.

References

1. Kappers, J. A.: The mammalian pineal gland a survey. *Acta Neurochir.*, 1976; 34: 109-149.
2. Reiter, R.J., Robinson, J.O.: *Melatonin (Your Body's Natural Wonder Drug)*. New York: Bantam Books, 1995; 3-146.
3. Reiter, R.J.: *The Pineal Gland (Anatomy and Biochemistry)*. Chapter 5, 6, Florida: CRC Press, INC. Boca Raton. 1981; 1: 122-169 .
4. Illnerova, H.: Entrainment of mammalian circadian rhythms in melatonin production by light. *Pineal Res. Rev.*, 1988; 6: 173-217.
5. Cassone, V.M.: Effects of melatonin on vertebrate circadian systems. *Trends Neurosci.*, 1990; 13: 457-464.
6. Cassone, V.M., Warren, W.S., Brooks, D.S., Lu, J.: Melatonin, the pineal gland and circadian rhythms. *J. Biol. Rhythms*, 1993; 8 Suppl: S73-81.
7. Honma, S., Kanematsu, N., Katsuno, Y., Honma, K.: Light suppression of nocturnal pineal and plasma melatonin in rats depends on wavelength and time of day. *Neurosci. Lett.*, 1992; 147: 201-204.
8. Underwood, H., Goldman, B.D.: Vertebrate circadian and photoperiodic systems. Role of the pineal and melatonin. *J. Biol. Rhythms.*, 1987; 2: 279-315.
9. Szel, A., Rohlich, P.: The cone types of retina detected by anti-vital pigment antibodies. *Exp. Eye Res.*, 1992; 55: 47-52.
10. Reiter, R.J.: Neuroendocrine effects of light. *Int. J. Biometeorol.*, 1991; 35: 169-175.
11. Brainard, G.C., Vaughan, M.K. Reiter, R.J.: Effect of light irradiance and wavelength on the Syrian hamster reproductive system. *Endocrinology*, 1986; 119: 648-654.
12. Vaughan, M.K., Brainard, G.C., Reiter, R.J.: Photoperiodic and light spectral conditions which inhibit circulating concentrations of thyroxine in the male hamster. *Life Sci.*, 1985; 36: 2183-2188.
13. Jarmak, A., Zawilska, J.B., Nowak, J.: Effect of light with various wavelengths and impulse times on nocturnal suppression of n-acetyltransferase activation by serotonin in the chick. *Klin. Oczna.*, 1996; 98: 417-422.
14. Sinhasane, S.V., Joshi, B.N.: Exposure to different spectra of light continuous light and treatment with melatonin affect reproduction in the Indian desert gerbil *Meriones hurrianae* (Jerdon). *Biol. Signals Recept.*, 1998; 7: 179-187.
15. Brainard, G.C., Richardson, B.A., King, T.S. Reiter, R.J.: The influence of different light spectra on the suppression of pineal melatonin content in the Syrian hamster. *Brain Res.*, 1984; 294: 333-339.

Acknowledgments

This study was supported by the Research Projects Unit of Osmangazi University, Eskişehir.

16. Bertoni, J.M., Sprenkle, P.M., Brainard, G.C.: Effects of different light wavelengths at equal irradiance on testis weight, protein content, and k-paranitrophenylphosphates activity in the Syrian hamster. *Life Sci.*, 1987; 40: 1479-1486.
17. Chenigi, S., Hofmann, B.E., Kaider, A.: Direct enzyme-linked immunosorbent assay and a radioimmunoassay for melatonin compared. *Clin. Chem.*, 1995; 41 :381-386.
18. Dominguez, S., Piezzi, R.S., Scardapane, L., Guzman, J.A.: A light and electron microscopic study of the pineal gland of the viscacha (*Lagostomus maximus maximus*) *J. Pineal Res.*, 1987; 4: 211-219.
19. Calvo, J., Boya J.: Ultrastructure of the pineal gland in the adult rat. *J. Anat.*, 1984; 138: 405-409.
20. Bhatnagar, K.P.: The ultrastructure of mammalian pinealocytes: A systematic investigation. *Microsc. Res. Tech.*, 1992; 21: 85-115.
21. Becker, U.G., Vollrath, L.: 24 hour-variations of pineal gland volume, pinealocyte nuclear volume and mitotic activity in male Spraque-Dawley rats. *J. Neural Transm.*, 1983; 56: 211-221.
22. Cardinali, D.P., Larin, F., Wurtman, J.R.: Control of the rat pineal gland by light spectra. *Proc. Natl. Acad. Sci. USA.*, 1972; 69: 2003-2005.
23. Karasek, M., Marek, K., Pevet, P.: Influence of a short light pulse at night on the ultrastructure of the rat pinealocytes: a quantitative study. *Cell Tissue Res.*, 1988; 254: 247-249.
24. Thiele, G., Holtorf, A., Steinlechner, S. Reiter, R.J.: The influence of different light irradiance on pineal N-acetyltransferase activity and melatonin levels in the cotton rat, *Sigmodon hispidus*. *Life Sci.*, 1983; 33: 1543-1547.
25. van-Benthem, J., Steinen, A.C., Sommer, M.C., De Koning, J., Ebels I., Balemans M.G.: The influence of light of different wavelengths on the methylating capacity of the pineal gland of male golden hamsters in relation to reproduction. *J. Neural Transm.*, 1989; 78: 145-158.
26. Poeggeler, B.H., Barlow-Walden, L.R., Reiter, R.J., Saarela, S., Menendez-Pelaez, A., Yaga, K., Manchester, L.C., Chen, L.D., Tan, D.X.: Red-light-induced suppression of melatonin synthesis is mediated by n-methyl-d-aspartate receptor activation in retinally normal and retinally degenerate rats. *J. Neurobiol.*, 1995; 28:1-8.
27. Karasek, M., Stankov, B., Lucini, V., Scaglione, F., Esposti, G., Mariani, M., Fraschini, F.: Comparison of the rat pinealocyte ultrastructure with melatonin concentrations during daytime and at night. *J. Pineal Res.*, 1990; 9: 251-257.
28. Csernus, V., Becher, P., Mess, B.: Wavelength dependency of light-induced changes in rhythmic melatonin secretion from chicken pineal gland in vitro. *Neuro Endocrinol. Lett.*, 1999; 20: 299-304.
29. Brainard, G.C., Hanifin, J.P., Greeson, J.M., Byrne, B., Glickman, G., Gerner, E., Rollag, M.D.: Action spectrum for melatonin regulation in humans. *J. Neurosci.*, 2001; 21: 6405-6412.
30. Thapan, K., Arendt, J., Skene, D.J.: An action spectrum for melatonin suppression: evidence for a novel non-rod, non-cone photoreceptor system in humans. *J. Physiol.*, 2001; 535: 261-267.