

1-1-2006

Some Random Fixed Point Theorems for Non-Self Nonexpansive Random Operators

POOM KUMAM

SOMYOT PLUBTIENG

Follow this and additional works at: <https://journals.tubitak.gov.tr/math>



Part of the [Mathematics Commons](#)

Recommended Citation

KUMAM, POOM and PLUBTIENG, SOMYOT (2006) "Some Random Fixed Point Theorems for Non-Self Nonexpansive Random Operators," *Turkish Journal of Mathematics*: Vol. 30: No. 4, Article 2. Available at: <https://journals.tubitak.gov.tr/math/vol30/iss4/2>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Mathematics by an authorized editor of TÜBİTAK Academic Journals. For more information, please contact academic.publications@tubitak.gov.tr.

Some Random Fixed Point Theorems for Non-Self Nonexpansive Random Operators*

Poom Kumam and Somyot Plubtieng

Abstract

Let (Ω, Σ) be a measurable space, with Σ a sigma-algebra of subsets of Ω , and let E be a nonempty bounded closed convex and separable subset of a Banach space X , whose characteristic of noncompact convexity is less than 1. We prove that a multivalued nonexpansive, non-self operator $T : \Omega \times E \rightarrow KC(X)$ satisfying an inwardness condition and itself being a $1-\chi$ -contractive nonexpansive mapping has a random fixed point. We also prove that a multivalued nonexpansive, non-self operator $T : \Omega \times E \rightarrow KC(X)$ with a uniformly convex X satisfying an inwardness condition has a random fixed point.

Key Words: Random fixed point, non-self mappings, Nonexpansive random operator, inwardness condition.

1. Introduction

Random fixed point theory has received much attention in recent years; see, Itoh [8] and Shahzad and Latif [15]. Research in this direction was initiated by the Prague School of Probabilists as the originator of random operator theory; see O. Hans [6, 7]. Since then, a lot of efforts have been devoted to random fixed point theory and applications; see Ramírez [11, 12], Tan and Yuan [16], Xu [17, 19, 20], Yuan and Yu [21].

AMS Mathematics Subject Classification: 47H10, 47H09, 47H04.

*This Research was Financially Supported by Faculty of Science, KMUTT.

In 2004, Domínguez Benavides and Ramírez [4] proved a fixed point theorem for a multivalued nonexpansive, non-self mapping and $1-\chi$ -contractive mapping in the framework of Banach spaces whose characteristic of noncompact convexity associated to the separation measure of noncompactness $\varepsilon_\alpha(X)$ is less than 1.

The purpose of the present paper is to prove some random fixed point theorems for nonexpansive non-self random operators. First, we will prove the existence of fixed point for multivalued non-self, nonexpansive random operators in the framework of a Banach spaces with characteristic of noncompact convexity associated to the Kuratowski measure of noncompactness $\varepsilon_\alpha(X)$ being less than 1 and satisfying an inwardness condition, and also being $1-\chi$ -contractive mapping. Moreover, if X is a separable subset of a uniformly convex Banach space, a similar result is proved. Finally we also prove that a multivalued nonexpansive non-self random operator $T : \Omega \times E \rightarrow KC(X)$ satisfying an inwardness condition has a random fixed point.

2. Preliminaries and notations

We begin with establishing some preliminaries. By a measurable space we mean a pair (Ω, Σ) , where Ω is a nonempty set and Σ is a sigma-algebra of subsets of Ω . Let X be a Banach space and E a nonempty subset of X . We shall denote by 2^E the family of nonempty closed subsets of E , by $CB(E)$ the family of nonempty closed bounded subsets of E , by $K(E)$ the family of nonempty compact subsets of E , and by $KC(E)$ the family of nonempty compact convex subsets of E . Let $H(\cdot, \cdot)$ be the Hausdorff distance on $CB(X)$, i.e.,

$$H(A, B) = \max \left\{ \sup_{a \in A} d(a, B), \sup_{b \in B} d(b, A) \right\}$$

for $A, B \in CB(X)$, where $d(x, E) = \inf\{d(x, y) | y \in E\}$ is the distance from x to $E \subset X$.

Let E be a nonempty closed subset of a Banach space X . Recall now that a multivalued mapping $T : E \rightarrow 2^X$ is said to be upper semicontinuous on E if $\{x \in E : Tx \subset V\}$ is open in E whenever $V \subset X$ is open; T is said to be lower semicontinuous if $T^{-1}(V) := \{x \in E : Tx \cap V \neq \emptyset\}$ is open in E whenever $V \subset X$ is open; and T is said to be continuous if it is both upper and lower semicontinuous (cf. [1] and [2] for details). There is another but different kind of continuity for a multivalued operator: $T : E \rightarrow CB(X)$ is said to be continuous on E (with respect to the Hausdorff metric H) if $H(Tx_n, Tx) \rightarrow 0$ whenever $x_n \rightarrow x$. It is not hard to see (see Deimling [2]) that both definitions of continuity are

equivalent if Tx is compact for every $x \in E$.

A multivalued operator $T : \Omega \rightarrow 2^X$ is called (Σ) -measurable if, for any open subset B of X ,

$$T^{-1}(B) := \{\omega \in \Omega : T(\omega) \cap B \neq \emptyset\}$$

belongs to Σ . A mapping $x : \Omega \rightarrow X$ is said to be a *measurable selector* of a measurable multivalued operator $T : \Omega \rightarrow 2^X$ if $x(\cdot)$ is measurable and $x(\omega) \in T(\omega)$ for all $\omega \in \Omega$. An operator $T : \Omega \times E \rightarrow 2^X$ is called a random operator if, for each fixed $x \in E$, the operator $T(\cdot, x) : \Omega \rightarrow 2^X$ is measurable. We will denote by $F(\omega)$ the fixed point set of $T(\omega, \cdot)$, i.e.,

$$F(\omega) := \{x \in E : x \in T(\omega, x)\}.$$

Note that, if we do not assume the existence of a fixed point for the deterministic mapping $T(\omega, \cdot) : E \rightarrow 2^X$, $F(\omega)$ may be empty. A measurable operator $x : \Omega \rightarrow E$ is said to be a *random fixed point of a operator* $T : \Omega \times E \rightarrow 2^X$ if $x(\omega) \in T(\omega, x(\omega))$ for all $\omega \in \Omega$. Recall that $T : \Omega \times E \rightarrow 2^X$ is continuous if, for each fixed $\omega \in \Omega$, the operator $T : (\omega, \cdot) \rightarrow 2^X$ is continuous.

If E is a closed convex subset of a Banach space X , then a multivalued mapping $T : E \rightarrow CB(X)$ is said to be a *contraction* if there exists a constant $k \in [0, 1)$ such that

$$H(Tx, Ty) \leq k\|x - y\|, \quad x, y \in E,$$

and T is said to be *nonexpansive* if

$$H(Tx, Ty) \leq \|x - y\|, \quad x, y \in E,$$

A random operator $T : \Omega \times E \rightarrow 2^X$ is said to be *nonexpansive* if, for each fixed $\omega \in \Omega$ the map $T : (\omega, \cdot) \rightarrow E$ is nonexpansive.

For later reference, we list the following results related to the concept of measurability.

Lemma 2.1 (cf. Wagner [17]) *Let (X, d) be a complete separable metric spaces and $F : \Omega \rightarrow CL(X)$ a measurable map. Then F has a measurable selector.*

Lemma 2.2 (cf. Itoh 1977, [8]) *Suppose $\{T_n\}$ is a sequence of measurable multivalued operator from Ω to $CB(X)$ and $T : \Omega \rightarrow CB(X)$ is an operator. If, for each $\omega \in \Omega$, $H(T_n(\omega), T(\omega)) \rightarrow 0$, then T is measurable.*

Lemma 2.3 (cf. Tan and Yuan [16]) *Let X be a separable metric spaces and Y a metric spaces. If $f : \Omega \times X \rightarrow Y$ is measurable in $\omega \in \Omega$ and continuous in $x \in X$, and if $x : \Omega \rightarrow X$ is measurable, then $f(\cdot, x(\cdot)) : \Omega \rightarrow Y$ is measurable.*

As an easy application of Proposition 3 of Itoh[8] we have the following result.

Lemma 2.4 *Let E be a closed separable subset of a Banach space X , $T : \Omega \times E \rightarrow E$ a random continuous operator and $F : \Omega \rightarrow 2^E$ a measurable closed-valued operator. Then for any $s > 0$, the operator $G : \Omega \rightarrow 2^E$ given by*

$$G(\omega) = \{x \in F(\omega) : \|x - T(\omega, x)\| < s\}, \quad \omega \in \Omega$$

is measurable and so is the operator $cl\{G(\omega)\}$ (the closure of $G(\omega)$).

Lemma 2.5 (cf. Domínguez Benavides and Lopez Acedo [5]) *Suppose E is a weakly closed nonempty separable subset of a Banach space X ; $F : \Omega \rightarrow 2^X$ is a measurable mapping with weakly compact values, and $f : \Omega \times E \rightarrow \mathbb{R}$ is a measurable, continuous and weakly lower semicontinuous function. Then the marginal function $r : \Omega \rightarrow \mathbb{R}$ defined by*

$$r(\omega) := \inf_{x \in F(\omega)} f(\omega, x)$$

and the marginal map $R : \Omega \rightarrow X$ defined by

$$R(\omega) := \{x \in F(\omega) : f(\omega, x) = r(\omega)\}$$

are measurable.

Recall that the Kuratowski and Hausdorff measures of noncompactness of a nonempty bounded subset B of X are defined, respectively, as the numbers

$$\alpha(B) = \inf \{r > 0 : B \text{ can be covered by finitely many sets of diameter } \leq r\},$$

$$\chi(B) = \inf \{r > 0 : B \text{ can be covered by finitely many balls of radius } \leq r\}.$$

The separation measure of noncompactness of a nonempty bounded subset B of X is defined by

$$\beta(B) = \sup \{\varepsilon : \text{there exists a sequence } \{x_n\} \text{ in } B \text{ such that } sep(\{x_n\}) \geq \varepsilon\}.$$

Let X be a Banach space and $\phi = \alpha, \beta$ or χ . The modulus of noncompact convexity associated to ϕ is defined as

$$\Delta_{X,\phi}(\varepsilon) = \inf \{1 - d(0, A) : A \subset B_X \text{ is convex, } \phi(A) \geq \varepsilon\},$$

where B_X is the unit ball of X .

The characteristic of noncompact convexity of X associated with the measure of noncompactness ϕ is defined by

$$\varepsilon_\phi(X) = \sup \{ \varepsilon \geq 0 : \Delta_{X,\phi}(\varepsilon) = 0 \}.$$

The following relationships among the different moduli are easy to obtain:

$$\Delta_{X,\alpha}(\varepsilon) \leq \Delta_{X,\beta}(\varepsilon) \leq \Delta_{X,\chi}(\varepsilon), \tag{2.1}$$

and consequently

$$\varepsilon_\alpha(X) \geq \varepsilon_\beta(X) \geq \varepsilon_\chi(X). \tag{2.2}$$

When X is a reflexive Banach space we have some alternative expressions for the moduli of noncompact convexity associated β and χ :

$$\Delta_{X,\beta}(\varepsilon) = \inf \{ 1 - \|x\| : \{x_n\} \subset B_X, x = w - \lim x_n, \text{sep}(\{x_n\}) \geq \varepsilon \},$$

$$\Delta_{X,\chi}(\varepsilon) = \inf \{ 1 - \|x\| : \{x_n\} \subset B_X, x = w - \lim x_n, \chi(\{x_n\}) \geq \varepsilon \}.$$

In order to study the fixed point theory for non-self mappings, we must introduce some terminology for boundary conditions. The inward set of E at $x \in E$ is defined by

$$I_E(x) := \{ x + \lambda(y - x) : \lambda \geq 0, y \in E \}.$$

Clearly $E \subset I_E(x)$, and it is not hard to show that whenever $I_E(x)$ is a convex set, so is E . A multivalued mapping $T : E \rightarrow 2^X \setminus \{\emptyset\}$ is said to be inward on E if

$$Tx \subset I_E(x) \quad \forall x \in E.$$

Let $\bar{I}_E(x) := x + \{ \lambda(z - x) : z \in E, \lambda \geq 1 \}$. Note that for a convex E , we have $\bar{I}_E(x) = \overline{I_E(x)}$, and T is said to be weakly inward on E if

$$Tx \subset \bar{I}_E(x) \quad \forall x \in E.$$

Let E be a nonempty bounded closed subset of Banach space X and $\{x_n\}$ a bounded sequence in X : we use $r(E, \{x_n\})$ and $A(E, \{x_n\})$ to denote the asymptotic radius and the asymptotic center of $\{x_n\}$ in E , respectively, i.e.

$$r(E, \{x_n\}) = \inf \left\{ \limsup_n \|x_n - x\| : x \in E \right\},$$

$$A(E, \{x_n\}) = \left\{ x \in E : \limsup_n \|x_n - x\| = r(E, \{x_n\}) \right\}.$$

If D is a bounded subset of X , the *Chebyshev radius* of D relative to E is defined by

$$r_E(D) := \inf \{ \sup \{ \|x - y\| : y \in D \} : x \in E \}.$$

Let S be a set $G \subset S$ and D be a directed set, we shall say that a net x_α in S eventually in G if there exist $\alpha_0 \in D$ such that $x_\alpha \in G$ for all $\alpha \geq \alpha_0$.

Definition 2.6 *A net $\{x_\alpha\}$ in a set S is called an ultranet if for each subset $E \subset S$, either $\{x_\alpha\}$ is eventually in E or $\{x_\alpha\}$ is eventually in $S \setminus E$.*

The following facts concerning ultranets can be found in [9]:

- (a) Every net in a set has an ultranet.
- (b) If $f : S_1 \rightarrow S_2$ is a map and if $\{x_\alpha\}$ is an ultranet in S_1 , then $\{f(x_\alpha)\}$ is a ultranet in S_2 .
- (c) If S is compact and $\{x_\alpha\}$ is a ultranet in S , then $\lim_\alpha x_\alpha$ exists.

Obviously, the convexity of E implies that $A(E, \{x_\alpha\})$ is convex. Notice that $A(E, \{x_\alpha\})$ is a nonempty weakly compact set if E is weakly compact, or E is a closed convex subset of a reflexive Banach spaces X .

Let E be a nonempty bounded closed subset of Banach spaces X . Then $\{x_n\} \subset X$ is called *regular* with respect to E if $r(E, \{x_n\}) = r(E, \{x_{n_i}\})$ for all subsequences $\{x_{n_i}\}$ of $\{x_n\}$.

Lemma 2.7 *(Geobel, Lim) Let $\{x_n\}$ and E be as above. Then, there always exists a subsequence of $\{x_n\}$ which is regular with respect to E .*

Moreover, we also need the following Lemma.

Theorem 2.8 (cf. T. D. Benavides and P. L. Ramírez; Theorem 4.3; [4].) *Let E be a closed convex subset of a reflexive Banach space X , and let $\{x_\beta : \beta \in D\}$ be a bounded ultranet. Then*

$$r_E(A(E, x_\beta)) \leq (1 - \Delta_{X,\alpha}(1^-))r(E, \{x_\beta\}). \tag{2.3}$$

The following results are now basic in fixed point theorems for multivalued mappings.

Lemma 2.9 (cf. Deimling 1992, [2]). *Let X be a Banach space and $\emptyset \neq D \subset X$ be a closed bounded convex. Let $F : D \rightarrow 2^X$ be upper semicontinuous γ -condensing with closed convex values, where $\gamma(\cdot) = \alpha(\cdot)$ or $\chi(\cdot)$. If $Fx \cap \overline{I_D(x)} \neq \emptyset$ for all $x \in E$, then F has a fixed point. (Here, $I_D(x)$ is called the inward set at x defined by $I_D := \{x + \lambda(y - x) : \lambda \geq 0, y \in D\}$)*

Proposition 2.10 (cf. Kirk-Massa Theorem [10]) *Let E be a nonempty weakly compact separable subset of a Banach space X , $T : E \rightarrow KC(E)$ a nonexpansive mapping, and $\{x_n\}$ a sequence in E such that $\lim_n d(x_n - Tx_n) = 0$. Then, there exists a subsequence $\{z_n\}$ of $\{x_n\}$ such that*

$$Tx \cap A \neq \emptyset, \forall x \in A := A(E, \{z_n\})$$

3. The Main Results

The following states the main result of this paper, and is the random version of theorem 3.4 of Domínguez Benavides and Lorenzo Ramírez ([4]).

Theorem 3.1 *Let E be a nonempty closed bounded, convex and separable subset of a Banach space X such that $\epsilon_\alpha(X) < 1$, and $T : \Omega \times E \rightarrow KC(X)$ be a nonexpansive random operator and 1 - χ -contractive mapping, such that for each $\omega \in \Omega$, $T(\omega, E)$ is a bounded set, which satisfies the inwardness condition. Then T has a random fixed point.*

Proof. For each $\omega \in \Omega$, and for every $n \geq 1$, we set

$$F(\omega) = \{x \in E : x \in T(\omega, x)\},$$

and

$$F_n(\omega) = \{x \in E : d(x, T(\omega, x)) \leq \frac{1}{n} \text{diam}C\}.$$

It follows from Theorem 3.4 of Benavides-Ramírez's [4] that $F(\omega)$ is nonempty. Clearly $F(\omega) \subseteq F_n(\omega)$, and $F_n(\omega)$ is closed and convex. Furthermore, by Lemma 2.4, each F_n is measurable. Then, by Lemma 2.1, each F_n admits a measurable selector $x_n(\omega)$ and $d(x_n(\omega), T(\omega, x_n(\omega))) \leq \frac{1}{n} \text{diam}E \rightarrow 0$ as $n \rightarrow \infty$.

Let $\{n_\alpha\}$ be an ultranet of the positive integer $\{n\}$. Defined a function $f : \Omega \times E \rightarrow \mathbb{R}^+ := [0, \infty)$ by

$$f(\omega, x) = \limsup_\alpha \|x_{n_\alpha}(\omega) - x\|, \quad x \in E.$$

Since $\{x_{n_\alpha}(\omega)\}$ is countable, it is easily seen that $f(\omega, \cdot)$ is measurable and $f(\omega, \cdot)$ is continuous and convex, therefore it is a weakly lower semicontinuous function. Hence by Lemma 2.5, the marginal functions

$$r(\omega) := \inf_{x \in E} f(\omega, x)$$

and

$$A(\omega) := \{x \in E : f(\omega, x) = r(\omega)\}$$

are measurable, and $A(\omega)$ is a weakly compact convex subset of E . Note that $A(\omega) = A(E, \{x_{n_\alpha}(\omega)\})$, and $r(\omega) = r(E, \{x_{n_\alpha}(\omega)\})$. Moreover, we can apply Lemma 2.8 to obtain

$$r_E(A(\omega)) \leq \lambda r(E, \{x_{n_\alpha}(\omega)\}), \tag{3.1}$$

where $\lambda = 1 - \Delta_{X,\alpha}(1^-) < 1$, since $\varepsilon_\alpha(X) < 1$. By Lemma 2.1 we can take $x_0(\omega)$ as a measurable selector of $A(\omega)$. For each $\omega \in \Omega$ and $n \geq 1$, we define the contraction $T_n(\omega, \cdot) : A(\omega) \rightarrow KC(X)$ defined by

$$T_n(\omega, x) = \frac{1}{n}x_0 + \left(\frac{n-1}{n}\right)T(\omega, x),$$

for each $(\omega, x) \in \Omega \times E$. We shall prove that the inwardness of T on E implies a weaker inwardness of T on A , i.e.,

$$T(\omega, x) \cap I_{A(\omega)}(x) \neq \emptyset, \forall x \in A(\omega). \tag{3.2}$$

Indeed, the compactness of $T(\omega, x_{n_\alpha}(\omega))$ implies that for each n_α fixed $\omega \in \Omega$, we can take $y_{n_\alpha}(\omega) \in T(\omega, x_{n_\alpha}(\omega))$ such that

$$\|x_{n_\alpha}(\omega) - y_{n_\alpha}(\omega)\| = d(x_{n_\alpha}(\omega), T(\omega, x_{n_\alpha}(\omega))).$$

Since $T(\omega, x(\omega))$ is compact, for each $x(\omega) \in A(\omega)$, we can find $z_{n_\alpha}(\omega) \in T(\omega, x(\omega))$ such that

$$\begin{aligned} \|y_{n_\alpha}(\omega) - z_{n_\alpha}(\omega)\| &= d(y_{n_\alpha}(\omega), T(\omega, x(\omega))) \\ &\leq H(T(\omega, x_{n_\alpha}(\omega)), T(\omega, x(\omega))) \\ &\leq \|x_{n_\alpha}(\omega) - x(\omega)\|. \end{aligned}$$

Let $z(\omega) = \lim_{\alpha} z_{n_{\alpha}}(\omega) \in T(\omega, x(\omega))$. It should remain to prove $z(\omega) \in I_{A(\omega)}(x)$. It follows that

$$\begin{aligned} \limsup_{\alpha} \|x_{n_{\alpha}}(\omega) - z(\omega)\| &= \limsup_{\alpha} \|y_{n_{\alpha}}(\omega) - z_{n_{\alpha}}(\omega)\| \\ &\leq \limsup_{\alpha} \|x_{n_{\alpha}}(\omega) - x(\omega)\| \\ &= r(\omega). \end{aligned}$$

Since $z(\omega) \in T(\omega, x(\omega)) \subseteq I_E(x)$, for each fixed $\omega \in \Omega$ there exist $\lambda \geq 0$ and $v(\omega) \in E$ such that

$$z(\omega) = x(\omega) + \lambda(v(\omega) - x(\omega)).$$

If $\lambda \leq 1$, then by the convexity of E , $z(\omega) \in E$ and hence $z(\omega) \in A(\omega) \subseteq I_{A(\omega)}(x)$ and we are done. So assume that $\lambda > 1$. Then we can write

$$v(\omega) = \mu z(\omega) + (1 - \mu)x(\omega) \text{ with } \mu = \frac{1}{\lambda} \in (0, 1).$$

It follows that

$$\begin{aligned} f(\omega, v(\omega)) &= \limsup_{\alpha} \|x_{n_{\alpha}}(\omega) - v(\omega)\| \\ &\leq \mu \limsup_{\alpha} \|x_{n_{\alpha}}(\omega) - z(\omega)\| + (1 - \mu) \limsup_{\alpha} \|x_{n_{\alpha}}(\omega) - x(\omega)\| \\ &\leq r(\omega). \end{aligned}$$

Therefore $v(\omega) \in A(\omega)$ and thus $z(\omega) = x(\omega) + \lambda(v(\omega) - x(\omega))$ belong to $I_{A(\omega)}(x)$. That is $T(\omega, x(\omega)) \cap I_{A(\omega)}(x) \neq \emptyset \forall x(\omega) \in A(\omega)$.

Now, we have a mapping $T(\omega, \cdot) : A(\omega) \rightarrow KC(X)$ which satisfies the boundary condition (3.2). Consequently, since $T_n(\omega, \cdot)$ is $1-\chi$ -contractive mapping, it is easily seen that $T_n(\omega, \cdot)$ is χ -condensing (see [3]). By lemma 2.9, $T_n(\omega, \cdot)$ has a fixed point $x_n(\omega) \in A(\omega)$, *i.e.* $F(\omega) \cap A(\omega) \neq \emptyset$. Also, we have

$$dist(x_n(\omega), T(\omega, x_n(\omega))) \leq \frac{1}{n} diam E \rightarrow 0 \text{ as } n \rightarrow \infty.$$

Thus, $F_n^1(\omega) := \{x \in A(\omega) : d(x, T(\omega, x)) \leq \frac{1}{n} diam E\} \neq \emptyset$ for each $n \geq 1$ is closed and measurable. Hence, by Lemma 2.1, we can choose x_n^1 a measurable selector of F_n^1 , and from definition of it we have $x_n^1(\omega) \in A(\omega)$ and $d(x_n^1(\omega), T(\omega, x_n^1(\omega))) \rightarrow 0$ as $n \rightarrow \infty$. Consider the function $f_2 : \Omega \times E \rightarrow \mathbb{R}^+$ defined by

$$f_2(\omega, x) = \limsup_{\alpha} \|x_{n_{\alpha}}^1(\omega) - x\|, \quad \forall \omega \in \Omega.$$

As above, f_2 is a measurable function and weakly lower semicontinuous function. Then the marginal function

$$r_2(\omega) := \inf_{x \in A(\omega)} f_2(\omega, x)$$

and

$$A^1(\omega) := \{x \in A(\omega) : f_2(\omega, x) = r_2(\omega)\}$$

are measurable. Since $A^1(\omega) = A(A(\omega), \{x_{n_\alpha}^1(\omega)\})$, it follows that $A^1(\omega)$ is a weakly compact and convex. Also $r_2(\omega) = r(A(\omega), \{x_{n_\alpha}^1(\omega)\})$. We proceed as before to obtain that

$$T(\omega, x(\omega)) \cap I_{A^1}(x(\omega)) \neq \emptyset \quad \forall x(\omega) \in A^1 = A(A(\omega), \{x_{n_\alpha}^1(\omega)\}),$$

and by (3.1) we obtain that

$$r_E(A^1) \leq \lambda r(A(\omega), \{x_{n_\alpha}^1(\omega)\}) \leq \lambda r_E(A(\omega)). \quad (3.3)$$

By induction, for each $m \geq 1$, we take a sequence $\{x_n^m(\omega)\}_n \subseteq A^{m-1}$ such that $\lim_n d(x_n^m(\omega), T(\omega, x_n^m(\omega))) = 0$ for each fixed $\omega \in \Omega$. By means of the ultranet $\{x_{n_\alpha}^m(\omega)\}_\alpha$ we construct the set $A^m := A(E, \{x_{n_\alpha}^m(\omega)\})$ such that

$$r_E(A^m) \leq \lambda^m r_E(A(\omega)). \quad (3.4)$$

Choose $x_m(\omega)$ as a measurable selector of A^m . We shall prove that $\{x_m(\omega)\}_m$ is a Cauchy sequence. For each $m \geq 1$, we have

$$\begin{aligned} \|x_{m-1}(\omega) - x_m(\omega)\| &\leq \|x_{m-1}(\omega) - x_n^m(\omega)\| + \|x_n^m(\omega) - x_m(\omega)\| \\ &\leq \text{diam}A_{m-1}(\omega) + \|x_n^m(\omega) - x_m(\omega)\|. \end{aligned}$$

Since $\text{diam}A^m \leq 2r_E(A^m)$, taking upper limit as $n \rightarrow +\infty$, we have

$$\begin{aligned} \|x_{m-1}(\omega) - x_m(\omega)\| &\leq \text{diam}A^{m-1} + \limsup_n \|x_n^m(\omega) - x_m(\omega)\| \\ &= \text{diam}A^{m-1} + r(E, \{x_n^m(\omega)\}) \\ &\leq \text{diam}A^{m-1} + r_E(A^{m-1}) \\ &\leq 2r_E(A^{m-1}) + r_E(A^{m-1}) \\ &= 3\lambda^{m-1}r_E(A(\omega)). \end{aligned}$$

Since $\lambda < 1$, hence $\{x_m(\omega)\}_{m \geq 1}$ is a Cauchy sequence we conclude that there exists $x(\omega) \in E$ such that $x_m(\omega)$ converges to $x(\omega)$. Finally, we will show that $x(\omega)$ is a random

fixed point of T . Indeed, for each $m \geq 1$, we have

$$\begin{aligned} d(x_m(\omega), T(\omega, x_m(\omega))) &\leq \|x_m(\omega) - x_n^m(\omega)\| + d(x_n^m(\omega), T(\omega, x_n^m(\omega))) \\ &\quad + H(T(\omega, x_n^m(\omega)), T(\omega, x_m(\omega))) \\ &\leq 2\|x_m(\omega) - x_n^m(\omega)\| + d(x_n^m(\omega), T(\omega, x_n^m(\omega))). \end{aligned}$$

Taking the upper limit as $n \rightarrow +\infty$,

$$\begin{aligned} d(x_m(\omega), T(\omega, x_m(\omega))) &\leq 2 \limsup_n \|x_m(\omega) - x_n^m(\omega)\| \\ &\leq 2\lambda^{m+1}r_E(A(\omega)). \end{aligned}$$

Now taking the limit $m \rightarrow +\infty$ on both sides we get $d(x_m(\omega), T(\omega, x_m(\omega))) = 0$, and the continuity of $T(\omega, \cdot)$ implies that $d(x(\omega), T(\omega, x(\omega))) = 0$, that is, $x(\omega) \in T(\omega, x(\omega))$. This completes the proof. \square

Corollary 3.2 (Domínguez Benavides and Lorenzo Ramírez [4, Theorem 3.4]) *Let X be a Banach space such that $\varepsilon_\alpha(X) < 1$, and E be a nonempty closed bounded convex subset of X . If $T : E \rightarrow KC(X)$ is a nonexpansive and $1 - \chi$ -contractive nonexpansive mapping, such that $T(E)$ is a bounded set, and which satisfies $Tx \subset I_E(x) \ \forall x \in E$, then T has a fixed point.*

Proof. Define a random operator $S : \Omega \times E \rightarrow KC(X)$ by $S(\omega, x) = T(x)$ for all $\omega \in \Omega$ and for all $x \in E$. Thus $S(\omega, \cdot)$ is a nonexpansive random operator and $1 - \chi$ -contractive mapping such that $S(\omega, E)$ is bounded for all $\omega \in \Omega$. Hence, by Theorem 3.1, $S(C_i)$ has a random fixed point $x(\omega) \in S(\omega, x) = T(x)$ for all $\omega \in \Omega$. Thus is completes of the proof. \square

Next we prove the random version of the following celebrated deterministic result due to Xu ([20, Theorem 3.4]). The proof below is inspired by same ideas in the proof of [20].

Theorem 3.3 *Let E be a nonempty closed, bounded and convex separable subset of a uniformly convex Banach space X and $T : \Omega \times E \rightarrow KC(X)$ be a multivalued nonexpansive random operator such that for each $\omega \in \Omega$, $T(\omega, E)$ is a bounded set, which satisfies the inwardness condition, i.e., for each $\omega \in \Omega$, $T(\omega, x) \subset \bar{I}_E(x)$, $\forall x \in E$. Then T has a random fixed point.*

Proof. Fix $x_0 \in E$ for each $n \geq 1$, define the mapping $T_n : E \rightarrow KC(X)$ by

$$T_n(\omega, x) = \frac{1}{n}x_0 + (1 - \frac{1}{n})T(\omega, x), \quad \omega \in \Omega, x \in E.$$

Then T_n is a multivalued random contraction satisfying the same boundary condition as T does, i.e. we have, $T_n(\omega, x) \subset \bar{I}_E(x)$ for all $x \in E$. Hence, by [20, Theorem 1.4], $T_n(\omega, \cdot)$ has a random fixed point denoted $x_n(\omega)$. Also it is easily seen that we have $dist(x_n(\omega), T(\omega, x_n(\omega))) \leq \frac{1}{n}diamE \rightarrow 0$ as $n \rightarrow \infty$. Let $\{n_\alpha\}$ be a universal subnet of the positive integers $\{n\}$. Define a function $f : \Omega \times E \rightarrow \mathbb{R}^+$ by

$$f(\omega, x) = \limsup_n \|x_{n_\alpha}(\omega) - x\|, \quad \forall \omega \in \Omega.$$

Since $\{x_{n_\alpha}\}$ is countable, it is easily seen that for each $x \in E$, $f(\cdot, x) : \Omega \rightarrow \mathbb{R}^+$ is measurable and each $\omega \in \Omega$, $f(\omega, \cdot) : E \rightarrow \mathbb{R}^+$ is continuous and convex (and hence weakly lower semicontinuous (w-l.s.c.)). Since the space X is uniformly convex and E is weakly compact and convex for each $\omega \in \Omega$, there exists exactly a point $x(\omega) \in E$ such that

$$f(\omega, x(\omega)) = \inf_{x \in E} f(\omega, x) =: r(\omega).$$

Note that $x(\omega)$ is an asymptotic center of the net $\{x_{n_\alpha}(\omega)\}$ with respect to E . Lim [12], and Kirk and Massa [10] actually proved that for each $\omega \in \Omega$, $x(\omega)$ is a fixed point of the map $T(\omega, \cdot)$. By using the same argument as in the proof of Xu ([20, p.1091]), we obtain $x(\omega)$ is measurable. Therefore $x(\omega)$ is a random fixed point of T . The proof of the theorem, is complete. \square

Corollary 3.4 (Xu cf. [20]) *Assume X is a uniformly convex Banach space, E is a closed bounded convex subset of X , and $T : E \rightarrow KC(X)$ is a nonexpansive mapping satisfying the inwardness condition, i.e., $Tx \subset \bar{I}_E(x)$, $x \in E$. Then T has a fixed point.*

Corollary 3.5 (Xu cf. [18]) *Let (Ω, Σ) be a measurable spaces with Σ a sigma-algebra of subsets of Ω . Let E be a nonempty, bounded, closed, convex and separable subset of a uniformly convex Banach space X , and let $T : \Omega \times E \rightarrow KC(E)$ be a multivalued nonexpansive random operator. Then T has a random fixed point.*

Proof. It follows from Theorem 3.3, since every self multivalued mappings satisfies the inwardness condition. \square

Acknowledgement

The second author would like to thank the Thailand Research Fund for financial support. Moreover, the authors also wishes to thank to an anonymous referee for his suggestions which led to an improved presentation of the manuscript.

References

- [1] Ayerbe Toledano, J. M., Domínguez Benavides, T. and López Acedo G.: “Measures of noncompactness in metric fixed point theory”: *Advances and Applications Topics in metric fixed point theory*, Birkhuser-Verlag, Basel, **99** (1997).
- [2] Deimling, K.: “*Nonlinear Functional Analysis*”, Springer-Verlag, Berlin (1974).
- [3] Domínguez Benavides, T. and Lorenzo Ramírez, P.: “Fixed point theorem for Multivalued nonexpansive mapping without uniform convexity”, *Abstr. Appl. Anal.* 6, 375-386 (2003).
- [4] Domínguez Benavides, T. and Lorenzo Ramírez, P.: “Fixed point theorem for multivalued nonexpansive mapping satisfying inwardness conditions”, *J. Math. Anal. Appl.* 291, 100-108 (2004).
- [5] Domínguez Benavides, T., Lopez Acedo G. and Xu, H.-K.: “Random fixed point of set-valued operator”, *Proc. Amer. Math. Soc.* 124, 838-838 (1996).
- [6] Hans, O.: “Random fixed point theorems”, *Transac. of the 1st Prague Conference*, pp. 105-125 (1957).
- [7] Hans, O.: “Random operator equations”, *Proc. of the 4th Berkeley Symposium*, vol.2, pp. 185-201 (1960).
- [8] Itoh, S.: “Random fixed point theorem for a multivalued contraction mapping”, *Pacific J. Math.* 68, 85-90 (1977).
- [9] Kelly, J.L.: “*General Topology*”, van Nostrand, Princeton, NJ, 1955.
- [10] Kirk, W.A. and Massa, S.: “Remarks on asymptotic and Chebyshev centers”, *Houston J. Math.* 16, 357-364 (1990).
- [11] Kumam, P. and Plubtieng, S.: “The Characteristic of noncompact convexity and random fixed point theorem for set-valued operators”, (Accepted) *Czechoslovak Math. J.*
- [12] Lim, T.C.: “A fixed point theorem for multivalued nonexpansive mappings in a uniformly convex Banach space”, *Bull. Amer. Math. Soc.* 80, 1123-1126 (1974).

- [13] Lorenzo Ramírez, P.: “Some random fixed point theorems for nonlinear mappings”, *Nonlinear Anal.* 50, 265-274 (2002).
- [14] Lorenzo Ramírez, P.: “Random fixed point of uniformly Lipschitzian mappings”, *Nonlinear Anal.* 57, 23-34 (2004).
- [15] Shahzad, N. and Latif, S.: “Random fixed points for several classes of 1-ball-contractive and 1-set-contractive random maps”, *J. Math. Anal. Appl.* 237, 83-92 (1999).
- [16] Tan, K.-K. and Yuan, X.Z.: “Some Random fixed point theorem ”, in: K.-K. Tan (Ed), *Fixed Point Theory and Applications*, World Sciedtific, Singapro, 334-345 1992.
- [17] Wagner, D.-H.: “Survey of measurable selection theorems”, *SIAM J. Control Optim.* 15, 859-903 (1977).
- [18] Xu, H. K.: “A Random fixed point theorem for multivalued nonexpansive operators in Uniformly convex Banach spaces ”, *Proc. Amer. Math. Soc.* 117, 4, 1089-1092 (1993).
- [19] Xu, H. K.: “Random fixed point theorems for nonlinear Uniform Lipschitzian mappings ”, *Nonlinear Anal.* 26, 7, 1301-1311 (1996).
- [20] Xu, H. K.: “Multivalued nonexpansive mappings in Banach spaces. *Nonlinear Anal.* 43, 693-706 (2001).
- [21] Yuan, X. and Yu, J.: “Random fixed point theorems for nonself mappings ”, *Nonlinear Anal.* 26, 6, 1097-1102 (1996).

Poom KUMAM
 Department of Mathematics King Mongkut’s
 University of Technology
 Bangkok 10140 THAILAND
 e-mail: poom.kum@kmutt.ac.th
 Somyot PLUBTIENG
 Department of Mathematics,
 Naresuan University,
 Pitsanulok 65000. THAILAND
 e-mail: Somyotp@nu.ac.th.

Received 08.03.2005