

1-1-2006

## On Finitary Permutation Groups

ALİ OSMAN ASAR

Follow this and additional works at: <https://journals.tubitak.gov.tr/math>



Part of the [Mathematics Commons](#)

---

### Recommended Citation

ASAR, ALİ OSMAN (2006) "On Finitary Permutation Groups," *Turkish Journal of Mathematics*: Vol. 30: No. 1, Article 10. Available at: <https://journals.tubitak.gov.tr/math/vol30/iss1/10>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Mathematics by an authorized editor of TÜBİTAK Academic Journals. For more information, please contact [academic.publications@tubitak.gov.tr](mailto:academic.publications@tubitak.gov.tr).

## On Finitary Permutation Groups

*Ali Osman Asar*

### Abstract

In this work we give some sufficient conditions under which the structure of a transitive group of finitary permutations on an infinite set can be determined from the structure of a point stabilizer. Also, we give some sufficient conditions for the existence of a proper subgroup having an infinite orbit in a totally imprimitive  $p$ -group of finitary permutations. These results, with the help of some known results, give sufficient conditions for the nonexistence of a perfect locally finite minimal non  $FC$  - ( $p$ -group).

**Key Words:** Finitary permutation, primitive, almost primitive, totally imprimitive.

### 1. Introduction

Let  $G$  be a transitive subgroup of  $FSym(\Omega)$ , where  $\Omega$  is infinite. Many authors have investigated  $G$  by imposing suitable conditions on a point stabilizer (see, for example, [1], [2], [4], [7], [9]). Also another problem which might be interesting is finding sufficient conditions under which  $G$  can have a proper subgroup having an infinite orbit. In view of the important reduction theorems given in [3] and [8], any solution of the last problem means a solution for the following well known problem: Does there exist a perfect minimal non  $FC$  - ( $p$ -group)? The aim of this work is to obtain some sufficient conditions about the problems described above.

Let  $\Omega$  be a (possibly infinite) set and let  $Sym(\Omega)$  be the symmetric group on  $\Omega$ . For each  $x \in Sym(\Omega)$  the set  $supp(x) = \{i \in \Omega : x(i) \neq i\}$  is called the *support* of  $x$  and if  $supp(x)$  is finite, then  $x$  is called a *finitary permutation*. The set of all the finitary permutations on  $\Omega$  forms a subgroup which is denoted by  $FSym(\Omega)$ . Let  $G$  be a subgroup

---

2000 *AMS Mathematics Subject Classification:* 20B 07 20B 35 20E 25

of  $Sym(\Omega)$  and  $a \in \Omega$ . Then  $G_a = \{g \in G : g(a) = a\}$  is called the *stabilizer* of  $a$  in  $G$  and  $G(a) = \{g(a) : a \in G\}$  is called the *orbit* of  $G$  containing  $a$ . More generally if  $\Delta$  is a nonempty subset of  $\Omega$ , then  $G_\Delta = \{g \in G : g(i) = i \text{ for all } i \in \Delta\}$  and  $G_{\{\Delta\}} = \{g \in G : g(\Delta) = \Delta\}$  are called the *pointwise* and the *setwise* stabilizers of  $\Delta$ . Furthermore if  $g(\Delta) = \Delta$  or  $g(\Delta) \cap \Delta = \emptyset$ , for every  $g \in G$ , then  $\Delta$  is called a *block* for  $G$ . A block  $\Delta$  which is not equal to  $G$  and contains at least two elements is called *non-trivial*.

Finally a group  $G$  is called a *minimal non FC - group* if  $G$  is not an *FC - group* but every proper subgroup of  $G$  is an *FC - group*.

The main results of this work are stated below.

**Theorem 1.1** *Let  $G$  be a transitive subgroup of  $FSym(\Omega)$ , where  $\Omega$  is infinite. Let  $F$  be a finite non-abelian subgroup of  $G$  and let  $\Delta$  be a non-trivial block such that  $supp(F) \subseteq \Delta$ . Let  $S$  be a normal solvable subgroup of  $G_{\{\Delta\}}$  of derived length  $d \geq 0$ . Then  $\langle S^x : x \in G \rangle \neq G$ .*

**Theorem 1.2** *Let  $G$  be a transitive subgroup of  $FSym(\Omega)$ , where  $\Omega$  is infinite. Then the following hold:*

- (a) *A point stabilizer cannot be solvable.*
- (b) *If  $G$  satisfies the normalizer condition, then it is a  $p$ -group and  $G'$  is a minimal non FC - group.*

If in Theorem 1.2(a)  $G$  is barely transitive (see[4] for the definition of a barely transitive group), then the result follows from [1, Theorem] or [4, Theorem 1]. Furthermore Theorem 1.2(b) is not true if the normalizer condition is satisfied only by a point stabilizer. Indeed in the example given at the end of Section 2 it is shown that in Wiegold's group[16, p.468] a point stabilizer satisfies the normalizer condition but the commutator subgroup of it, which is a perfect proper subgroup of the group, is not a minimal non *FC - group*.

**Theorem 1.3** *Let  $G$  be a transitive subgroup of  $FSym(\Omega)$ , where  $\Omega$  is infinite. Then the following hold:*

- (a) *If a point stabilizer is locally solvable, then  $G$  is locally solvable.*
- (b) *If a point stabilizer is locally nilpotent - by - solvable, then  $G$  is a  $p$ -group for some prime  $p$ .*

Pinnock[12] shows that if a transitive subgroup of  $FSym(\Omega)$  is either locally (nilpotent - by - abelian) or locally supersolvable, then it is a  $p$  - group. These results can be generalized as follows:

**Corollary 1.4** *Let  $G$  be a transitive subgroup of  $FSym(\Omega)$ , where  $\Omega$  is infinite. Then the following hold:*

- (a) *If a point stabilizer is locally (nilpotent - by - abelian), then  $G$  is a  $p$ -group for some prime  $p$ .*
- (b) *If a point stabilizer is locally supersolvable then  $G$  is a  $p$ -group for some prime  $p$ .*

**Theorem 1.5** *Let  $G$  be a totally imprimitive  $p$  - subgroup of  $FSym(\Omega)$ , where  $\Omega$  is infinite. Suppose that for every non-normal finite subgroup  $F$  of  $G$  there exists  $y \in G \setminus N_G(F)$  such that  $y^p \in C_G(F)$ . Then  $G$  contains a proper subgroup that has an infinite orbit.*

**Corollary 1.6** *Let  $G$  be a totally imprimitive  $p$  - subgroup of  $FSym(\Omega)$ , where  $\Omega$  is infinite. Suppose that for every non-normal finite subgroup  $F$  of  $G$  there exists  $y \in G \setminus N_G(F)$  such that  $y^p \in FC_G(F)$ . Then  $G$  contains a proper subgroup that has an infinite orbit.*

**Corollary 1.7** *Let  $G$  be a totally imprimitive  $p$  - subgroup of  $FSym(\Omega)$ , where  $\Omega$  is infinite. Suppose that there exists an infinite properly ascending chain of non-trivial blocks  $\Delta_1 \subset \Delta_2 \subset \dots \subset \Delta_k \subset \dots$  for  $G$  such that the following holds: For each  $k \geq 1$   $\langle F_k^x : x \in G \rangle$  is the largest normal subgroup of  $G$  that is contained in  $G_{\{\Delta_k\}}$ , where  $F_k = \{x \in G : \text{supp}(x) \subseteq \Delta_k\}$ . Then  $G$  contains a proper subgroup that has an infinite orbit.*

**Corollary 1.8** *Let  $G$  be a totally imprimitive  $p$  - subgroup of  $FSym(\Omega)$ , where  $\Omega$  is infinite. Suppose that there exists an infinite properly ascending chain of non-trivial blocks  $\Delta_1 \subset \Delta_2 \subset \dots \subset \Delta_k \subset \dots$  for  $G$  such that the following holds: For each  $k \geq 1$  there exists  $y_k \in G \setminus G_{\Delta_k}$  such that  $\langle y_k \rangle \cap G_{\{\Delta_k\}} \leq G_{\Delta_k}$ . Then  $G$  contains a proper subgroup that has an infinite orbit.*

Corollary 1.8 can be used to prove the following:

**Corollary 1.9** ([6, Theorem 3]) *Let  $G$  be a totally imprimitive  $p$ -subgroup of  $FSym(\Omega)$ , where  $\Omega$  is infinite. Suppose that  $G = \langle x \in G : x^p = 1 \rangle$ . Then  $G$  cannot be a minimal non  $FC$ -group.*

**Proof.** Assume that  $G$  is a minimal non  $FC$ -group. Then every orbit of every proper subgroup of  $G$  is finite by [5, Lemma 8.3D] or [16, Theorem 1]. Let  $X = \{x \in G : x^p = 1\}$ . Assume that  $G = \langle X \rangle$ . Let  $\Delta_1 \subset \Delta_2 \subset \dots \subset \Delta_k \subset \dots$  be an infinite properly ascending chain of non-trivial blocks for  $G$ . By hypothesis for each  $k \geq 1$  there exists an  $x \in X$  such that  $x \in G \setminus G_{\{\Delta_k\}}$  and  $\langle x \rangle \cap G_{\{\Delta_k\}} = 1$ . But then  $G$  contains a proper subgroup that has an infinite orbit by Corollary 1.8, which is a contradiction.  $\square$

**Remark 1.10** An easy induction shows that the group  $G$  of Corollary 1.9 cannot be generated by a subset of finite exponent.

Theorem 1.5 together with the important reduction theorems given in [3, Theorem 1] or [8, Theorem] gives the following:

**Theorem 1.11** *Let  $G$  be a locally finite  $p$ -group that is also a minimal non  $FC$ -group. Assume that for every finite non-normal subgroup  $F$  of  $G$  there exists  $y \in G \setminus N_G(F)$  such that  $y^p \in FC_G(F)$ . Then  $G$  cannot be perfect.*

The notation and the definitions are standard and may be found in [5], [10], [11] and [13]. Finally for a nonempty subset  $X$  we define  $exp(X)$  to be the maximum of the set  $\{o(x) : x \in X\}$ , if it exists, otherwise we set it equal to  $\infty$ .

## 2. Proofs of Theorems 1, 2, 3

Let  $G$  be a transitive subgroup of  $FSym(\Omega)$ , where  $\Omega$  is infinite. If  $G$  has no non-trivial blocks, then it is called **primitive**, and if  $G$  has non-trivial blocks, then it is called **imprimitive**. If  $G$  is primitive, then it is isomorphic to  $Alt(\Omega)$  or  $FSym(\Omega)$  by [5, Lemma 8.3A] or [10, Theorem 2.3]. Next suppose that  $G$  is imprimitive. Then any non-trivial block is finite. If  $G$  has a maximal non-trivial block  $\Delta$ , then  $\Sigma = \{x(\Delta) : x \in G\}$  has a **system of blocks** for  $G$ . It is easy to see that  $G$  acts primitively on  $\Sigma$ , and so it has an epimorphic image which is isomorphic to  $Alt(\Omega)$  or  $FSym(\Omega)$  by [5, Lemma 8.3A] or [10, Theorem 2.3] or [14, Proposition]. In the remaining case there exists a strictly increasing infinite ascending chain

$$\Delta_1 \subset \Delta_2 \subset \cdots \subset \Delta_k \subset \dots (1)$$

of finite blocks for  $G$  such that  $\Omega = \bigcup_{k=1}^{\infty} \Delta_k$ . In this case, both of  $\Omega$  and  $G$  are countably infinite. Let  $k \geq 1$  and put  $\Sigma_k = \{x(\Delta_k) : x \in G\}$ . For each  $g \in G$  the equality  $\bar{g}(x(\Delta_k)) = (gx)(\Delta_k)$  defines a permutation  $\bar{g}$  on  $\Sigma_k$  and the correspondence  $g \rightarrow \bar{g}$  defines a representation of  $G$  into  $FSym(\Sigma_k)$ . Let  $N_k$  denote the kernel of this representation. Then

$$N_1 \leq N_2 \leq \cdots \leq N_k \leq \dots (2)$$

is an ascending chain of proper normal subgroups of  $G$  such that  $G = \bigcup_{k=1}^{\infty} N_k$  and each  $N_k$  is isomorphic to a restricted direct product of isomorphic copies of a finite epimorphic image of itself. In particular, each  $N_k$  is an  $FC$  - group (see [5, Lemma 8.3B(i)] or [11, Theorem 2.4]). The two cases of the imprimitive case are called *almost primitive* and *totally imprimitive* by P.M. Neumann. In the rest of this work these will be used without further explanation.

Most of the basic properties of infinite finitary permutation groups can be found in [5], [10], [11] and [6]. Some of them are collected in Lemmas 1, 2, 3 for the convenience of the reader.

**Lemma 2.1** *Let  $G$  be a transitive subgroup of  $FSym(\Omega)$  for some set  $\Omega$ . Let  $\Delta$  be a non-trivial block for  $G$  and let  $H$  be a non-trivial subgroup of  $G$  such that  $\text{supp}(H) \subseteq \Delta$ . Then the following hold:*

- (a) *Let  $\Gamma = \text{supp}(H)$  then  $G_\Gamma \leq C_G(H)$ ;*
- (b)  *$N_G(H) \leq G_{\{\Delta\}}$ ;*
- (c)  *$H \cap G_\Delta = 1$  and  $G_\Delta \leq C_G(H)$ ;*
- (d)  *$H^x \leq G_\Delta$  and so  $[H^x, H] = 1$  for all  $x \in G \setminus G_{\{\Delta\}}$ ;*
- (e) *If  $x \in G \setminus G_{\{\Delta\}}$ , then  $H' \leq [H, x]$*

**Proof.** (a) Let  $h \in H$  and  $x \in G_\Gamma$ . If  $i \in \Gamma$  then  $h(i) \in \Gamma$  and hence  $x(h(i)) = h(x(i))$  since  $x(i) = i$ . If  $i \notin \Gamma$  then  $x(i) \notin \Gamma$  and hence  $h(x(i)) = x(i) = x(h(i))$ . (b) and (c) are left to the reader. (d) Let  $h \in H$  and  $x \in G \setminus G_{\{\Delta\}}$ . Then  $\Delta \cap x(\Delta) = \Delta \cap x^{-1}(\Delta) = \emptyset$ .

Let  $i \in \Delta$ , then  $x^{-1}hx(i) = x^{-1}h(x(i)) = x^{-1}x(i) = i$  and so  $x^{-1}hx \in G_\Delta$ . Hence it follows that  $H^x \leq G_\Delta$  and so  $[H, H^x] = 1$  by (c). (e) See the proof of [5, Lemma 8.3C(i)].  $\square$

**Lemma 2.2** *Let  $G$  be a totally imprimitive subgroup of  $FSym(\Omega)$  and  $a \in \Omega$ , where  $\Omega$  is infinite. Then the following hold:*

- (a) *Every orbit of  $G_a$  is finite;*
- (b) *If  $K \leq G$  and  $K(a)$  is finite then  $[K : K \cap G_a]$  is finite.*

**Proof.** (a) Put  $H = G_a$  and choose  $b \in \Omega$ . By (1) there exists a finite block  $\Delta$  such that  $a, b \in \Delta$ . Then  $H \leq G_{\{\Delta\}}$ . Hence  $H(b) \subseteq H(\Delta) = \Delta$  and so  $H(b)$  is finite. (b) This follows from the fact that  $[K : K_a] = |K(a)|$  and  $K \cap G_a = K_a$ .  $\square$

**Lemma 2.3** *Let  $G$  be a subgroup of  $FSym(\Omega)$ , where  $F$  is a finite subgroup of  $G$  and  $\Delta$  is a block for  $G$  such that  $\text{supp}(F) \subseteq \Delta$ . Then the following hold:*

- (a)  $U = \{u \in G_{\{\Delta\}} : \text{supp}(u) \subseteq \Delta\}$  is a normal subgroup of  $G_{\{\Delta\}}$  with  $F \subseteq U$ .
- (b) Let  $y \in G$  and let  $t$  be the smallest positive integer such that  $y^t \in G_{\{\Delta\}}$ . Assume that  $y^t$  normalizes  $F$ . Then  $F^{\langle y \rangle} = F \times F^y \times \dots \times F^{y^{t-1}}$ .

**Proof.** (a) Clearly  $F \subseteq U$ . For any  $u, v \in U$  and  $x \in G_{\{\Delta\}}$  it is easy to check that  $uv^{-1}, u^x \in U$ .

(b) Since  $y^t$  normalizes  $F$  it follows that  $F^{\langle y \rangle} = \langle F^{y^k} : 0 \leq k \leq t-1 \rangle$ . Since  $y, y^2, \dots, y^{t-1}$  are not contained in  $G_{\{\Delta\}}$ ,  $F^{y^k} \leq G_\Delta$  and  $[F, F^{y^k}] = 1$  for all  $1 \leq k \leq t-1$  and so  $F \cap \langle F^{y^k} : 1 \leq k \leq t-1 \rangle = 1$  by Lemma 2.1. Also the above properties hold if  $F$  is replaced by  $F^{y^k}$  for any  $1 \leq k \leq t-1$ . Therefore  $F^{\langle y \rangle}$  is the direct product of  $F, F^y, \dots, F^{y^{t-1}}$ .  $\square$

**Lemma 2.4** *Let  $G$  be a transitive subgroup of  $FSym(\Omega)$ , where  $\Omega$  is infinite. Let  $F$  be a finite non-abelian subgroup of  $G$  and let  $\Delta$  be a non-trivial block for  $G$  such that  $\text{supp}(F) \subseteq \Delta$ . If  $A$  is a normal abelian subgroup of  $G_{\{\Delta\}}$ , then  $\langle A^x : x \in G \rangle \leq G_{\{\Delta\}}$ .*

**Proof.** Put  $H = G_{\{\Delta\}}$ . Let  $x \in G \setminus H$ . Then  $F \leq H^x$  by Lemma 2.1(d) and  $A^x \triangleleft H^x$ . If there is an  $a \in A^x \setminus H$  then  $F' \leq [F, a]$  and then  $A^x \leq C_G(F') \leq H$  by Lemma 2.1, which is a contradiction. Hence it follows that  $A^x \leq H$  for any  $x \in G$  and so the assertion follows.  $\square$

**Lemma 2.5** *Let  $G$  be a transitive subgroup of  $FSym(\Omega)$ , where  $\Omega$  is infinite and  $K$  be an ascendant subgroup of  $G$ . If  $K$  has an infinite orbit on  $\Omega$ , then  $K$  is a transitive normal subgroup of  $G$ .*

**Proof.** Assume that  $K(i)$  is infinite for some  $i \in \Omega$  but  $K$  is not normal in  $G$ . Then there exist ascendant subgroups  $K_1$  and  $K_2$  of  $G$  such that  $K < K_1 < K_2$ ,  $K \triangleleft K_1 \triangleleft K_2$  but  $K$  is not normal in  $K_2$ . Since  $K \triangleleft K_1$   $K(i)$  is a block for the action of  $K_1$  on  $K_1(i)$ . So if  $K(i) \neq K_1(i)$ , then there exists  $x \in K_1$  such that  $x(K(i)) \cap K(i) = \emptyset$  and so  $K(i) \subseteq \text{supp}(x)$  which is impossible since  $\text{supp}(x)$  is finite. Hence it follows that  $K(i) = K_1(i)$  and so  $K$  acts transitively on  $K_1(i)$ . Similarly  $K_1$  acts transitively on  $K_2$  and so  $K_1(i) = K_2(i)$  which yields that  $K(i) = K_2(i)$ . Continuing in this way it follows that  $K(i) = \Omega$  and so  $K$  is transitive on  $\Omega$ . Now since  $K_1(i)$  and  $K_2(i)$  are transitive subgroups of  $FSym(\Omega)$  having the property that  $K \triangleleft K_1 \triangleleft K_2$  it follows from [11, Theorem 3.3] that  $K \triangleleft K_2$  which is a contradiction.  $\square$

**Proof of Theorem 1.1** Let  $G$  be a transitive subgroup of  $FSym(\Omega)$ , where  $\Omega$  is infinite. Let  $F$  be a non-abelian subgroup of  $G$  and let  $\Delta$  be a non-trivial block such that  $\text{supp}(F) \subseteq \Delta$ . Then  $G$  cannot be primitive. First suppose that  $G$  has a maximal block  $\Gamma$  with  $\Delta \subseteq \Gamma$ . Put  $\Sigma = \{x(\Gamma) : x \in G\}$ . Let  $K$  be the kernel of the representation of  $G$  into  $FSym(\Sigma)$ . Then  $G/K$  is isomorphic to  $Alt(\Sigma)$  or  $FSym(\Sigma)$  by [5, Lemma 8.3(B)] or [14, Proposition]. Then also  $(G/K)_\Gamma$  is isomorphic to  $Alt(\Sigma \setminus \Gamma)$  or  $FSym(\Sigma \setminus \Gamma)$ . Moreover it is easy to see that  $(G/K)_\Gamma = G_{\{\Gamma\}}/K$ . Therefore  $G_{\{\Gamma\}}/K$  contains a unique subgroup of index  $\leq 2$  which is isomorphic to the simple group  $Alt(\Sigma \setminus \Gamma)$ . Since  $G_{\{\Delta\}}$  has finite index in  $G_{\{\Gamma\}}$  it follows that  $Alt(\Sigma \setminus \Gamma)$  is isomorphic to a subgroup of  $G_{\{\Delta\}}K/K$ . Hence it follows that any solvable normal subgroup of  $G_{\{\Delta\}}$  is contained in  $K$ .

Next suppose that  $G$  is totally imprimitive. Put  $H = G_{\{\Delta\}}$ . First suppose that  $G = G'$ . We use induction on  $d$ . By Lemma 2.4 the assertion is true for  $d \leq 1$ . So suppose that  $d > 1$  and the assertion is true for smaller derived lengths. Put  $L = \langle (S^{(d-1)})^x : x \in G \rangle$ . Again  $L \leq H$  by Lemma 2.4. Let  $M$  be the kernel of the representation of  $G$  into  $FSym(\Lambda)$ , where  $\Lambda = \{x(\Delta) : x \in G\}$  and put  $\bar{G} = G/M$ .



Then  $L \leq M$ . Clearly  $\bar{G}$  is totally imprimitive and the derived length of  $\bar{S}$  is less than  $d$ . Let  $\bar{F}_1$  be a finite non-abelian subgroup of  $\bar{G}$  and  $\Gamma$  be a non-trivial block for  $\bar{G}$  such that  $\Delta \subseteq \Gamma$  and  $\text{supp}(\bar{F}_1) \subseteq \Gamma$ . There exist  $1 = x_1, x_2, \dots, x_r \in G$  such that  $\Gamma = \{x_1(\Delta), x_2(\Delta), \dots, x_r(\Delta)\}$ . Put  $\Delta_1 = x_1(\Delta) \cup x_2(\Delta) \cup \dots \cup x_r(\Delta)$ . Then  $\Delta_1$  is a block for  $G$  and  $\bar{G}_{\{\Gamma\}} = \overline{G_{\{\Delta_1\}}}$ . Put  $\bar{T} = \bar{G}_\Gamma$ . Then  $\bar{G}_{\{\Gamma\}}/\bar{T}$  is a finite group and  $\bar{T} \leq \bar{G}_{x_i(\Delta)}$  for all  $i \geq 1$ . Thus  $\bar{G}_{\{\Gamma\}} = \bar{X}\bar{T}$  for some finite subgroup  $\bar{X}$  of  $\bar{G}_{\{\Gamma\}}$ . Since  $\bar{G}_{\{\Gamma\}}$  is an *FC*-group we may suppose that  $\bar{X}$  is normal in  $\bar{G}_{\{\Gamma\}}$ .

Put  $D_S = S \cap T$ . Since  $T \leq G_{\{\Delta\}}$ ,  $D_S \triangleleft T$  and  $S/D_S$  is finite. Now  $\overline{D_S} \cap C_{\bar{T}}(\bar{X})$  is normalized by  $\bar{X}\bar{T} = \bar{G}_{\{\Gamma\}}$  and  $[\overline{D_S} : \overline{D_S} \cap C_{\bar{T}}(\bar{X})]$  is finite since  $C_{\bar{T}}(\bar{X}) \cap \bar{T}$  has finite index in  $\bar{G}_{\{\Gamma\}}$ . Since the derived length of  $\overline{D_S} \cap C_{\bar{T}}(\bar{X})$  is less than  $d$ , it follows by induction hypothesis that  $\bar{R} = \langle (\overline{D_S} \cap C_{\bar{T}}(\bar{X}))^x : x \in G \rangle \neq \bar{G}$ . Clearly every orbit of  $R$  on  $\Omega$  is finite by Lemma 2.5 and [11, Theorem 1] since  $G = G'$ . Since  $\bar{T}$  and  $C_{\bar{T}}(\bar{X})$  have finite index in  $\bar{G}_{\{\Gamma\}}$ ,  $[\bar{G}_{\{\Gamma\}} : C_{\bar{T}}(\bar{X}) \cap \bar{T}] = m$  for some  $m \geq 1$ . Thus  $[\bar{S} : \bar{S} \cap (C_{\bar{T}}(\bar{X}) \cap \bar{T})] = [\bar{S} : \overline{D_S} \cap C_{\bar{T}}(\bar{X})] \leq m$  and hence  $|\bar{S}\bar{R}/\bar{R}| \leq m$ . Since  $G$  is totally imprimitive it is the union of an ascending chain of proper normal subgroups. Therefore there exists a proper normal subgroup  $N$  of  $G$  such that  $\bar{R} \leq \bar{N}$  and  $\bar{S}\bar{R}/\bar{R} \leq \bar{N}$ , since every orbit of  $R$  is finite. Clearly then  $\langle S^x : x \in G \rangle \neq G$ .

Now suppose that  $G' < G$ . If  $G'S \neq G$  then we are done. So suppose that  $G'S = G$ . Let  $W = G'$  and  $S_1 = S \cap W$ . Then  $U = \langle S_1^x : x \in G' \rangle \neq W$  by the above paragraph. Since every orbit of  $U$  is finite there exists a non-trivial block  $\Pi$  for  $G$  containing  $\Delta$  such that  $U \leq G_{\{\Pi\}}$ . Let  $B$  be the kernel of the representation of  $G$  into  $\text{FSym}(Y)$ , where  $Y = \{x(\Pi) : x \in G\}$ . Put  $\bar{G} = G/B$ . Since  $\bar{H} = \bar{S}(\bar{H}) \cap \bar{W}$  and  $\bar{S} \cap \bar{W} = 1$ , it follows that  $\bar{S} \leq Z(\bar{H})$ . Put  $V = G_{\{\Pi\}}$ . Then  $[V : H]$  is finite. Let  $A$  be the largest normal subgroup of  $V$  contained in  $H$ . Then  $V/A$  is finite. Put  $\bar{Z} = Z(\bar{A})$ . Then  $[\bar{S} : \bar{S} \cap \bar{Z}]$  is finite. Since  $\bar{Z} \triangleleft \bar{V}$ , if  $X = \langle Z^x : x \in G \rangle$  then  $\bar{X} \neq \bar{G}$  by Lemma 2.4. Now consider  $\bar{G}/\bar{X}$ . Then  $\bar{S}\bar{X}/\bar{X}$  is finite. Also  $\bar{G}/\bar{X}$  is an ascending union of proper normal subgroups. Therefore  $\bar{S}\bar{X}/\bar{X}$  is contained in a proper normal subgroup of  $\bar{G}/\bar{X}$ , which implies that  $S$  is contained in a proper normal subgroup of  $G$ . This completes the proof of the theorem.  $\square$

As an easy consequence of the above proof one can verify the following easily: Suppose that in Theorem 1.1  $G$  is perfect. Let  $W_d = \{S \leq G_{\{\Delta\}} : S \triangleleft G_{\{\Delta\}} \text{ and } S^{(d)} = 1\}$ . Then  $\langle S^x : S \in W_d \text{ and } x \in G \rangle \neq G$ .

**Proof of Theorem 1.2** Let  $G$  be a transitive subgroup of  $FSym(\Omega)$ , where  $\Omega$  is infinite.

(a) Let  $a \in \Omega$  and suppose that  $G_a$  is solvable. By [5, Lemma 8.3B(i)] or [10, Theorem 2.3]  $G$  cannot be primitive. First suppose that  $G$  is almost primitive. Then  $G$  has a maximal non-trivial block  $\Gamma$  containing  $a$ . Then, as in the proof of Theorem 1.1,  $G$  contains a normal subgroup  $K$  such that  $G/K$  is isomorphic to  $Alt(\Sigma)$  or  $FSym(\Sigma)$ , where  $\Sigma = \{x(\Gamma) : x \in G\}$  and then  $(G/K)_\Gamma$  is isomorphic to  $Alt(\Sigma \setminus \{\Gamma\})$  or  $FSym(\Sigma \setminus \{\Gamma\})$ . But since  $G_a K/K$  has finite index in  $(G/K)_\Gamma$ , this gives a contradiction.

Next suppose that  $G$  is totally imprimitive. Then  $G$  contains a non-abelian finite subgroup  $F$  and a non-trivial block  $\Delta$  such that  $supp(F) \cup \{a\} \subseteq \Delta$ . Clearly  $G_a$  has finite index in  $G_{\{\Delta\}}$  and so  $G_{\{\Delta\}}$  contains a normal solvable subgroup  $S$  of finite index. By Theorem 1.1  $M = \langle S^x : x \in G \rangle \neq G$ . Since  $G_{\{\Delta\}}/M$  is finite  $G_{\{\Delta\}}$  is contained in a proper normal subgroup  $N$  of  $G$ . In particular  $G_a \leq N$ . Since  $G_b$  is conjugate to  $G_a$  for any  $b \in \Omega$  it follows that  $G = \langle G_b : b \in \Omega \rangle \leq N$ , which is a contradiction.  $\square$

(b) Suppose that  $G$  satisfies the normalizer condition. Then  $G$  is locally nilpotent by [13, 12.2.2] and so it is a  $p$ -group for some prime  $p$  by [15, Theorem 1]. It is easy to see that  $G'$  is transitive and hence perfect by [11, Theorem 1]. Let  $H$  be a proper subgroup of  $G'$ . If every orbit of  $H$  is finite, then  $H$  is an  $FC$ -group by [5, Lemma 8.3(D)] or [16, Theorem 1]. So suppose that  $H(a)$  is infinite for some  $a \in \Omega$ . Then  $H$  is normal in  $G$  by Lemma 2.5, since it is ascendant in  $G$  by assumption. This implies that  $H(a)$  is a block for  $G$  and so  $H(a) = \Omega$  since any non-trivial block for  $G$  is finite. In particular then  $H$  is transitive on  $\Omega$ . But now  $G' \leq H$  by [5, Lemma 8.3C] or [11, Theorem 1], which is a contradiction. This completes the proof of the theorem.

**Proof of Theorem 1.3** Let  $G$  be a transitive subgroup of  $FSym(\Omega)$ , where  $\Omega$  is infinite.

Let  $a \in \Omega$  and suppose that  $G_a$  satisfies (a) or (b) of the theorem. Then it follows as in the proof of Theorem 1.2 that  $G$  can be neither primitive nor almost primitive. Therefore we may assume that  $G$  is totally imprimitive. Then  $G = \bigcup_{k=1}^{\infty} N_k$  by (2). For each  $k \geq 1$   $[N_k : N_k \cap G_a]$  is finite by Lemma 2.2(b). Let  $M_k$  be the largest normal subgroup of  $N_k$  contained in  $N_k \cap G_a$ . Then  $N_k/M_k$  is finite.

(a) Suppose that  $G_a$  is locally solvable. Then  $M_k$  is locally solvable. In fact then  $M_k$  is solvable since  $N_k$  is isomorphic to a subgroup of the direct product of isomorphic copies of a finite group as was explained in the introduction. Let  $S_k$  be the product of all the normal solvable subgroups of  $N_k$  for all  $k \geq 1$ . Then  $S_k$  is normal in  $G$  and  $N_k S_k/S_k$  is finite since  $M_k \leq S_k$ . Define  $S = \langle S_k : k \geq 1 \rangle$ . Then  $S$  is locally solvable. Also  $S \triangleleft G$  and  $G/S$  is an  $FC$ -group since  $G/S = \bigcup_{k=1}^{\infty} N_k S/S$  and each  $N_k S/S$  is finite. In

this case  $G/S$  cannot be represented as a group of finitary permutations on an infinite set by [5, Lemma 8.3D] or [16, Theorem 1] and so  $S$  must be transitive by [11, Lemma 2.1]. This implies that  $G' \leq S$  and so  $G$  is locally solvable, which completes the proof of (a).

(b) Suppose that  $G_a$  is locally nilpotent-by-solvable of derived length  $t \geq 0$ , say. Then  $G$  is locally solvable by (a). Let  $\eta(G_a)$  be the Hirsh - Plotkin radical of  $G_a$ . Then  $G_a/\eta(G_a)$  is solvable of derived length  $\leq t$ . Clearly if  $G'$  is locally nilpotent then the assertion follows from [15, Theorem 1] and [12, Lemma 2.1], since  $G'$  is transitive on  $\Omega$ . Therefore it suffices to show that  $G'$  is locally nilpotent. Without loss of generality we may suppose that  $G = G'$ .

Since  $G_a$  is locally nilpotent-by-solvable of derived length  $t$  and  $M_k \leq G_a$  it follows that  $M_k/\eta(M_k)$  is solvable of derived length  $\leq t$ . Also  $\eta(M_k) \leq \eta(N_k)$  for all  $k \geq 1$ . Define  $K = \langle \eta(N_k) : k \geq 1 \rangle$ . Then  $K$  is a locally nilpotent normal subgroup of  $G$ . So if  $K$  is transitive on  $\Omega$ , then  $G = G' \leq K$  and thus  $G$  is locally nilpotent. Assume if possible that  $K$  is not transitive. Then every orbit of  $K$ , being a block for  $G$ , must be finite.

Let  $a \in \Omega$ ,  $\Delta = K(a)$  and  $\Sigma = \{x(\Delta) : x \in G\}$ . Let  $L$  be the kernel of the representation of  $G$  into  $FSym(\Sigma)$ . Then  $K \leq L$ . Put  $\bar{G} = G/L$ . Then  $\bar{G}$  is transitive on  $\Sigma$ , in fact it is totally imprimitive since  $G$  is locally solvable. Now  $\bar{G} = \bigcup_{k=1}^{\infty} \bar{N}_k$ , and for each  $k \geq 1$ ,  $\bar{M}_k$  is a solvable normal subgroup of finite index of  $\bar{N}_k$  with derived length  $\leq t$ . Therefore each  $\bar{N}_k$  contains a characteristic subgroup of finite index with derived length  $\leq t^2$  by [4, Lemma 4]. But since  $\bar{G}$  is perfect, it follows that each  $\bar{N}_k$  is solvable of derived length  $\leq t^2 + 1$  for all  $k \geq 1$  and thus  $\bar{G}$  is solvable, which is a contradiction.  $\square$

**Proof of Corollary 1.4** (a) Let  $a \in \Omega$  and put  $H = G_a$ . Suppose that  $H$  is locally (nilpotent-by-abelian). Then  $G$  is locally solvable by Theorem 1.3(a), which implies that  $G$  is totally imprimitive and hence countably infinite. Let  $F_1 \leq F_2 \leq \dots \leq F_i \leq \dots$  be an ascending chain of finite subgroups of  $H$  whose union is equal to  $H$ . By hypothesis  $F_i'$  is nilpotent for each  $i \geq 1$ . Since

$$H' = \left( \bigcup_{i=1}^{\infty} F_i \right)' = \bigcup_{i=1}^{\infty} F_i',$$

and  $(F_i)' \leq (F_{i+1})'$  for all  $i \geq 1$ , it follows that  $H'$  is locally nilpotent and so  $H$  is locally nilpotent-by-abelian. Therefore by Theorem 1.3(b)  $G$  is a  $p$ -group for some prime  $p$ .

(b) Suppose that  $G_a$  is locally supersolvable. Since a supersolvable group is nilpotent-by-abelian by [13, 5.4.10],  $G_a$  is locally (nilpotent-by-abelian) and so  $G$  is a  $p$ -group for some prime  $p$  by (a), which was to be shown.  $\square$

**Example** Let  $G$  be the 2 - group constructed by Wiegold in [16, p. 468]. Then  $G$  is a totally imprimitive subgroup of  $FSym(\mathbb{N})$ . By Lemma 2.2(a) every orbit of a point stabilizer is finite and so it is an  $FC$ -group. We show that  $G'$  is not a minimal non  $FC$ -group, which will show that the condition of Theorem 1.2(b) cannot be restricted to a point stabilizer. We will adopt the description of Wiegold's group given in [5, Exercise 8.3.1]. Thus for each  $k \geq 1$  let  $T_k$  be the subset of  $FSym(\mathbb{N})$  defined by

$$T_k = \{x_{k,n} : n = 1, 2, 3, \dots\}, \text{ where } x_{k,n} = \prod_{i=0}^{2^k-1} (i + n2^k, i + 2^k + 2^k - 1)$$

for each  $n \geq 0$ . Thus for example,

$$\begin{aligned} T_1 &= \{(01), (23), (45), \dots\}, T_2 = \{(02)(13), (46)(57), (810)(911), \dots\} \\ T_3 &= \{(04)(15)(26)(37), (812)(913)(1014)(1115), \dots\}. \end{aligned}$$

For each  $k \geq 1$  let  $G_k = \langle T_1, \dots, T_k \rangle$ . Then  $G = \bigcup_{k=1}^{\infty} G_k$ . Clearly  $G$  is a transitive 2 -subgroup of  $FSym(\mathbb{N})$ . It is easy to see that each  $G_k$  is normal in  $G$  and every orbit of it is finite. Furthermore each  $T_k$  is a set of disjoint involutions which are conjugate in  $G$  (see [16, p. 468]).

Next we show that  $G'$  is not a minimal non  $FC$ -group. Let  $X = \langle x \in G' : x^2 = 1 \rangle$ . Then  $X \triangleleft G$ . Let  $k \geq 1$ . Then for each  $n \geq 0$  there exists  $g \in G$  such that  $x_{k,n} = g^{-1}x_{k,0}g \in T_k$ , and hence  $x_{k,0}x_{k,n} = x_{k,0}g^{-1}x_{k,0}g \in X$  for all  $n \geq 0$  since any two elements of  $T_k$  commute which implies that  $x_{k,n} \in x_{k,0}X$ . Clearly it follows from this that  $G/X = \langle x_{k,0}X : k \geq 1 \rangle$ . We claim that  $G/X$  is abelian. Let  $k, t \geq 1$ . Clearly  $x_{t,0}x_{k,0}x_{t,0}x_{k,0} \in X$  as above. Hence it follows that  $x_{k,0}x_{t,0}X = (x_{k,0}x_{t,0})(x_{t,0}x_{k,0}x_{t,0}x_{k,0}X = x_{t,0}x_{k,0}X$  which shows that  $G/X$  is abelian and so  $G' \leq X$ . Since  $X \leq G'$  it follows that  $G' = X$  and so  $G'$  cannot be a minimal non  $FC$ -group by Corollary 1.9. Note that  $G' \neq G$  since  $G'$  is generated by even permutations and so  $T_1 \cap G' = \emptyset$ .

### 3. Proofs of Theorems 4, 5

**Lemma 3.1** *Let  $G$  be a totally imprimitive subgroup of  $FSym(\Omega)$ , where  $\Omega$  is infinite, and let  $A$  be a generating subset for  $G$ . Then there exists  $a \in A$  such that  $\langle a \rangle^G$  is not abelian.*

**Proof.** Assume that  $\langle a \rangle^G$  is abelian for all  $a \in A$ . Let  $F$  be a non-abelian finite subgroup of  $G$  and let  $\Delta$  be a non-trivial block for  $G$  such that  $supp(F) \subseteq \Delta$ . Since  $\langle A \rangle = G$  and  $G_{\{\Delta\}} \neq G$ , there exists  $a \in A \setminus G_{\{\Delta\}}$ . Then  $F' \leq [F, a] \leq \langle a \rangle^G$  and so  $\langle a \rangle^G \leq G_{\{\Delta\}}$  by Lemma 2.1(b), which is a contradiction.  $\square$

**Lemma 3.2** *Let  $G$  be a subgroup of  $FSym(\Omega)$ ,  $F$  a subgroup of  $G$  and  $\Delta$  be a non-trivial block for  $G$  such that  $supp(F) \subseteq \Delta$ . Let  $y \in G \setminus G_{\{\Delta\}}$  and let  $t$  be the smallest positive integer such that  $y^t \in G_{\{\Delta\}}$ . Assume that  $y^t \in FC_G(F)$ . Then there exists  $x \in F$  such that  $(yx^{-1})^t \in C_G(F)$ , and  $t$  is the smallest positive integer with this property.*

**Proof.** By hypothesis there exists  $h \in F$  and  $c \in C_G(F)$  such that  $y^t = ch$ . Also  $y(\Delta), \dots, y^{t-1}(\Delta)$  are pairwise disjoint by the choice of  $t$ . Let  $x \in F$  and  $i \in \Delta$ . We claim that  $(yx)^k(i) = y^k(x(i))$  for all  $1 \leq k \leq t$ . Assume that it holds for  $1 \leq k < t$ . Then

$$(yx)^{k+1}(i) = (yx)((yx)^k(i)) = (yx)(y^k(x(i))) = y(y^k(x(i))) = y^{k+1}(x(i))$$

since  $y^k(x(i)) \notin \Delta$  and  $supp(F) \subseteq \Delta$ . Thus the induction is complete. Now letting  $k = t$  and  $x = h^{-1}$  we get

$$(yh^{-1})^t(i) = y^t(h^{-1}(i)) = (y^t h^{-1})(i) = c(i).$$

Since  $i$  is any element of  $\Delta$  it follows that  $(yh^{-1})^t|_{\Delta} = c|_{\Delta}$ . Hence  $(yh^{-1})^t = (c|_{\Delta})d$  where  $d \in FSym(\Omega \setminus \Delta)$  which implies that  $(yh^{-1})^t \in C_G(F)$ . Also it is clear from the induction that  $t$  is the smallest such number.  $\square$

**Proof of Theorem 1.5** Let  $G$  be a totally imprimitive  $p$ -subgroup of  $FSym(\Omega)$ , where  $\Omega$  is infinite. Suppose that every orbit of every proper subgroup of  $G$  is finite. Assume that for every non-normal finite subgroup  $F$  of  $G$  there exists  $y \in G \setminus N_G(F)$  such that  $y^p \in C_G(F)$ . By Lemma 3.1 there exists  $c_1 \in G$  such that  $\langle c_1 \rangle^G$  is not abelian. Put  $F_1 = \langle c_1 \rangle$  and let  $\Lambda_1$  be a member of (1) containing  $supp(F_1)$ . Let  $U_1$  be the normal

closure of  $F_1$  in  $G_{\{\Delta\}}$ . Then  $\text{supp}(U_1) \subseteq \Delta_1$  by Lemma 2.3(a) and also  $N_G(U_1) = G_{\{\Lambda_1\}}$  by Lemma 2.1(a). By the hypothesis there exists  $c_2 \in G \setminus G_{\{\Lambda_1\}}$  such that  $c_2^p \in C_G(U_1)$ . In particular,  $c_2^p$  centralizes  $F_1$  since  $F_1 \subseteq U_1$ . Therefore

$$F_1^{\langle c_2 \rangle} = \langle c_1 \rangle \times \langle c_1 \rangle^{c_2} \times \cdots \times \langle c_1 \rangle^{c_2^{p-1}},$$

by Lemma 2.3(b). Next, put  $F_2 = F_1^{\langle c_2 \rangle} \langle c_2 \rangle = \langle F_1, c_2 \rangle$ . Let  $\Lambda_2$  be a member of (1) containing  $\text{supp}(F_2)$  and  $U_2$  be the normal closure of  $\text{supp}(F_2)$  in  $G_{\{\Lambda_2\}}$ . Then, as in the first case, there exists  $c_3 \in G \setminus G_{\{\Lambda_2\}}$  such that  $c_3^p \in C_G(U_2)$  and so  $F_2^{\langle c_3 \rangle} = F_2 \times F_2^{c_3} \times \cdots \times F_2^{c_3^{p-1}}$ . Put  $F_3 = F_2^{\langle c_3 \rangle} \langle c_3 \rangle = \langle F_2, c_3 \rangle$ . Continuing in this way we obtain properly increasing infinite chains

$$\Lambda_1 \subset \Lambda_2 \subset \cdots \text{ and } F_1 \subset F_2 \subset \cdots$$

of non-trivial blocks  $\Lambda_i$  and finite subgroups  $F_i = \langle c_1, c_2, \dots, c_i \rangle$  of  $G$  such that  $c_{i+1} \in G \setminus G_{\{\Lambda_i\}}$  for every  $i \geq 1$ . Obviously  $\Omega = \bigcup_{k=1}^{\infty} \Lambda_k$ . Put  $F = \bigcup_{i=1}^{\infty} F_i$ . Clearly  $F = G$  since the orbit of  $F$  containing any element of  $\text{supp}(c_1)$  is infinite by the choice of the  $c_i$ .

We claim that  $\langle c_1 \rangle^F$  is abelian. It suffices to show that  $\langle c_1 \rangle^{F_i}$  is abelian for all  $i \geq 1$ . This is obvious for  $i = 1$  since  $F_1 = \langle c_1 \rangle$ . Assume that it holds for some  $i \geq 1$ . We must show that it holds for  $i + 1$ . Thus  $\langle c_1 \rangle^{F_i}$  is abelian by assumption. Now

$$F_{i+1} = (F_i^{\langle c_{i+1} \rangle} \langle c_{i+1} \rangle \text{ and } F_i^{\langle c_{i+1} \rangle} = F_i \times F_i^{c_{i+1}} \times \cdots \times F_i^{c_{i+1}^{p-1}}.$$

Hence we can write  $F_{i+1}$  as  $F_{i+1} = (F_i \times F_i^{c_{i+1}} \times \cdots \times F_i^{c_{i+1}^{p-1}}) \langle c_{i+1} \rangle$ . Let  $x, y \in F_{i+1}$ . It suffices to show that  $[c_1^x, c_1^y] = 1$ . Now there exists  $a_0, a_1, \dots, a_{p-1}$  and  $b_0, b_1, \dots, b_{p-1} \in F_i$  and  $0 \leq r, s \leq p - 1$  such that

$$x = (a_0 a_1^{c_{i+1}} \cdots a_{p-1}^{c_{i+1}^{p-1}}) c_{i+1}^r \text{ and } y = (b_0 b_1^{c_{i+1}} \cdots b_{p-1}^{c_{i+1}^{p-1}}) c_{i+1}^s.$$

Since  $c_1^{a_0} \in F_i$  and  $[F_i, F_i^{c_{i+1}^u}] = 1$  for  $1 \leq u \leq p - 1$  by Lemma 2.1(d), it is easy to see that  $c_1^x = c_1^{a_0 c_{i+1}^r}$ . Similarly  $c_1^y = c_1^{b_0 c_{i+1}^s}$ . Now  $[c_1^x, c_1^y] = 1$  if  $r \neq s$  by Lemma 2.1(d), since  $[F_i^{c_{i+1}^r}, F_i^{c_{i+1}^s}] = 1$  due to the fact that  $0 \leq r, s \leq p - 1$ . If  $r = s$ , then  $[c_1^x, c_1^y] = [c_1^{a_0}, c_1^{b_0}]^{c_{i+1}^r} = 1$  since  $\langle c_1 \rangle^{F_i}$  is abelian. This completes the induction and so

it follows that  $\langle c_1 \rangle^F$  is abelian. But since  $\langle c_1 \rangle^G$  is not abelian this contradicts the fact that  $F = G$  and so the proof is complete.  $\square$

**Proof of Corollary 1.6** Let  $F$  be a non-normal finite subgroup of  $G$  and  $\Delta$  be a minimal block such that  $\text{supp}(F) \subseteq \Delta$ . By hypothesis there exists  $y \in G \setminus G_{\{\Delta\}}$  such that  $y^p \in FC_G(F)$ . By Lemma 3.2 there exists  $x \in F$  such that  $(yx)^p \in C_G(F)$ . Also  $yx \notin G_{\{\Delta\}}$  since  $x \in G_{\{\Delta\}}$ . Thus it follows that  $yx \notin N_G(F)$  but  $(yx)^p \in C_G(F)$  since  $N_G(F) \leq G_{\{\Delta\}}$  by Lemma 2.1. Since  $F$  is any finite non-normal subgroup of  $G$  applying Theorem 1.5 yields a proper subgroup of  $G$  that has an infinite orbit.  $\square$

**Proof of Corollary 1.7** Let  $k \geq 1$  and  $H = G_{\{\Delta_k\}}$ . Let  $M_k$  be the kernel of the representation of  $G$  into  $FSym(\Sigma_k)$ , where  $\Sigma_k = \{x(\Delta_k) : x \in G\}$ . Then  $M_k$  is the largest normal subgroup of  $G$  contained in  $H$  and so  $M_k = \langle F_k^G \rangle$  by hypothesis. Put  $\bar{G} = G/M_k$ . Let  $\bar{R}$  be a proper normal subgroup of  $\bar{G}$  such that  $\bar{R} \neq 1$ . Since  $\bar{R}$  is nilpotent  $\Omega_1(Z(\bar{R})) \neq 1$  and so it is not contained in  $\bar{H}$  by definition of  $M_k$ . Choose  $\bar{z} \in \Omega_1(Z(\bar{R})) \setminus \bar{H}$  such that  $o(\bar{z}) = p$ . Then  $z \notin H$  but  $z^p \in M_k$ . Since  $M_k = \langle F_k^G \rangle = \prod_{x \in G} F^x \leq F_k G_{\Delta_k}$  it follows that  $z^p \in F_k C_G(F_k)$ .

Let now  $F$  be any finite non - normal subgroup of  $G$ . There exists  $k \geq 1$  such that  $\text{supp}(F) \subseteq \Delta_k$  and so by the above paragraph there there exists  $y \in G \setminus N_G(F)$  such that  $y^p \in FC_G(F)$ . Therefore Corollary 1.6 yields a proper subgroup of  $G$  that has an infinite orbit.  $\square$

**Proof of Corollary 1.8** Let  $F$  be a non - normal finite subgroup of  $G$ . There exists a  $k \geq 1$  such that  $\text{supp}(F) \subseteq \Delta_k$ . By hypothesis there exists  $y \in G \setminus G_{\{\Delta_k\}}$  such that  $y^p \in G_{\Delta_k} \leq C_G(F)$ . Since  $N_G(F) \leq G_{\{\Delta_k\}}$ , the hypothesis of Theorem 1.5 is satisfied and so  $G$  contains a proper subgroup having an infinite orbit.  $\square$

**Proof of Theorem 1.11** Let  $G$  be a locally finite  $p$  - group that is also a minimal no  $FC$  - group. Then every proper subgroup of  $G$  is an  $FC$ -group. Assume that  $G$  is perfect and satisfies the hypothesis of the theorem. Then  $Z(G/Z(G)) = 1$ . Put  $\bar{G} = G/Z(G)$ . Then  $Z(\bar{G}) = 1$ . Since  $G$  is locally nilpotent, it has a proper normal subgroup  $\bar{N} \neq 1$ . Then since  $Z(\bar{N}) \neq 1$ , we can choose  $\bar{a} \in Z(\bar{N})$  with  $o(\bar{a}) = p$ . Put  $\Omega = \{(\bar{a})^{\bar{x}} : \bar{x} \in \bar{G}\}$ . Clearly  $\Omega$  is infinite and the conjugation action of  $\bar{G}$  on  $\Omega$  defines a finitary permutation group on  $\Omega$  by [3, Theorem1] or [8, Theorem]. Let  $K$  be the kernel of this representation. Then

$\bar{G}/\bar{K}$  is isomorphic to a totally imprimitive  $p$ -subgroup of  $FSym(\Omega)$ . Since  $\bar{G}/\bar{K} \cong G/K$  we may consider  $G/K$  instead of  $\bar{G}/\bar{K}$ .

Now we show that  $G/K$  satisfies the hypothesis of Corollary 1.6. By hypothesis for every non-normal finite subgroup  $F$  of  $G$  there exists  $y \in G \setminus N_G(F)$  such that  $y^p \in FC_G(F)$ . Let  $X/K$  be a finite non-normal subgroup of  $G/K$  and let  $T$  be a finite subgroup of  $X$  such that  $X = TK$ . Let  $\Delta$  be a non-trivial block for  $G/K$  such that  $supp(X/K) \subseteq \Delta$ . Put  $L/K = (G/K)_{\{\Delta\}}$ . Let  $V$  be the normal closure of  $T$  in  $L$ . Then  $V$  is a finite normal subgroup of  $L$  and  $X \leq VK$ . Thus  $L/K = N_{L/K}(VK/K)$  since  $supp(VK/K) \subseteq \Delta$ . In particular  $N_G(V) = L$ . By hypothesis there exists  $y \in G \setminus L$  such that  $y^p \in VC_G(V)$ . Then  $yK \notin L/K$  but  $y^pK \in (VK/K)C_{G/K}(X/K)$  since  $X/K \leq VK/K$ . Thus it follows that  $G/K$  satisfies the hypothesis of Corollary 1.6. But then  $G/K$  contains a proper subgroup having an infinite orbit, which is impossible by [5, Lemma 3.8D] or [16, Theorem 1] since  $G/K$  is a minimal non  $FC$ -group. This contradiction completes the proof of the theorem.  $\square$

The author is grateful to the referee for a careful reading of the manuscript and pointing out some errors.

### References

- [1] Arıkan A.: On barely transitive  $p$ -groups with soluble point stabilizer. J. Group Theory 5, 441-442 (2002).
- [2] Asar, A. O.: Barely transitive locally nilpotent  $p$ -groups. J. London Math. Soc.(2) 55, 357-362(1997)
- [3] Belyaev, V. V.: On the question of existence of minimal non  $FC$ -groups. Siberian Math. J.39, 1093-1095 (1998).
- [4] Belyaev, V. V., Kuzucuoglu, M.: Locally finite barely transitive groups. Algebra and Logic 42, 147-152 (2003)
- [5] Dixon, J. D., Mortimer, B.: Permutation Groups. New York, Springer-Ferlag, 1996.
- [6] Hartly, B., Kuzucuoglu, M.: Non-simplicity of locally finite barely transitive groups. Proceeding of the Edinburg Math. Soc. 40, 483-490 (1997).
- [7] Kuzucuoglu, M.: On torsion-free barely transitive groups. Turk. J. Math. 24, 273-276 (2000).



- [8] Leinen, F.: A reduction theorem for perfect locally finite minimal non  $FC$ -groups. Glasgow Math. J. 4, 81-83(1999).
- [9] Leinen, F., Puglisi, O.: Finitary representations and images of transitive finitary permutation groups. J. Algebra 222, 524-549 (1999).
- [10] Neumann, P. M.: The lawlessness of groups of finitary permutations. Arch. Math. 26, 561-566 (1975).
- [11] Neumann, P. M.: The structure of finitary permutation groups. Arch. Math. 27, 3-17 (1976).
- [12] Pinnock, C. J. E.: Irreducible and transitive locally-nilpotent by abelian groups. Arch. Math. 74, 168-172 (2000).
- [13] Robinson, D. J. S.: A Course in the Theory of Groups. New York, Heidelberg, Berlin. Springer-Verlag, 1980
- [14] Segal, D.: Normal subgroups of finitary permutation groups. Math. Z. 140, 81-85 (1974).
- [15] Suprunenko, D. A.: Locally nilpotent subgroups of infinite symmetric groups. Soviet Math. Dokl. 7, 392-394 (1966).
- [16] Wiegold, J.: Groups of finitary permutations. Arch. Math. 25, 466-469 (1974).

Ali Osman ASAR  
Gazi University, Gazi Education Faculty  
06500 Teknikokullar, Ankara-TURKEY  
e-mail: aliasar@gazi.edu.tr

Received 17.02.2005