

1-1-2005

On Generalization of The Quasi Homogeneous Riesz Potential

HÜSEYİN YILDIRIM

Follow this and additional works at: <https://journals.tubitak.gov.tr/math>



Part of the [Mathematics Commons](#)

Recommended Citation

YILDIRIM, HÜSEYİN (2005) "On Generalization of The Quasi Homogeneous Riesz Potential," *Turkish Journal of Mathematics*: Vol. 29: No. 4, Article 4. Available at: <https://journals.tubitak.gov.tr/math/vol29/iss4/4>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Mathematics by an authorized editor of TÜBİTAK Academic Journals. For more information, please contact academic.publications@tubitak.gov.tr.

On Generalization of The Quasi Homogeneous Riesz Potential

Hüseyin Yıldırım

Abstract

In this paper, a generalization of the quasi homogeneous Riesz Potential has been defined using non-isotropic quasi-distance and its L_p ($p \geq 1$) continuity study.

Key words and phrases: Non-isotropic quasi-distance, Riesz Potential, Quasi-metric.

1. Introduction

Continuity properties of the classical Riesz Potential was studied in [1],[2],[3] and [4]. In this article we have defined a generalization of the quasi homogeneous Riesz Potential and we studied the L_p ($p \geq 1$)– continuity of this potential.

2. Preliminaries

Firstly, we give some notations and definitions. We define a quasimetric (a non-isotropic quasi-distance) in \mathbb{R}^n by

$$\|x - y\|_{\lambda, \gamma} = \left(|x_1 - y_1|^{\frac{2\lambda_1}{\gamma}} + \dots + |x_n - y_n|^{\frac{2\lambda_n}{\gamma}} \right)^{\frac{\gamma}{2n}} \left| \frac{1}{\lambda} \right|, \quad (1)$$

where $\lambda_1, \dots, \lambda_n$ and γ are the positive real numbers and $\left| \frac{1}{\lambda} \right| = \frac{1}{\lambda_1} + \dots + \frac{1}{\lambda_n}$. This quasimetric is named non-isotropic quasi-distance [5].

2000 Mathematics Subject Classification: 31B10, 46E30, 42B35 and 47B37

For $\rho \in \mathbb{R}$ and $x \in \mathbb{R}^n$ define $\rho^{\frac{\gamma}{2\lambda}} x = (\rho^{\frac{\gamma}{2\lambda_1}} x_1, \rho^{\frac{\gamma}{2\lambda_2}} x_2, \dots, \rho^{\frac{\gamma}{2\lambda_n}} x_n)$.

1. $\|x\|_{\lambda, \gamma} = 0 \Leftrightarrow x = \theta$
2. $\|\rho^{\frac{\gamma}{2\lambda}} x\|_{\lambda, \gamma} = |\rho|^{\frac{\gamma}{2n} |\frac{1}{\lambda}|} \|x\|_{\lambda, \gamma}$
3. $\|x + y\|_{\lambda, \gamma} \leq C(\|x\|_{\lambda, \gamma} + \|y\|_{\lambda, \gamma})$,

where $\lambda_{\max} = \max\{\lambda_1, \dots, \lambda_n\}$ and $C = 2^{(2\lambda_{\max} + \gamma) \frac{1}{2n} |\frac{1}{\lambda}|}$. It is clear that if $\lambda_1 = \dots = \lambda_n = 2$ and $\gamma = 2$ then, quasimetric (a non-isotropic quasi-distance) is the Euclidean metric (Euclidean norm) on \mathbb{R}^n .

For a $f : \mathbb{R}^n \rightarrow \mathbb{R}$ and a $K : \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R}^+$ we write

$$(I_{\varphi}^{\lambda, \gamma} f)(x) = \int_{\mathbb{R}^n} f(y) K(x, y) dy, \tag{2}$$

where

$$K(x, y) = \frac{1}{\varphi(\|x - y\|_{\lambda, \gamma})}. \tag{3}$$

For $\mathbb{R}_0^+ = [0, \infty)$, function $\varphi : \mathbb{R}_0^+ \rightarrow \mathbb{R}_0^+$ has the finite derivative and the following are valid:

$$\varphi(0) = 0 \tag{4}$$

$$\frac{\varphi'(t)}{\varphi(t)} \searrow \text{ on } (0, \Delta) \ (\Delta > 0) \tag{5}$$

$$\frac{t \varphi'(t)}{\varphi(t)} \sim c > 0, t \rightarrow +0 \tag{6}$$

$$\int_0^r \frac{t^{\gamma |\frac{1}{\lambda}| - 1}}{\varphi^p(t)} dt = O\left(\frac{r^{\gamma |\frac{1}{\lambda}|}}{\varphi^p(r)}\right), r \rightarrow +0 \tag{7}$$

for any $\beta \in \mathbb{R}_0^+$,

$$\varphi(\beta t) = \beta^m \varphi(t), \quad m \in \mathbb{R}. \tag{8}$$

The operator $I_\varphi^{\lambda, \gamma} f$, when

$$\varphi(t) = t^{n-\alpha}, \quad 0 < \alpha < n, t \in [0, +\infty) \tag{9}$$

is called the generalized Riesz potential with non-isotropic quasi-distance of function f . If φ satisfies the conditions (4), (5), (6), (7) and (8), then $I_\varphi^{\lambda, \gamma} f$ is the generalization of the following quasi homogeneous Riesz potential

$$I^{\lambda, \gamma} f = \int_{\mathbb{R}^n} \|x - y\|_{\lambda, \gamma}^{\alpha-n} f(y) dy. \tag{10}$$

The L_p -continuity, or continuity in the L_p norm, of a real function g at point $x \in \mathbb{R}^n$ thus:

$$\left\{ \frac{1}{|B_r|} \int_{B_r} |g(x+s) - g(x)|^p ds \right\}^{\frac{1}{p}} \rightarrow 0, \quad r \rightarrow +0, \tag{11}$$

where $|B_r|$ is the volume of a ball of radius r with center at zero [2].

Here we consider spherical coordinates by the following formulas:

$$y_1 = (\rho \cos \theta_1)^{\frac{\gamma}{\lambda_1}}, \dots, y_n = (\rho \sin \theta_1, \sin \theta_2, \dots, \sin \theta_{n-1})^{\frac{\gamma}{\lambda_n}},$$

where we obtained that $\|x\|_\lambda = \rho^{\frac{\gamma}{\lambda}} |\frac{1}{\lambda}|$. It can be seen that the Jacobian $J_\lambda(\rho, \theta)$ of this transformation is $J_\lambda(\rho, \theta) = \rho^{\gamma |\frac{1}{\lambda}| - 1} \Omega_\lambda(\theta)$, where $\Omega_\lambda(\theta)$ is the bounded function and

$$\Omega_\lambda(\theta) = \gamma^n \cdot \frac{1}{\lambda_1} \frac{1}{\lambda_2} \dots \frac{1}{\lambda_n} (\cos \theta_1)^{\frac{\gamma}{\lambda_1} - 1} \cdot (\cos \theta_2)^{\frac{\gamma}{\lambda_2} - 1} \dots (\cos \theta_{n-1})^{\frac{\gamma}{\lambda_n} - 1} (\sin \theta_1)^{\frac{1}{\lambda_1} \frac{1}{\lambda_2} \dots \frac{1}{\lambda_n} - 1} \dots (\sin \theta_1)^{\frac{1}{\lambda_n} - 1}$$

Now we will prove that, depending on conditions imposed on f and φ , the operator $I_\varphi^{\lambda, \gamma} f$ is L_p continuous at every point at which it exists.

Theorem 2.1: If $p \geq 1$ and $I_\varphi^{\lambda, \gamma} f$ exists at point $x \in \mathbb{R}^n$, then $I_\varphi^{\lambda, \gamma} f$ is L_p continuous at x .

Proof: From (11), it is sufficient to consider the case when $x = 0$ and to prove that ([2] and [6])

$$\left\{ \int_{\|s\|_{\lambda,\gamma} < r} |(I_{\varphi}^{\lambda,\gamma} f)(s) - (I_{\varphi}^{\lambda,\gamma} f)(0)| ds \right\}^{\frac{1}{p}} = o(r^{\frac{2}{p}|\frac{1}{\lambda}|}) \quad (12)$$

holds when $r \rightarrow +0$. From (12), (2) and (3), we have following inequality:

$$\begin{aligned} & \left\{ \int_{\|s\|_{\lambda,\gamma} < r} |(I_{\varphi}^{\lambda,\gamma} f)(s) - (I_{\varphi}^{\lambda,\gamma} f)(0)|^p ds \right\}^{\frac{1}{p}} \\ &= \left\{ \int_{\|s\|_{\lambda,\gamma} < r} \left| \int_{\mathbb{R}^n} \left[\frac{1}{\varphi(\|s-y\|_{\lambda,\gamma})} - \frac{1}{\varphi(\|y\|_{\lambda,\gamma})} \right] f(y) dy \right|^p ds \right\}^{\frac{1}{p}} \\ &\leq \left\{ \int_{\|s\|_{\lambda,\gamma} < r} \left| \int_{\|y\|_{\lambda,\gamma} > 2r} \left[\frac{1}{\varphi(\|s-y\|_{\lambda,\gamma})} - \frac{1}{\varphi(\|y\|_{\lambda,\gamma})} \right] f(y) dy \right|^p ds \right\}^{\frac{1}{p}} \\ &+ \left\{ \int_{\|s\|_{\lambda,\gamma} < r} \left| \int_{\|y\|_{\lambda,\gamma} < 2r} \left[\frac{1}{\varphi(\|s-y\|_{\lambda,\gamma})} - \frac{1}{\varphi(\|y\|_{\lambda,\gamma})} \right] f(y) dy \right|^p ds \right\}^{\frac{1}{p}} \quad (13) \\ &\leq \left\{ \int_{\|s\|_{\lambda,\gamma} < r} \left| \int_{\|y\|_{\lambda,\gamma} < 2r} \left[\frac{|f(y)|}{\varphi(\|s-y\|_{\lambda,\gamma})} \right] dy \right|^p ds \right\}^{\frac{1}{p}} \\ &+ \left\{ \int_{\|s\|_{\lambda,\gamma} < r} \left| \int_{\|y\|_{\lambda,\gamma} < 2r} \left[\frac{|f(y)|}{\varphi(\|y\|_{\lambda,\gamma})} \right] dy \right|^p ds \right\}^{\frac{1}{p}} \\ &+ \left\{ \int_{\|s\|_{\lambda,\gamma} < r} \left| \int_{\|y\|_{\lambda,\gamma} > 2r} \left[\frac{1}{\varphi(\|s-y\|_{\lambda,\gamma})} - \frac{1}{\varphi(\|y\|_{\lambda,\gamma})} \right] |f(y)| dy \right|^p ds \right\}^{\frac{1}{p}} \\ &= \mathbf{I}_1 + \mathbf{I}_2 + \mathbf{I}_3. \end{aligned}$$

By Minkowsky inequality and (7) we have

$$\begin{aligned}
 \mathbf{I}_1 &\leq \int_{\|y\|_{\lambda,\gamma} < 2r} |f(y)| \left(\int_{\|s\|_{\lambda,\gamma} < r} \frac{1}{\varphi^p(\|s-y\|_{\lambda,\gamma})} ds \right)^{\frac{1}{p}} dy \\
 &\leq \int_{\|y\|_{\lambda,\gamma} < 2r} |f(y)| \left(\int_{\|y-s\|_{\lambda} \leq 3Cr} \frac{ds}{\varphi^p(\|s-y\|_{\lambda,\gamma})} \right)^{\frac{1}{p}} dy \\
 &= \int_{\|y\|_{\lambda,\gamma} < 2r} |f(y)| \left(|B_{1,\lambda}| \int_0^{3Cr} \frac{t^{\gamma|\frac{1}{\lambda}|-1}}{\varphi^p(t)} dt \right)^{\frac{1}{p}} dy \\
 &= O \left(\frac{(3Cr)^{\frac{\gamma}{p}|\frac{1}{\lambda}|}}{\varphi^p(3Cr)} \varphi(2r) \int_{\|y\|_{\lambda,\gamma} < 2r} \frac{|f(y)|}{\varphi(\|y\|_{\lambda,\gamma})} dy \right) = o(r^{\frac{\gamma}{p}|\frac{1}{\lambda}|}), \quad (r \rightarrow 0),
 \end{aligned} \tag{14}$$

since the following is valid:

$$\int_{\|x\|_{\lambda,\gamma} < r} h(\|x\|_{\lambda,\gamma}) dx = |B_{1,\lambda}| \int_0^r t^{\gamma|\frac{1}{\lambda}|-1} h(t) dt, \tag{15}$$

if $t^{\gamma|\frac{1}{\lambda}|-1}h(t)$ is non-negative and measurable or integrable function at $(0, r)$, where, $|B_{1,\lambda}|$ denote the volume of the ball of $B_{1,\lambda}(x) = \{x \in \mathbb{R}^n : \|x\|_{\lambda,\gamma} < 1\}$.

For the \mathbf{I}_2

$$\mathbf{I}_2 \leq \int_{\|y\|_{\lambda,\gamma} < 2r} \frac{|f(y)|}{\varphi(\|y\|_{\lambda,\gamma})} \left(\int_{\|s\|_{\lambda,\gamma} < r} ds \right)^{\frac{1}{p}} dy = o \left(\int_{\|s\|_{\lambda,\gamma} < r} ds \right)^{\frac{1}{p}} = o \left(r^{\frac{\gamma}{p}|\frac{1}{\lambda}|} \right), \quad r \rightarrow +0. \tag{16}$$

If we use the Minkowski inequality at the \mathbf{I}_3 , then we obtain,

$$\begin{aligned}
 \mathbf{I}_3 &= \left\{ \int_{\|s\|_{\lambda,\gamma} < r} \left| \int_{\|y\|_{\lambda,\gamma} > 2r} \left[\frac{1}{\varphi(\|s-y\|_{\lambda,\gamma})} - \frac{1}{\varphi(\|y\|_{\lambda,\gamma})} \right] |f(y)| dy \right|^p ds \right\}^{\frac{1}{p}} \\
 &= O \left\{ \int_{\|s\|_{\lambda,\gamma} < r} \left| \int_{\|y\|_{\lambda,\gamma} > 2r} \frac{\varphi'(\|y\|_{\lambda,\gamma})}{\varphi^2(\|y\|_{\lambda,\gamma})} \|s\|_{\lambda,\gamma} |f(y)| dy \right|^p ds \right\}^{\frac{1}{p}} \\
 &\leq O \left\{ \int_{\|y\|_{\lambda,\gamma} > 2r} \frac{\varphi'(\|y\|_{\lambda,\gamma})}{\varphi^2(\|y\|_{\lambda,\gamma})} |f(y)| \left[\int_{\|s\|_{\lambda,\gamma} < r} \|s\|_{\lambda,\gamma}^p ds \right]^{\frac{1}{p}} dy \right\} \tag{17} \\
 &\leq O \left\{ \int_{\|y\|_{\lambda,\gamma} > 2r} \frac{\varphi'(\|y\|_{\lambda,\gamma})}{\varphi^2(\|y\|_{\lambda,\gamma})} |f(y)| \cdot r \cdot \left(\int_0^r t^{\gamma|\frac{1}{\lambda}|-1} dt \right)^{\frac{1}{p}} dy \right\} \\
 &= O \left\{ r^{\frac{2}{p}|\frac{1}{\lambda}|+1} \int_{\|y\|_{\lambda,\gamma} > 2r} \frac{\varphi'(\|y\|_{\lambda,\gamma})}{\varphi^2(\|y\|_{\lambda,\gamma})} |f(y)| dy \right\}.
 \end{aligned}$$

for $0 < r < \min\{1, \Delta\}$, from (5) and (6) follows:

$$\begin{aligned}
 \int_{\|y\|_{\lambda,\gamma} > 2r} \frac{\varphi'(\|y\|_{\lambda,\gamma})|f(y)|}{\varphi(\|y\|_{\lambda,\gamma})^2} dy &= \left(\int_{\|y\|_{\lambda,\gamma} > 2\sqrt{r}} + \int_{2\sqrt{r} \geq \|y\|_{\lambda,\gamma} > 2r} \right) \frac{\varphi'(\|y\|_{\lambda,\gamma})|f(y)|}{\varphi(\|y\|_{\lambda,\gamma})^2} dy \\
 &\leq \frac{\varphi'(2\sqrt{r})}{\varphi(2\sqrt{r})} \int_{\|y\|_{\lambda,\gamma} > 2\sqrt{r}} \frac{|f(y)|}{\varphi(\|y\|_{\lambda,\gamma})} dy \\
 &\quad + \frac{\varphi'(2r)}{\varphi(2r)} \int_{2\sqrt{r} \geq \|y\|_{\lambda,\gamma} > 2r} \frac{|f(y)|}{\varphi(\|y\|_{\lambda,\gamma})} dy \tag{18} \\
 &\leq \frac{\varphi'(2\sqrt{r})}{\varphi(2\sqrt{r})} \int_{\mathbb{R}^n} \frac{|f(y)|}{\varphi(\|y\|_{\lambda,\gamma})} dy \\
 &\quad + \frac{\varphi'(2r)}{\varphi(2r)} \int_{\|y\|_{\lambda,\gamma} < 2\sqrt{r}} \frac{|f(y)|}{\varphi(\|y\|_{\lambda,\gamma})} dy \\
 &= O\left(\frac{1}{2\sqrt{r}}\right) + o\left(\frac{1}{r}\right) = o\left(\frac{1}{r}\right).
 \end{aligned}$$

Then from (17) and (18) follows:

$$\mathbf{I}_3 = o\left(r^{\frac{2}{p}|\frac{1}{\lambda}|}\right), \quad (r \rightarrow +0) \tag{19}$$

Hence Theorem is obtained from (14), (16) and (19). \square

Specially, if φ is defined by $\varphi = t^{n-\alpha}$, $\gamma = 2$ and $\lambda_1 = \dots = \lambda_n = 2$, then $I_\varphi f$ is classical Riesz potential of f . If φ is defined by $\varphi = t^{n-\alpha}$, i.e. if $I_\varphi^{\lambda,\gamma} f$ is a non-isotrop

Riesz potentials of f , then the conditions (4),(5),(6),(7) and (8) are valid if $1 - \frac{1}{p} < \frac{\alpha}{p} < 1$, and if $I_{\varphi}^{\lambda, \gamma} f$ exists at point $x \in \mathbb{R}^n$, then the function $I_{\varphi}^{\lambda, \gamma} f$ is L_p continuous at x .

References

- [1] Mizuta Y.: Continuity properties of Riesz Potentials and boundary limits of Beppo Levi function, Math. Scand, 63(1988), 238-260.
- [2] Zigmund, A.: Trigonometric series, I-II, Cambridge, (1968).
- [3] Divnić, T. and Djurić, Z.: A Generalization of the Riesz Potential, Kragujevac J. Math., 22(2000)83-86.
- [4] Stein, E.M.: Singular Integrals Differential Properties of Functions, Princeton University Press, Princeton, New Jersey. (1970).
- [5] Fabes, E.B. and Riviere, N.M.: Symbolic Calculus of Kernels with Mixed Homogeneity, Proceeding of Symposion in Pure Math. P:107-127, 1967.
- [6] Love, E.R. and Tuan, V.K.: L^p -Continuity of Riesz potentials, Integral Trasforms and Special Functions, Vol. 1, No 1, pp 27-31, (1993).

Hüseyin YILDIRIM
 Department of Mathematics,
 Faculty of Sciences and Arts,
 Kocatepe University,
 A.N. Sezer Campus, Afyon - Turkey
 e-mail: hyildir@aku.edu.tr

Received 13.07.2004