

1-1-2005

On δ -I-Continuous Functions

ŞAZİYE YÜKSEL

A. AÇIKGÖZ

T. NOIRI

Follow this and additional works at: <https://journals.tubitak.gov.tr/math>



Part of the [Mathematics Commons](#)

Recommended Citation

YÜKSEL, ŞAZİYE; AÇIKGÖZ, A.; and NOIRI, T. (2005) "On δ -I-Continuous Functions," *Turkish Journal of Mathematics*: Vol. 29: No. 1, Article 4. Available at: <https://journals.tubitak.gov.tr/math/vol29/iss1/4>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Mathematics by an authorized editor of TÜBİTAK Academic Journals. For more information, please contact academic.publications@tubitak.gov.tr.

On δ -I-Continuous Functions

S. Yüksel, A. Açıkgöz and T. Noiri

Abstract

In this paper, we introduce a new class of functions called δ -I-continuous functions. We obtain several characterizations and some of their properties. Also, we investigate its relationship with other types of functions.

Key words and phrases: δ -I-cluster point, R-I-open set, δ -I-continuous, strongly θ -I-continuous, almost-I-continuous, SI-R space, AI-R space.

1. Introduction

Throughout this paper $Cl(A)$ and $Int(A)$ denote the closure and the interior of A , respectively. Let (X, τ) be a topological space and let I an ideal of subsets of X . An ideal is defined as a nonempty collection I of subsets of X satisfying the following two conditions: (1) If $A \in I$ and $B \subset I$, then $B \in I$; (2) If $A \in I$ and $B \in I$, then $A \cup B \in I$. An ideal topological space is a topological space (X, τ) with an ideal I on X and is denoted by (X, τ, I) . For a subset $A \subset X$, $A^*(I) = \{x \in X \mid U \cap A \notin I \text{ for each neighborhood } U \text{ of } x\}$ is called the local function of A with respect to I and τ [4]. We simply write A^* instead of $A^*(I)$ to be brief. X^* is often a proper subset of X . The hypothesis $X = X^*[1]$ is equivalent to the hypothesis $\tau \cap I = \emptyset$ [5]. For every ideal topological space (X, τ, I) , there exists a topology $\tau^*(I)$, finer than τ , generated by $\beta(I, \tau) = \{U \setminus I \mid U \in \tau \text{ and } I \in I\}$, but in general $\beta(I, \tau)$ is not always a topology [2]. Additionally, $Cl^*(A) = A \cup (A^*)^*$ defines a Kuratowski closure operator for $\tau^*(I)$.

AMS Mathematics Subject Classification: 54C10.

In this paper, we introduce the notions of δ -I-open sets and δ -I-continuous functions in ideal topological spaces. We obtain several characterizations and some properties of δ -I-continuous functions. Also, we investigate the relationships with other related functions.

2. δ -I-open sets

In this section, we introduce δ -I-open sets and the δ -I-closure of a set in an ideal topological space and investigate their basic properties. It turns out that they have similar properties with δ -open sets and the δ -closure due to Veličko [6].

Definition 2.1 *A subset A of an ideal topological space (X, τ, I) is said to be an R -I-open (resp. regular open) set if $\text{Int}(Cl^*(A)) = A$ (resp. $\text{Int}(Cl(A)) = A$). We call a subset A of X R -I-closed if its complement is R -I-open.*

Definition 2.2 *Let (X, τ, I) be an ideal topological space, S a subset of X and x a point of X .*

(1) *x is called a δ -I-cluster point of S if $S \cap \text{Int}(Cl^*(U)) \neq \emptyset$ for each open neighborhood U of x ;*

(2) *The family of all δ -I-cluster points of S is called the δ -I-closure of S and is denoted by $[S]_{\delta-I}$ and*

(3) *A subset S is said to be δ -I-closed if $[S]_{\delta-I} = S$. The complement of a δ -I-closed set of X is said to be δ -I-open.*

Lemma 2.1 *Let A and B be subsets of an ideal topological space (X, τ, I) . Then, the following properties hold:*

- (1) *$\text{Int}(Cl^*(A))$ is R -I-open;*
- (2) *If A and B are R -I-open, then $A \cap B$ is R -I-open;*
- (3) *If A is regular open, then it is R -I-open;*
- (4) *If A is R -I-open, then it is δ -I-open and*
- (5) *Every δ -I-open set is the union of a family of R -I-open sets.*

Proof. (1) Let A be a subset of X and $V = \text{Int}(Cl^*(A))$. Then, we have $\text{Int}(Cl^*(V)) = \text{Int}(Cl^*(\text{Int}(Cl^*(A)))) \subset \text{Int}(Cl^*(Cl^*(A))) = \text{Int}(Cl^*(A)) = V$ and also $V = \text{Int}(V) \subset \text{Int}(Cl^*(V))$. Therefore, we obtain $\text{Int}(Cl^*(V)) = V$.

(2) Let A and B be R-I-open. Then, we have $A \cap B = \text{Int}(Cl^*(A)) \cap \text{Int}(Cl^*(B)) = \text{Int}(Cl^*(A) \cap Cl^*(B)) \supset \text{Int}(Cl^*(A \cap B)) \supset \text{Int}(A \cap B) = A \cap B$. Therefore, we obtain $A \cap B = \text{Int}(Cl^*(A \cap B))$. This shows that $A \cap B$ is R-I-open.

(3) Let A be regular open. Since $\tau^* \supset \tau$, we have $A = \text{Int}(A) \subset \text{Int}(Cl^*(A)) \subset \text{Int}(Cl(A)) = A$ and hence $A = \text{Int}(Cl^*(A))$. Therefore, A is R-I-open.

(4) Let A be any R-I-open set. For each $x \in A$, $(X-A) \cap A = \emptyset$ and A is R-I-open. Hence $x \notin [X-A]_{\delta-I}$ for each $x \in A$. This shows that $x \notin (X-A)$ implies $x \notin [X-A]_{\delta-I}$. Therefore, we have $[X-A]_{\delta-I} \subset (X-A)$. Since in general, $S \subset [S]_{\delta-I}$ for any subset S of X , $[X-A]_{\delta-I} = (X-A)$ and hence A is δ -I-open.

(5) Let A be a δ -I-open set. Then $(X-A)$ is δ -I-closed and hence $(X-A) = [X-A]_{\delta-I}$. For each $x \in A$, $x \notin [X-A]_{\delta-I}$ and there exists an open neighborhood V_x such that $\text{Int}(Cl^*(V_x)) \cap (X-A) = \emptyset$. Therefore, we have $x \in V_x \subset \text{Int}(Cl^*(V_x)) \subset A$ and hence $A = \cup \{\text{Int}(Cl^*(V_x)) \mid x \in A\}$. By (1), $\text{Int}(Cl^*(V_x))$ is R-I-open for each $x \in A$. \square

Lemma 2.2 *Let A and B be subsets of an ideal topological space (X, τ, I) . Then, the following properties hold:*

- (1) $A \subset [A]_{\delta-I}$;
- (2) If $A \subset B$, then $[A]_{\delta-I} \subset [B]_{\delta-I}$;
- (3) $[A]_{\delta-I} = \cap \{F \subset X \mid A \subset F \text{ and } F \text{ is } \delta\text{-I-closed}\}$;
- (4) If A is a δ -I-closed set of X for each $\alpha \in \Delta$, then $\cap \{A_\alpha \mid \alpha \in \Delta\}$ is δ -I-closed;
- (5) $[A]_{\delta-I}$ is δ -I-closed.

Proof. (1) For any $x \in A$ and any open neighborhood V of x , we have $\emptyset \neq A \cap V \subset A \cap \text{Int}(Cl^*(V))$ and hence $x \in [A]_{\delta-I}$. This shows that $A \subset [A]_{\delta-I}$.

(2) Suppose that $x \notin [B]_{\delta-I}$. There exists an open neighborhood V of x such that $\emptyset = \text{Int}(Cl^*(V)) \cap B$; hence $\text{Int}(Cl^*(V)) \cap A = \emptyset$. Therefore, we have $x \notin [A]_{\delta-I}$.

(3) Suppose that $x \in [A]_{\delta-I}$. For any open neighborhood V of x and any δ -I-closed set F containing A , we have $\emptyset \neq A \cap \text{Int}(Cl^*(V)) \subset F \cap \text{Int}(Cl^*(V))$ and hence $x \in [F]_{\delta-I} = F$. This shows that $x \in \cap \{F \subset X \mid A \subset F \text{ and } F \text{ is } \delta\text{-I-closed}\}$. Conversely, suppose that $x \notin [A]_{\delta-I}$. There exists an open neighborhood V of x such that $\text{Int}(Cl^*(V)) \cap A = \emptyset$. By Lemma 2.1, $X - \text{Int}(Cl^*(V))$ is a δ -I-closed set which contains A and does not contain x . Therefore, we obtain $x \notin \cap \{F \subset X \mid A \subset F \text{ and } F \text{ is } \delta\text{-I-closed}\}$. This completes the proof. \square

(4) For each $\alpha \in \Delta$, $[\bigcap_{\alpha \in \Delta} A_\alpha]_{\delta-I} \subset [A_\alpha]_{\delta-I} = A_\alpha$ and hence $[\bigcap_{\alpha \in \Delta} A_\alpha]_{\delta-I} \subset [\bigcap_{\alpha \in \Delta} A_\alpha]$. By (1), we obtain $[\bigcap_{\alpha \in \Delta} A_\alpha]_{\delta-I} = [\bigcap_{\alpha \in \Delta} A_\alpha]$. This shows that $\bigcap_{\alpha \in \Delta} A_\alpha$ is δ -I-closed.

(5) This follows immediately from (3) and (4).

A point x of a topological space (X, τ) is called a δ -cluster point of a subset S of X if $\text{Int}(\text{Cl}(V)) \cap S \neq \emptyset$ for every open set V containing x . The set of all δ -cluster points of S is called the δ -closure of S and is denoted by $Cl_\delta(S)$. If $Cl_\delta(S) = S$, then S is said to be δ -closed [6]. The complement of a δ -closed set is said to be δ -open. It is well-known that the family of regular open sets of (X, τ) is a basis for a topology which is weaker than τ . This topology is called the *semi-regularization* of τ and is denoted by τ_S . Actually, τ_S is the same as the family of δ -open sets of (X, τ) .

Theorem 2.1 *Let (X, τ, I) be an ideal topological space and $\tau_{\delta-I} = \{A \subset X \mid A \text{ is a } \delta\text{-I-open set of } (X, \tau, I)\}$. Then $\tau_{\delta-I}$ is a topology such that $\tau_S \subset \tau_{\delta-I} \subset \tau$.*

Proof. By Lemma 2.1, we obtain $\tau_S \subset \tau_{\delta-I} \subset \tau$. Next, we show that $\tau_{\delta-I}$ is a topology.

(1) It is obvious that $\emptyset, X \in \tau_{\delta-I}$.

(2) Let $V_\alpha \in \tau_{\delta-I}$ for each $\alpha \in \Delta$. Then $X - V_\alpha$ is δ -I-closed for each $\alpha \in \Delta$. By Lemma 2.2, $\bigcap_{\alpha \in \Delta} (X - V_\alpha)$ is δ -I-closed and $\bigcap_{\alpha \in \Delta} (X - V_\alpha) = X - \bigcup_{\alpha \in \Delta} V_\alpha$. Hence $\bigcup_{\alpha \in \Delta} V_\alpha$ is δ -I-open.

(3) Let $A, B \in \tau_{\delta-I}$. By Lemma 2.1, $A = \bigcup_{\alpha \in \Delta_1} A_\alpha$ and $B = \bigcup_{\beta \in \Delta_2} B_\beta$, where A_α and B_β are R-I-open sets for each $\alpha \in \Delta_1$ and $\beta \in \Delta_2$. Thus $A \cap B = \bigcup \{A_\alpha \cap B_\beta \mid \alpha \in \Delta_1, \beta \in \Delta_2\}$. Since $A_\alpha \cap B_\beta$ is R-I-open, $A \cap B$ is a δ -I-open set by Lemma 2.1. \square

Example 2.1 *Let $X = \{a, b, c, d\}$, $\tau = \{\emptyset, X, \{a\}, \{c\}, \{a, c\}\}$ and $I = \{\emptyset, \{a\}\}$. Then $A = \{a, c\}$ is a δ -I-open set which is not R-I-open. Since $\{a\}$ and $\{c\}$ are regular open sets, A is a δ -open set and hence δ -I-open. But A is not R-I-open. Because $A^* = \{b, c, d\}$ and $Cl^*(A) = A \cup A^* = X$. Therefore, we have $\text{Int}(Cl^*(A)) = X \neq A$.*

For some special ideals, we have the following properties.

Proposition 2.1 *Let (X, τ, I) be an ideal topological space.*

(1) *If $I = \{\emptyset\}$ or the ideal N of nowhere dense sets of (X, τ) , then $\tau_{\delta-I} = \tau_S$.*

(2) *If $I = P(X)$, then $\tau_{\delta-I} = \tau$.*

Proof. (1) Let $I = \{\emptyset\}$, then $S^* = Cl(S)$ for every subset S of X . Let A be R - I -open. Then $A = Int(Cl^*(A)) = (A \cup A^*) = Int(Cl(A))$ and hence A is regular open. Therefore, every δ - I -open set is δ -open and we obtain $\tau_{\delta-I} \subset \tau_S$. By Theorem 2.1, we obtain $\tau_{\delta-I} = \tau_S$. Next, Let $I = N$. It is well-know that $S^* = Cl(Int(Cl(S)))$ for every subset S of X . Let A be any R - I -open set. Then since A is open, $A = Int(Cl^*(A)) = Int(A \cup A^*) = Int(A \cup Cl(Int(Cl(A)))) = Int(Cl(Int(Cl(A)))) = Int(Cl(A))$. Hence A is regular open. Similarly to the case of $I = \{\emptyset\}$, we obtain $\tau_{\delta-I} = \tau_S$.

(2) Let $I = P(X)$. Then $S^* = \emptyset$ for every subset S of X . Now, let A be any open set of X . Then $A = Int(A) = Int(A \cup A^*) = Int(Cl^*(A))$ and hence A is R - I -open. By Theorem 2.1, we obtain $\tau_{\delta-I} = \tau$. \square

3. δ - I -continuous functions

Definition 3.1 A function $f:(X,\tau,I) \rightarrow (Y,\Phi,J)$ is said to be δ - I -continuous if for each $x \in X$ and each open neighborhood V of $f(x)$, there exists an open neighborhood U of x such that $f(Int(Cl^*(U))) \subset Int(Cl^*(V))$.

Theorem 3.1 For a function $f:(X,\tau,I) \rightarrow (Y,\Phi,J)$, the following properties are equivalent:

- (1) f is δ - I -continuous;
- (2) For each $x \in X$ and each R - I -open set V containing $f(x)$, there exists an R - I -open set containing x such that $f(U) \subset V$;
- (3) $f([A]_{\delta-I}) \subset [f(A)]_{\delta-I}$ for every $A \subset X$;
- (4) $[f^{-1}(B)]_{\delta-I} \subset f^{-1}([B]_{\delta-I})$ for every $B \subset Y$;
- (5) For every δ - I -closed set F of Y , $f^{-1}(F)$ is δ - I -closed in X ;
- (6) For every δ - I -open set V of Y , $f^{-1}(V)$ is δ - I -open in X ;
- (7) For every R - I -open set V of Y , $f^{-1}(V)$ is δ - I -open in X ;
- (8) For every R - I -closed set F of Y , $f^{-1}(F)$ is δ - I -closed in X .

Proof. (1) \Rightarrow (2): This follows immediately from Definition 3.1.

(2) \Rightarrow (3): Let $x \in X$ and $A \subset X$ such that $f(x) \in [A]_{\delta-I}$. Suppose that $f(x) \notin [f(A)]_{\delta-I}$. Then, there exists an R - I -open neighborhood V of $f(x)$ such that $f(A) \cap V = \emptyset$. By (2), there exists an R - I -open neighborhood U of x such that $f(U) \subset V$. Since $f(A) \cap f(U) \subset f(A) \cap V = \emptyset$, $f(A) \cap f(U) = \emptyset$. Hence, we get that $U \cap A \subset f^{-1}(f(U)) \cap f^{-1}(f(A)) = f^{-1}(f(U) \cap f(A))$

$= \emptyset$. Hence we have $U \cap A = \emptyset$ and $x \notin [A]_{\delta-I}$. This shows that $f(x) \notin f([A]_{\delta-I})$. This is a contradiction. Therefore, we obtain that $f(x) \in [f(A)]_{\delta-I}$.

(3) \Rightarrow (4): Let $B \subset Y$ such that $A = f^{-1}(B)$. By (3), $f([f^{-1}(B)]_{\delta-I}) \subset [f(f^{-1}B)]_{\delta-I} \subset [B]_{\delta-I}$. From here, we have $[f^{-1}(B)]_{\delta-I} \subset f^{-1}([f(f^{-1}(B))]_{\delta-I}) \subset f^{-1}([B]_{\delta-I})$. Thus we obtain that $[f^{-1}(B)]_{\delta-I} \subset f^{-1}([B]_{\delta-I})$.

(4) \Rightarrow (5): Let $F \subset Y$ be δ -I-closed. By (4), $[f^{-1}(F)]_{\delta-I} \subset f^{-1}([F]_{\delta-I}) = f^{-1}(F)$ and always $f^{-1}(F) \subset [f^{-1}(F)]_{\delta-I}$. Hence we obtain that $[f^{-1}(F)]_{\delta-I} = f^{-1}(F)$. This shows that $f^{-1}(F)$ is δ -I-closed.

(5) \Rightarrow (6): Let $V \subset Y$ be δ -I-open. Then $Y-V$ is δ -I-closed. By (5), $f^{-1}(Y-V) = X-f^{-1}(V)$ is δ -I-closed. Therefore, $f^{-1}(V)$ is δ -I-open.

(6) \Rightarrow (7): Let $V \subset Y$ be R-I-open. Since every R-I-open set is δ -I-open, V is δ -I-open, By (6), $f^{-1}(V)$ is δ -I-open.

(7) \Rightarrow (8): Let $F \subset Y$ be an R-I-closed set. Then $Y-F$ is R-I-open. By (7), $f^{-1}(Y-F) = X-f^{-1}(F)$ is δ -I-open. Therefore, $f^{-1}(F)$ is δ -I-closed.

(8) \Rightarrow (1): Let $x \in X$ and V be an open set containing $f(x)$. Now, set $V_o = \text{Int}(Cl^*(V))$, then by Lemma 2.1 $Y-V_o$ is an R-I-closed set. By (8), $f^{-1}(Y-V_o) = X-f^{-1}(V_o)$ is a δ -I-closed set. Thus we have $f^{-1}(V_o)$ is δ -I-open. Since $x \in f^{-1}(V_o)$, by Lemma 2.1, there exists an open neighborhood U of x such that $x \in U \subset \text{Int}(Cl^*(U)) \subset f^{-1}(V_o)$. Hence we obtain that $f(\text{Int}(Cl^*(U))) \subset \text{Int}(Cl^*(V))$. This shows that f is a δ -I-continuous function.

□

Corollary 3.1 *A function $f:(X,\tau,I) \rightarrow (Y,\Phi,J)$ is δ -I-continuous if and only if $f:(X,\tau,I) \rightarrow (Y,\Phi,J)$ is continuous.*

Proof. This is an immediate consequence of Theorem 2.1.

The following lemma is known in [3, as Lemma 4.3].

□

Lemma 3.1 *Let (X,τ,I) be an ideal topological space and A,B subsets of X such that $B \subset A$. Then $B^*(\tau/A,I/A) = B^*(\tau,I) \cap A$.*

Proposition 3.1 *Let (X, τ, I) be an ideal topological space, A, X_o subsets of X such that $A \subset X_o$ and X_o is open in X .*

- (1) *If A is R-I-open in (X, τ, I) , then A is R-I-open in $(X_o, \tau/X_o, I/X_o)$,*
- (2) *If A is δ -I-open in (X, τ, I) , then A is δ -I-open in $(X_o, \tau/X_o, I/X_o)$.*

Proof. (1) Let A be R-I-open in (X, τ, I) . Then $A = \text{Int}(Cl^*(A))$ and $Cl_{X_o}^*(A) = A \cup A^*(\tau/X_o, I/X_o) = A \cup [A^*(\tau, I) \cap X_o] = (A \cap X_o) \cup (A^* \cap X_o) = (A \cup A^*) \cap X_o = Cl^*(A) \cap X_o$. Hence we have $\text{Int}_{X_o}(Cl_{X_o}^*(A)) = \text{Int}(Cl_{X_o}^*(A)) = \text{Int}((Cl^*(A) \cap X_o)) = \text{Int}((Cl^*(A)) \cap X_o) = A$. Therefore, A is R-I-open in $(X_o, \tau/X_o, I/X_o)$.

(2) Let A be a δ -I-open set of (X, τ, I) . By Lemma 2.1, $A = \bigcup_{\alpha \in \Delta} A_\alpha$, where A_α is R-I-open set of (X, τ, I) for each $\alpha \in \Delta$. By (1), A is R-I-open in $(X_o, \tau/X_o, I/X_o)$ for each $\alpha \in \Delta$ and hence A is δ -I-open in $(X_o, \tau/X_o, I/X_o)$. \square

Theorem 3.2 *If $f: (X, \tau, I) \rightarrow (Y, \Phi, J)$ is a δ -I-continuous function and X_o is a δ -I-open set of (X, τ, I) , then the restriction $f/X_o: (X_o, \tau/X_o, I/X_o) \rightarrow (Y, \Phi, J)$ is δ -I-continuous.*

Proof. Let V be any δ -I-open set of (Y, Φ, J) . Since f is δ -I-continuous, $f^{-1}(V)$ is δ -I-open in (X, τ, I) . Since X_o is δ -I-open, by Theorem 2.1 $X_o \cap f^{-1}(V)$ is δ -I-open in (X, τ, I) and hence $X_o \cap f^{-1}(V)$ is δ -I-open in $(X_o, \tau/X_o, I/X_o)$ by Proposition 3.1. This shows that $(f/X_o)^{-1}(V)$ is δ -I-open in $(X_o, \tau/X_o, I/X_o)$ and hence f/X_o is δ -I-continuous. \square

Theorem 3.3 *If $f: (X, \tau, I) \rightarrow (Y, \Phi, J)$ and $g: (Y, \Phi, J) \rightarrow (Z, \varphi, K)$ are δ -I-continuous, then so is $g \circ f: (X, \tau, I) \rightarrow (Z, \varphi, K)$.*

Proof. It follows immediately from Cor. 3.1. \square

Theorem 3.4 *If $f, g: (X, \tau, I) \rightarrow (Y, \Phi, J)$ are δ -I-continuous functions and Y is a Hausdorff space, then $A = \{x \in X : f(x) = g(x)\}$ is a δ -I-closed set of (X, τ, I) .*

Proof. We prove that $X-A$ is δ -I-open set. Let $x \in X-A$. Then, $f(x) \neq g(x)$. Since Y is Hausdorff, there exist open sets V_1 and V_2 containing $f(x)$ and $g(x)$, respectively, such that $V_1 \cap V_2 = \emptyset$. From here we have $\text{Int}(Cl(V_1)) \cap \text{Int}(Cl(V_2)) = \emptyset$. Thus, we obtain that $\text{Int}(Cl^*(V_1)) \cap \text{Int}(Cl^*(V_2)) = \emptyset$. Since f and g are δ -I-continuous, there exists an open

neighborhood U of x such that $f(\text{Int}(Cl^*(U))) \subset \text{Int}(Cl^*(V_1))$ and $g(\text{Int}(Cl^*(U))) \subset \text{Int}(Cl^*(V_2))$. Hence we obtain that $\text{Int}(Cl^*(U)) \subset f^{-1}(\text{Int}(Cl^*(V_1)))$ and $\text{Int}(Cl^*(U)) \subset g^{-1}(\text{Int}(Cl^*(V_2)))$. From here we have $\text{Int}(Cl^*(U)) \subset f^{-1}(\text{Int}(Cl^*(V_1))) \cap g^{-1}(\text{Int}(Cl^*(V_2)))$. Moreover $f^{-1}(\text{Int}(Cl^*(V_1))) \cap g^{-1}(\text{Int}(Cl^*(V_2))) \cap A = \emptyset$. Suppose that $f^{-1}(\text{Int}(Cl^*(V_1))) \cap g^{-1}(\text{Int}(Cl^*(V_2))) \cap A \neq \emptyset$. Hence there exists a point z such that $z \in f^{-1}(\text{Int}(Cl^*(V_1))) \cap g^{-1}(\text{Int}(Cl^*(V_2))) \cap A$. Thus, $f(z) \in \text{Int}(Cl^*(V_1))$, $g(z) \in \text{Int}(Cl^*(V_2))$ and $z \in A$. Since $z \in A$, $f(z) = g(z)$. Therefore, we have $f(z) \in \text{Int}(Cl^*(V_1)) \cap \text{Int}(Cl^*(V_2))$ and $\text{Int}(Cl^*(V_1)) \cap \text{Int}(Cl^*(V_2)) \neq \emptyset$. This is a contradiction to $\text{Int}(Cl^*(V_1)) \cap \text{Int}(Cl^*(V_2)) = \emptyset$. Hence we obtain that $f^{-1}(\text{Int}(Cl^*(V_1))) \cap g^{-1}(\text{Int}(Cl^*(V_2))) \cap A = \emptyset$. Thus $f^{-1}(\text{Int}(Cl^*(V_1))) \cap g^{-1}(\text{Int}(Cl^*(V_2))) \subset X - A$. Since $\text{Int}(Cl^*(U)) \subset f^{-1}(\text{Int}(Cl^*(V_1))) \cap g^{-1}(\text{Int}(Cl^*(V_2)))$, we have that there exists an open neighborhood of x such that $x \in U$ $\text{Int}(Cl^*(U)) \subset X - A$. Therefore, $X - A$ is a δ -I-open set. This shows that A is δ -I-closed. \square

4. Comparisons

Definition 4.1 A function $f: (X, \tau, I) \rightarrow (Y, \Phi, J)$ is said to be strongly θ -I-continuous (resp. θ -I-continuous, almost I-continuous) if for each $x \in X$ and each open neighborhood V of $f(x)$, there exists an open neighborhood U of x such that $f(Cl^*(U)) \subset V$ (resp. $f(Cl^*(U)) \subset Cl^*(V)$, $f(U) \subset \text{Int}(Cl^*(V))$).

Definition 4.2 A function $f: (X, \tau, I) \rightarrow (Y, \Phi, J)$ is said to be almost-I-open if for each R -I-open set U of X , $f(U)$ is open in Y .

Theorem 4.1 (1) If $f: (X, \tau, I) \rightarrow (Y, \Phi, J)$ is strongly θ -I-continuous and $g: (Y, \Phi, J) \rightarrow (Z, \varphi, K)$ is almost I-continuous, then $g \circ f: (X, \tau, I) \rightarrow (Z, \varphi, K)$ is δ -I-continuous.

(2) The following implications hold:

$$\text{strongly } \theta - I - \text{continuous} \Rightarrow \delta - I - \text{continuous} \Rightarrow \text{almost} - I - \text{continuous}. \quad (4.1)$$

Proof. (1) Let $x \in X$ and W be any open set of Z containing $(g \circ f)(x)$. Since g is almost I-continuous, there exists an open neighborhood $V \subset Y$ of $f(x)$ such that $g(V) \subset \text{Int}(Cl^*(W))$.

Since f is strongly θ -I-continuous, there exists an open neighborhood $U \subset X$ of x such that $f(Cl^*(U)) \subset V$. Hence we have $g(f(Cl^*(U))) \subset g(V)$ and $g(f(Int(Cl^*(U)))) \subset g(f(Cl^*(U))) \subset g(V) \subset Int(Cl^*(W))$. Thus, we obtain $g(f(Int(Cl^*(U)))) \subset Int(Cl^*(W))$. This shows that $g \circ f$ is δ -I-continuous.

(2) Let f be strongly θ -I-continuous. Let $x \in X$ and V be any open neighborhood of $f(x)$. Then, there exists an open neighborhood $U \subset X$ of x such that $f(Cl^*(U)) \subset V$. Since always $f(Int(Cl^*(U))) \subset f(Cl^*(U))$, $f(Int(Cl^*(U))) \subset V$. Since V is open, we have $f(Int(Cl^*(U))) \subset Int(Cl^*(V))$. Thus, f is δ -I-continuous. Let f be δ -I-continuous. Now we prove that f is almost I-continuous. Then, for each $x \in X$ and each open neighborhood $V \subset Y$ of $f(x)$, there exists an open neighborhood $U \subset X$ of x such that $f(Int(Cl^*(U))) \subset Int(Cl^*(V))$. Since $U \subset Int(Cl^*(U))$, $f(U) \subset Int(Cl^*(V))$. Thus, f is almost I-continuous. \square

Remark 4.1 *The following examples enable us to realize that none of these implications in Theorem 4.1 (2) is reversible.*

Example 4.1 *Let $X = \{a, b, c\}$, $\tau = \{\emptyset, X, \{a\}, \{a, c\}\}$, $I = \{\emptyset, \{c\}\}$, $\Phi = \{\emptyset, X, \{a, b\}\}$ and $J = \{\emptyset, \{b\}, \{c\}, \{b, c\}\}$. The identity function $f: (X, \tau, I) \rightarrow (X, \Phi, J)$ is δ -I-continuous but it is not strongly θ -I-continuous.*

(i) *Let $a \in X$ and $V = \{a, b\} \in \Phi$ such that $f(a) \in V$. $V^* = (\{a, b\})^* = \{a, b, c\} = X$, $Cl^*(V) = V \cup V^* = X$ and $Int(Cl^*(V)) = Int(X) = X$. Then, there exists an open $U = \{a, c\} \subset X$ such that $a \in U$. We have $U^* = (\{a\})^* = \{a, b, c\}$, $Cl^*(U) = U \cup U^* = \{a, b, c\}$ and $Int(Cl^*(U)) = \{a, c\}$. Since $f(Int(Cl^*(U))) = f(\{a, c\}) = \{a, c\}$ and $\{a, c\} \subset Int(Cl^*(V)) = X$.*

(ii) *Let $b \in X$ and $V = \{a, b\} \in \Phi$ such that $f(b) \in V$. $V^* = (\{a, b\})^* = \{a, b, c\} = X$, $Cl^*(V) = V \cup V^* = X$ and $Int(Cl^*(V)) = Int(X) = X$. Then, there exists an open $U = X$ such that $b \in U$. We have $Cl^*(U) = Cl^*(X) = X$ and $Int(Cl^*(U)) = Int(X)$. Since $f(Int(Cl^*(U))) = f(X) = X$ and $X \subset Int(Cl^*(V)) = X$.*

(iii) *Let $x = a, b$ or c and $V = X \in \Phi$ such that $f(x) \in V$. $Cl^*(V) = V \cup V^* = X$ and $Int(Cl^*(V)) = Int(X) = X$. Then, there exists an open $U = X$ such that $x \in U$. We have $Cl^*(U) = Cl^*(X) = X$ and $Int(Cl^*(U)) = Int(X)$. Since $f(Int(Cl^*(U))) = f(X) = X$ and $X \subset Int(Cl^*(V)) = X$. By (i), (ii) and (iii), f is δ -I-continuous. On the other hand by (i), since $f(Cl^*(U)) = f(Cl^*(\{a\})) = f(\{a, b, c\}) = \{a, b, c\}$ is not subset of $V = \{a, b\}$, f is not strongly θ -I-continuous.*

Example 4.2 Let $X = \{a, b, c, d\}$, $\tau = \{\emptyset, X, \{a, c\}, \{d\}, \{a, b, c\}, \{a, c, d\}\}$, $I = \{\emptyset, \{d\}\}$ and $J = \{\emptyset, \{c\}, \{d\}, \{c, d\}\}$. The identity function $f: (X, \tau, I) \rightarrow (X, \tau, J)$ is almost I -continuous but it is not δ - I -continuous. (i) Let $x = a$ or $c \in X$ and $V = \{a, c\} \in \Phi = \tau$ such that $f(x) \in V$. $V^* = (\{a, c\})^* = \{a, b, c\}$, $Cl^*(V) = \bigvee V^* = \{a, b, c\}$ and $Int(Cl^*(V)) = \{a, c\}$. Then, there exists an open $U = \{a, c\} \subset X$ such that $x \in U$. We have $U^* = (\{a, c\})^* = \{a, b, c\}$ and $Int(Cl^*(U)) = Int(\{a, b, c\}) = \{a, b, c\}$. Since $f(U) = f(\{a, c\}) = \{a, c\} \subset Int(Cl^*(V)) = \{a, c\}$.

(ii) Let $x = a, c$ or $d \in X$ and $V = \{a, c, d\} \in \Phi = \tau$ such that $f(x) \in V$. $V^* = (\{a, c, d\})^* = \{a, b, c\}$ and $Cl^*(V) = \bigvee V^* = \{a, b, c, d\}$ and $Int(Cl^*(V)) = X$. Then, there exists an open $U = \{a, c, d\} \subset X$ such that $x \in U$. We have $U^* = (\{a, c, d\})^* = \{a, b, c, d\} = X$ and $Int(Cl^*(U)) = Int(X) = X$. Since $f(U) = f(\{a, c, d\}) = \{a, c, d\} \subset Int(Cl^*(V)) = \{a, b, c, d\}$.

(iii) Let $x = a, b$ or $c \in X$ and $V = \{a, b, c\} \in \Phi = \tau$ such that $f(x) \in V$. $V^* = (\{a, b, c\})^* = \{a, b, c\}$ and $Cl^*(V) = \bigvee V^* = \{a, b, c\}$ and $Int(Cl^*(V)) = \{a, b, c\}$. Then, there exists an open $U = \{a, b, c\} \subset X$ such that $x \in U$. We have $U^* = (\{a, b, c\})^* = \{a, b, c\}$ and $Int(Cl^*(U)) = \{a, b, c\}$. Since $f(U) = f(\{a, b, c\}) = \{a, b, c\} \subset Int(Cl^*(V)) = \{a, b, c\}$.

(iv) Let $d \in X$ and $V = \{d\} \in \Phi = \tau$ such that $f(d) \in V$. $V^* = (\{d\})^* = \emptyset$ and $Cl^*(V) = \bigvee V^* = \{d\}$ and $Int(Cl^*(V)) = \{d\}$. Then, there exists an open $U = \{d\} \subset X$ such that $d \in U$. We have $U^* = (\{d\})^* = \emptyset$ and $Int(Cl^*(U)) = \{d\}$. Since $f(U) = f(\{d\}) = \{d\} \subset Int(Cl^*(V)) = \{d\}$. By (i), (ii), (iii) and (iv), f is almost I -continuous. On the other hand by (i), since $f(Int(Cl^*(U))) = f(\{a, b, c\}) = \{a, b, c\}$ is not subset of $Int(Cl^*(V))$ and $Int(Cl^*(V)) = \{a, c\}$, f is not δ - I -continuous.

Definition 4.3 An ideal topological space (X, τ, I) is said to be an SI - R space if for each $x \in X$ and each open neighborhood V of x , there exists an open neighborhood U of x such that $x \in U \subset Int(Cl^*(U)) \subset V$.

Theorem 4.2 For a function $f: (X, \tau, I) \rightarrow (Y, \Phi, J)$, the following are true:

- (1) If Y is an SI - R space and f is δ - I -continuous, then f is continuous.
- (2) If X is an SI - R space and f is almost I -continuous, then f is δ - I -continuous.

Proof. (1) Let Y be an SI - R space. Then, for each open neighborhood V of $f(x)$, there exists an open neighborhood V_o of $f(x)$ such that $f(x) \in V_o \subset Int(Cl^*(V_o)) \subset V$. Since f is δ - I -continuous, there exists an open neighborhood U_o of x such that $f(Int(Cl^*(U_o))) \subset Int(Cl^*(V_o))$. Since U_o is an open set, $f(U_o) \subset f(Int(Cl^*(U_o))) \subset Int(Cl^*(V_o)) \subset V$. Thus, $f(U_o) \subset V$ and hence f is continuous.

(2) Let $x \in X$ and V be an open neighborhood of $f(x)$. Since f is almost I -continuous, there exists an open neighborhood U of x such that $f(U) \subset \text{Int}(Cl^*(V))$. Since X is an SI - R space, there exists an open neighborhood U_1 of x such that $\text{Int}(Cl^*(U_1)) \subset U$. Thus $f(\text{Int}(Cl^*(U_1))) \subset f(U) \subset \text{Int}(Cl^*(V))$. Therefore f is δ - I -continuous. \square

Corollary 4.1 *If (X, τ, I) and (Y, Φ, J) are SI - R spaces, then the following concepts on a function $f: (X, \tau, I) \rightarrow (Y, \Phi, J)$: δ - I -continuity, continuity and almost I -continuity are equivalent.*

Definition 4.4 *An ideal topological space (X, τ, I) is said to be an AI - R space if for each R - I -closed set $F \subset X$ and each $x \notin F$, there exist disjoint open sets U and V in X such that $x \in U$ and $F \subset V$.*

Theorem 4.3 *An ideal topological space (X, τ, I) is an AI - R space if and only if each $x \in X$ and each R - I -open neighborhood V of x , there exists an R - I -open neighborhood U of x such that $x \in U \subset Cl^*(U) \subset Cl(U) \subset V$.*

Proof. **Necessity.** Let $x \in V$ and V be R - I -open. Then $\{x\} \cap (X - V) = \emptyset$. Since X is an AI - R space, there exist open sets U_1 and U_2 containing x and $(X - V)$, respectively, such that $U_1 \cap U_2 = \emptyset$. Then $Cl(U_1) \cap U_2 = \emptyset$ and hence $Cl^*(U_1) \subset Cl(U_1) \subset (X - U_2) \subset V$. Thus $x \in U_1 \subset Cl^*(U_1) \subset Cl(U_1) \subset V$ and we obtain that $U_1 \subset \text{Int}(Cl^*(U_1)) \subset Cl^*(U_1)$. Let $\text{Int}(Cl^*(U_1)) = U$. Thus, we have $Cl(U) = Cl(\text{Int}(Cl^*(U_1))) \subset Cl(Cl^*(U_1)) \subset Cl(Cl(U_1)) = Cl(U_1) \subset Cl(U)$ and $U_1 \subset U \subset Cl^*(U) \subset Cl^*(U_1) \subset Cl(U_1) \subset V$. Therefore, there exists an R - I -open set U such that $x \in U \subset Cl^*(U) \subset Cl(U) \subset V$.

Sufficiency. Let $x \in X$ and an R - I -closed set F such that $x \notin F$. Then, $X - F$ is an R - I -open neighborhood of x . By hypothesis, there exists an R - I -open neighborhood V of x such that $x \in V \subset Cl^*(V) \subset Cl(V) \subset X - F$. From here we have $F \subset X - Cl(V) \subset (X - Cl^*(V))$, where $X - Cl(V)$ is an open set. Moreover, we have that $V \cap (X - Cl(V)) = \emptyset$ and V is open. Therefore, X is an AI - R space. \square

Theorem 4.4 *For a function $f: (X, \tau, I) \rightarrow (Y, \Phi, J)$, the following are true:*

- (1) *If Y is an AI - R space and f is θ - I -continuous, then f is δ - I -continuous.*

(2) If X is an AI-R space, Y is an SI-R space and f is δ -I-continuous, then f is strongly θ -I-continuous.

Proof. (1) Let Y be an AI-R space. Then, for each $x \in X$ and each R-I-open neighborhood V of $f(x)$, there exists an R-I-open neighborhood V_o of $f(x)$ such that $f(x) \in V_o \subset Cl^*(V_o) \subset V$. Since f is θ -I-continuous, there exists an open neighborhood U_o of x such that $f(Cl^*(U_o)) \subset Cl^*(V_o)$. Hence, we obtain that $f(Int(Cl^*(U_o))) \subset f(Cl^*(U_o)) \subset Cl^*(V_o) \subset V$ and thus $f(Int(Cl^*(U_o))) \subset V$. By Theorem 3.1, f is δ -I-continuous.

(2) Let X be an AI-R space and Y an SI-R space. For each $x \in X$ and each open neighborhood V of $f(x)$, there exists an open set V_o such that $f(x) \in V_o \subset Int(Cl^*(V_o)) \subset V$ since Y is an SI-R space. Since f is δ -I-continuous, there exists an open set U containing x such that $f(Int(Cl^*(U))) \subset Int(Cl^*(V_o))$. By Lemma 2.1, $Int(Cl^*(U))$ is R-I-open and since X is AI-R, by Theorem 4.3 there exists an R-I-open set U_o such that $x \in V_o \subset Cl^*(U_o) \subset Int(Cl^*(U))$. Every R-I-open set is open and hence U_o is open. Moreover, we have $f(Cl^*(U_o)) \subset V$. This shows that f is strongly θ -I-continuous. \square

Theorem 4.5 *If a function $f: (X, \tau, I) \rightarrow (Y, \Phi, J)$ is θ -I-continuous and almost-I-open, then it is δ -I-continuous.*

Proof. Let $x \in X$ and V be an open neighborhood of $f(x)$. Since f is θ -I-continuous, there exists an open neighborhood of x such that $f(Cl^*(U)) \subset Cl^*(V)$; therefore, $f(Int(Cl^*(U))) \subset Cl^*(V)$. Since f is almost-I-open, we have $f(Int(Cl^*(U))) \subset Int(Cl^*(V))$. This shows that f is δ -I-continuous. \square

References

- [1] E. Hayashi, Topologies defined by local properties, Math. Ann., 156(1964), 205-215.
- [2] D. Janković and T. R. Hamlett, New topologies from old via ideals, Amer. Math. Monthly, 97 (1990), 295-310.
- [3] D. Janković and T.R. Hamlett, Compatible extensions of ideals, Boll. Un. Mat. Ital.(7), 6-B (1992), 453-465.
- [4] K. Kuratowski, Topology Vol. 1 (transl.), Academic Press, New York, 1966.

YÜKSEL, AÇIKGÖZ, NOIRI

- [5] P. Samuels, A topology formed from a given topology and ideal, J. London Math. Soc. (2),10 (1975), 409-416.
- [6] N.V. Veličko, H-closed topological spaces, Amer. Math. Soc. Transl. (2), 78 (1968), 103-118.

S. YÜKSEL, A. AÇIKGÖZ
Department of Mathematics,
University of Selçuk,
42079 Konya-TURKEY
e-mail: syuksel@selcuk.edu.tr.

Received 03.09.2003

T. NOIRI
Department of Mathematics,
Yatsushiro College of Technology,
Yatsushiro, Kumamoto,
866-8501 JAPAN
e-mail: noiri@as.yatsushiro-nct.ac.jp