

1-1-2005

## Spacelike Normal Curves in Minkowski Space $E^3_1$

KAZIM İLARSLAN

Follow this and additional works at: <https://journals.tubitak.gov.tr/math>



Part of the [Mathematics Commons](#)

---

### Recommended Citation

İLARSLAN, KAZIM (2005) "Spacelike Normal Curves in Minkowski Space  $E^3_1$ ," *Turkish Journal of Mathematics*: Vol. 29: No. 1, Article 5. Available at: <https://journals.tubitak.gov.tr/math/vol29/iss1/5>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Mathematics by an authorized editor of TÜBİTAK Academic Journals. For more information, please contact [academic.publications@tubitak.gov.tr](mailto:academic.publications@tubitak.gov.tr).

## Spacelike Normal Curves in Minkowski Space $\mathbb{E}_1^3$

*Kazım İlarşlan*

Dedicated to Professor Dr. H. Hilmi Hacısalihoğlu

### Abstract

In the Euclidean space  $\mathbb{E}^3$ , it is well known that normal curves, i.e., curves with position vector always lying in their normal plane, are spherical curves [3]. Necessary and sufficient conditions for a curve to be a spherical curve in Euclidean 3-space are given in [10] and [11].

In this paper, we give some characterizations of spacelike normals curves with spacelike, timelike or null principal normal in the Minkowski 3-space  $\mathbb{E}_1^3$ .

**Key words and phrases:** Normal Curves, Position Vector and Minkowski Space.

### 1. Introduction

In the Euclidean space  $\mathbb{E}^3$ , it is well-known that to each unit speed curve  $\alpha : I \subset \mathbb{R} \rightarrow \mathbb{E}^3$  with at least four continuous derivatives, one can associate three mutually orthogonal unit vector fields  $T$ ,  $N$  and  $B$ , called respectively the tangent, the principal normal and the binormal vector fields. At each point  $\alpha(s)$  of curve  $\alpha$ , the planes spanned by  $\{T, N\}$ ,  $\{T, B\}$  and  $\{N, B\}$  are known respectively as the osculating plane, the rectifying plane and the normal plane. The curves  $\alpha : I \subset \mathbb{R} \rightarrow \mathbb{E}^3$  for which the position vector  $\alpha$  always lie in their rectifying plane, are for simplicity called *rectifying curves*, (see [3]). Similarly, the curves for which the position vector  $\alpha$  always lie in their osculating plane, are for simplicity called *osculating curves*; and finally, the curves for which the position vector

---

1991 *Mathematics Subject Classification*: 53C50, 53C40.

always lie in their normal plane, are for simplicity called *normal curves*. By definition, for a normal curve, the position vector  $\alpha$  satisfies

$$\alpha(s) = \lambda(s)N(s) + \mu(s)B(s),$$

for some differentiable functions  $\lambda$  and  $\mu$  of  $s \in I \subset \mathbb{R}$ .

Characterization of rectifying curves is given in [3] and these curves are studied in Minkowski space  $\mathbb{E}_1^3$  in [5]. In this paper, we characterize *spacelike normal curves*, lying fully in the Minkowski space  $\mathbb{E}_1^3$ .

## 2. Preliminaries

The Minkowski 3-space  $\mathbb{E}_1^3$  is the Euclidean 3-space  $\mathbb{E}^3$  provided with the standard flat metric given by

$$g = -dx_1^2 + dx_2^2 + dx_3^2,$$

where  $(x_1, x_2, x_3)$  is a rectangular coordinate system of  $\mathbb{E}_1^3$ .

Since  $g$  is an indefinite metric, recall that a vector  $v \in \mathbb{E}_1^3$  can have one of three Lorentzian causal characters: it can be spacelike if  $g(v, v) > 0$  or  $v = 0$ , timelike if  $g(v, v) < 0$  and null (lightlike) if  $g(v, v) = 0$  and  $v \neq 0$ . Similarly, an arbitrary curve  $\alpha = \alpha(s)$  in  $\mathbb{E}_1^3$  can locally be spacelike, timelike or null (lightlike), if all of its velocity vectors  $\alpha'(s)$  are respectively spacelike, timelike or null (lightlike). Denote by  $\{T, N, B\}$  the moving Frenet frame along the curve  $\alpha(s)$  in the space  $\mathbb{E}_1^3$ . For an arbitrary curve  $\alpha(s)$  in the space  $\mathbb{E}_1^3$ , the following Frenet formulae are given in [4, 9].

If  $\alpha$  is a spacelike curve with a spacelike or timelike principal normal  $N$ , then the Frenet formulae read

$$\begin{bmatrix} T' \\ N' \\ B' \end{bmatrix} = \begin{bmatrix} 0 & k_1 & 0 \\ -\epsilon k_1 & 0 & k_2 \\ 0 & k_2 & 0 \end{bmatrix} \begin{bmatrix} T \\ N \\ B \end{bmatrix}, \quad (1)$$

where  $g(T, T) = 1, g(N, N) = \epsilon = \pm 1, g(B, B) = -\epsilon, g(T, N) = 0, g(T, B) = 0, g(N, B) = 0$ .

If  $\alpha$  is a spacelike curve with a null (lightlike) principal normal  $N$ , the Frenet formulae

are

$$\begin{bmatrix} T' \\ N' \\ B' \end{bmatrix} = \begin{bmatrix} 0 & k_1 & 0 \\ 0 & k_2 & 0 \\ -k_1 & 0 & -k_2 \end{bmatrix} \begin{bmatrix} T \\ N \\ B \end{bmatrix}, \quad (2)$$

where  $g(T, T) = 1, g(N, N) = 0, g(B, B) = 0, g(T, N) = 0, g(T, B) = 0, g(N, B) = 1$ . In this case,  $k_1$  can take only two values:  $k_1 = 0$  when  $\alpha$  is a straight line;  $k_1 = 1$  in all other cases.

Let  $m$  be a fixed point in  $\mathbb{E}_1^3$  and  $r > 0$  be a constant. The pseudo-Riemannian sphere is defined by

$$\mathbb{S}_1^2(m, r) = \{u \in \mathbb{E}_1^3 : g(u - m, u - m) = r^2\};$$

the pseudo-Riemannian hyperbolical space is defined by

$$\mathbb{H}_0^2(m, r) = \{u \in \mathbb{E}_1^3 : g(u - m, u - m) = -r^2\};$$

the pseudo-Riemannian lightlike cone (quadric cone) is defined by

$$C(m) = \{u \in \mathbb{E}_1^3 : g(u - m, u - m) = 0\}.$$

### 3. The spacelike normal curves in $\mathbb{E}_1^3$

In this section, we give some characterization theorems for spacelike normal curves.

**Theorem 3.1** *Let  $\alpha = \alpha(s)$  be a unit speed spacelike normal curve in  $\mathbb{E}_1^3$  with spacelike or timelike principal normal  $N$  and with curvatures  $k_1(s) > 0, k_2(s) \neq 0$  for each  $s \in I \subset \mathbb{R}$ . Then the following statements hold:*

- (i) The curvatures  $k_1(s)$  and  $k_2(s)$  satisfy the following equality

$$\frac{1}{k_1(s)} = c_1 \cosh\left(\int k_2(s) ds\right) + c_2 \sinh\left(\int k_2(s) ds\right), \quad c_1, c_2 \in \mathbb{R};$$

- (ii) The principal normal and binormal component of the position vector of the curve are given respectively by

$$g(\alpha(s), N) = a_1 \cosh\left(\int k_2(s) ds\right) + a_2 \sinh\left(\int k_2(s) ds\right)$$

$$g(\alpha(s), B) = a_1 \sinh\left(\int k_2(s) ds\right) + a_2 \cosh\left(\int k_2(s) ds\right), \quad a_1, a_2 \in \mathbb{R};$$

(iii) If the position vector of the curve is null vector, then  $\alpha$  lies on pseudo-Riemannian lightlike cone  $C(m)$  and the curvatures  $k_1(s)$  and  $k_2(s)$  satisfy

$$\frac{1}{k_1(s)} = c_1 [\cosh(\int k_2(s)ds) \pm \sinh(\int k_2(s)ds)].$$

Conversely if  $\alpha(s)$  is a unit speed spacelike curve in  $\mathbb{E}_1^3$  with spacelike or timelike principal normal  $N$ , the curvatures  $k_1(s) > 0$ ,  $k_2(s) \neq 0$  for each  $s \in I \subset \mathbb{R}$  and one of the statements (i), (ii) and (iii) hold, then  $\alpha$  is a normal curve or congruent to a normal curve.

**Proof.** Let us first suppose that  $\alpha(s)$  is a unit speed spacelike normal curve in  $\mathbb{E}_1^3$  with spacelike or timelike principal normal  $N$ , where  $s$  is pseudo arclength parameter. Then by definition we have

$$\alpha(s) = \lambda(s)N(s) + \mu(s)B(s).$$

Differentiating this with respect to  $s$  and using the corresponding Frenet equations (1), we find

$$\epsilon\lambda k_1 = -1, \quad \lambda' + \mu k_2 = 0, \quad \mu' + \lambda k_2 = 0. \quad (3)$$

From the first and second equation in (3), we get

$$\lambda = -\frac{\epsilon}{k_1}, \quad \mu = \frac{\epsilon}{k_2} \left( \frac{1}{k_1} \right)'. \quad (4)$$

Thus

$$\alpha(s) = -\frac{\epsilon}{k_1}N + \frac{\epsilon}{k_2} \left( \frac{1}{k_1} \right)' B. \quad (5)$$

Further, from the third equation in (3) and using (4), we find the following differential equation

$$\left[ \frac{1}{k_2} \left( \frac{1}{k_1} \right)' \right]' - \frac{k_2}{k_1} = 0. \quad (6)$$

Putting  $y(s) = \frac{1}{k_1}$  and  $p(s) = \frac{1}{k_2}$ , equation (6) can be written as

$$(p(s)y'(s))' - \frac{y(s)}{p(s)} = 0.$$

If we change variables in the above equation as  $t = \int \frac{1}{p(s)} ds$ , then we get

$$\frac{d^2 y}{dt^2} - y = 0.$$

The solution of the previous differential equation is

$$y = c_1 \cosh(t) + c_2 \sinh(t),$$

where  $c_1, c_2 \in \mathbb{R}$ . Therefore,

$$\frac{1}{k_1(s)} = c_1 \cosh\left(\int k_2(s) ds\right) + c_2 \sinh\left(\int k_2(s) ds\right). \quad (7)$$

Thus we have proved statement (i). Next, substituting (7) into (4) and (5), we get

$$\begin{aligned} \lambda &= -\epsilon[c_1 \cosh\left(\int k_2(s) ds\right) + c_2 \sinh\left(\int k_2(s) ds\right)], \\ \mu &= \epsilon[c_1 \sinh\left(\int k_2(s) ds\right) + c_2 \cosh\left(\int k_2(s) ds\right)], \end{aligned}$$

and

$$\begin{aligned} \alpha &= -\epsilon(c_1 \cosh\left(\int k_2(s) ds\right) + c_2 \sinh\left(\int k_2(s) ds\right))N \\ &\quad + \epsilon(c_1 \sinh\left(\int k_2(s) ds\right) + c_2 \cosh\left(\int k_2(s) ds\right))B. \end{aligned} \quad (8)$$

Therefore, from (8) we easily find that

$$g(\alpha, \alpha) = \epsilon(c_1^2 - c_2^2), \quad (9)$$

$$g(\alpha, N) = a_1 \cosh\left(\int k_2(s) ds\right) + a_2 \sinh\left(\int k_2(s) ds\right), \quad (10)$$

$$g(\alpha, B) = a_1 \sinh\left(\int k_2(s) ds\right) + a_2 \cosh\left(\int k_2(s) ds\right), \quad (11)$$

where  $a_1 = -c_1 \in \mathbb{R}$ ,  $a_2 = -c_2 \in \mathbb{R}$ . Consequently, we have proved (ii).

Next, suppose that  $\alpha$  is a normal curve with a null (lightlike) position vector. Then we have  $g(\alpha, \alpha) = 0$ . Substituting this into equation (9), we obtain  $c_1^2 = c_2^2$ . Then (7) becomes

$$\frac{1}{k_1(s)} = c_1[\cosh(\int k_2(s)ds) \pm \sinh(\int k_2(s)ds)]. \quad (12)$$

On the other hand, let us consider the vector

$$m = \alpha(s) + \frac{\epsilon}{k_1}N - \frac{\epsilon}{k_2} \left(\frac{1}{k_1}\right)' B.$$

Differentiating this with respect to  $s$  and using corresponding Frenet equations (1), we find  $m' = 0$ , and therefore  $m = \text{constant}$ . Then  $g(\alpha - m, \alpha - m) = 0$ , which means that  $\alpha$  lies on  $C(m)$ . Consequently, we have proved statement (iii).

Conversely, suppose that statement (i) holds. Then we have

$$\frac{1}{k_1(s)} = c_1 \cosh(\int k_2(s)ds) + c_2 \sinh(\int k_2(s)ds).$$

Differentiating this with respect to  $s$ , we get

$$\left[ \frac{1}{k_2} \left(\frac{1}{k_1}\right)' \right]' = \frac{k_2}{k_1}.$$

By applying Frenet equations (1), we obtain

$$\frac{d}{ds} \left[ \alpha(s) + \frac{\epsilon_1}{k_1}N - \frac{\epsilon_1}{k_2} \left(\frac{1}{k_1}\right)' B \right] = 0.$$

Consequently,  $\alpha$  is congruent to a normal curve. Next, assume that statement (ii) holds. Then the equations (9) and (10) are satisfied. Differentiating (9) with respect to  $s$  and using (10), we find  $g(\alpha, T) = 0$ , which means that  $\alpha$  is normal curve. Finally, assume that statement (iii) holds. Then  $\alpha$  lies on light cone  $C(m)$  with vertex at  $m$ ,  $m = \text{constant}$  and curvatures  $k_1(s)$  and  $k_2(s)$  satisfy the equation (12). Hence we have

$$g(\alpha - m, \alpha - m) = 0.$$

Differentiating this four times with respect to  $s$  and using Frenet equations (1), we get

$$\alpha(s) - m = -\frac{\epsilon}{k_1}N + \left(\frac{\epsilon}{k_2}\right)\left(\frac{1}{k_1}\right)'B.$$

This means that, up to a translation for vector  $m$ , curve  $\alpha$  is congruent to a normal curve. Let us put  $m = 0$ . Then using (12) we easily find  $g(\alpha, \alpha) = 0$ , which proves the theorem.  $\square$

**Theorem 3.2** *Let  $\alpha = \alpha(s)$  be unit speed spacelike normal curve in  $\mathbb{E}_1^3$  with curvatures  $k_1(s) > 0$ ,  $k_2(s) \neq 0$ , non-null principal normal  $N$  and non-null position vector. Then:*

- (i) The position vector  $\alpha$  is spacelike if and only if the curve  $\alpha$  lies on the pseudo-Riemannian sphere  $\mathbb{S}_1^2(m, r)$  and there holds

$$\frac{1}{k_1(s)} = \pm\sqrt{c^2 + \epsilon r^2} \cosh\left(\int k_2(s)ds\right) + c \sinh\left(\int k_2(s)ds\right), \quad c \in \mathbb{R}, \quad \epsilon = \pm 1; \quad (13)$$

- (ii) The position vector  $\alpha$  is timelike if and only if the curve  $\alpha$  lies on the pseudohyperbolic space  $\mathbb{H}_0^2(m, r)$  and there holds

$$\frac{1}{k_1(s)} = \pm\sqrt{c^2 - \epsilon r^2} \cosh\left(\int k_2(s)ds\right) + c \sinh\left(\int k_2(s)ds\right), \quad c \in \mathbb{R}, \quad \epsilon = \pm 1. \quad (14)$$

**Proof.** Let us first assume that the position vector  $\alpha$  is spacelike. Then  $g(\alpha, \alpha) = r^2$ ,  $r \in \mathbb{R}^+$ . Substituting this into (9), we get  $c_1 = \pm\sqrt{c_2^2 + \epsilon r^2}$ . By using the last equation and (7), we obtain that (13) holds. Next, let us consider the vector

$$m = \alpha + (\epsilon/k_1)N - (\epsilon/k_2)(1/k_1)'B.$$

Differentiating this and using the corresponding Frenet equations, we get  $m' = 0$ . Consequently,  $m = \text{constant}$ . It follows that  $g(\alpha - m, \alpha - m) = r^2$ , which means that  $\alpha$  lies on pseudo-Riemannian sphere  $S_1^2(m, r)$  with center  $m$  and of radius  $r$ . Conversely, assume that (13) holds and that  $\alpha$  lies on  $\mathbb{S}_1^2(m, r)$ . Then  $g(\alpha - m, \alpha - m) = r^2$ , where  $r \in \mathbb{R}^+$ . Differentiating this four times with respect to  $s$  and using Frenet equations, we find

$$\alpha - m = -(\epsilon/k_1)N + (\epsilon/k_2)(1/k_1)'B.$$



Therefore, up to a translation for a vector  $m$ ,  $\alpha$  is congruent to a normal curve. In particular, let us put  $m = 0$ . Then (13) implies that  $g(\alpha, \alpha) = r^2$ , which proves statement (i).

The proof of statement (ii) is analogous to the proof of statement (i).  $\square$

**Remark.** The spacelike curves with a null principal normal  $N$ , in the space  $\mathbb{E}_1^3$  can have the first curvature  $k_1 = 0$  or  $k_1 = 1$  [7]. If  $k_1 = 0$ , then  $\alpha(s)$  is straight line. Therefore  $\alpha(s)$  is in direction of  $T(s)$  for each  $s$ . For straight line we have  $N = B = 0$ , so we do not have normal plane  $\{N, B\}$ . Therefore, if  $k_1 = 0$  then  $\alpha(s)$  can not be normal curve.

**Theorem 3.3** *Let  $\alpha(s)$  be unit speed spacelike normal curve in  $\mathbb{E}_1^3$  with a null principal normal  $N$  and  $k_1 = 1$ . Then  $\alpha$  is normal curve if and only if the principal normal and binormal component of the position vector are, respectively,  $g(\alpha, N) = -1$ ,  $g(\alpha, B) = c$ ,  $c \in \mathbb{R}$ .*

**Proof.** Let us first assume that  $\alpha(s)$  is normal curve. Then we have

$$\alpha(s) = \lambda(s)N(s) + \mu(s)B(s). \quad (15)$$

Differentiating this with respect to  $s$  and using Frenet equations (2), we get

$$\mu = -1, \quad \lambda' + \lambda k_2 = 0 \quad \text{and} \quad \mu' - \mu k_2 = 0 \quad (16)$$

We obtain from the third equation in (16) that  $k_2 = 0$ . Then the second equation in (16) implies  $\lambda' = 0$ . Thus  $\lambda = c$ ,  $c \in \mathbb{R}$  and therefore

$$\alpha = cN - B. \quad (17)$$

Finally, we obtain  $g(\alpha, N) = -1$ ,  $g(\alpha, B) = c$ .

Conversely, let  $g(\alpha, N) = -1$ ,  $g(\alpha, B) = c$ . Then differentiating with respect to  $s$ , we find  $k_2 = 0$  and  $g(\alpha, T) = 0$ , which means that  $\alpha$  is normal curve.  $\square$

**Theorem 3.4** *Let  $\alpha(s)$  be unit speed spacelike normal curve in  $\mathbb{E}_1^3$  with a null principal normal  $N$  and  $k_1 = 1$ . Then  $\alpha$  lies on pseudo-Riemannian sphere  $\mathbb{S}_1^2(m, r)$  if and only if  $\alpha$  is plane normal curve with the equation  $\alpha - m = -\frac{r^2}{2}N - B$ .*

**Proof.** Suppose that  $\alpha$  lies on pseudo-Riemannian sphere  $\mathbb{S}_1^2(m, r)$ . Then we have

$$g(\alpha - m, \alpha - m) = r^2, \quad r \in \mathbb{R}^+.$$

Differentiating this and applying Frenet formulae, we find

$$k_2 g(N, \alpha - m) = 0.$$

Thus  $k_2 = 0$ , and  $\alpha$  is plane curve. We will prove that it is normal curve. Decompose the vector  $\alpha - m$  by

$$\alpha - m = aT + bN + cB,$$

where  $a = a(s), b = b(s), c = c(s)$  are arbitrary functions of  $s$ .

Then  $g(\alpha - m, T) = 0 = a$ ,  $g(\alpha - m, N) = c = -1$ ,  $g(\alpha - m, B) = b$ . Differentiating  $g(\alpha - m, B) = b$ , we get  $b = b_0 = \text{constant}$ . We obtain that

$$\alpha - m = b_0 N - B,$$

and since  $g(\alpha - m, \alpha - m) = r^2$ , we have  $g(\alpha - m, \alpha - m) = -2b_0 = r^2$  and  $b_0 = -\frac{r^2}{2}$ .

Finally,  $\alpha$  has the equation

$$\alpha - m = -\frac{r^2}{2}N - B,$$

and it is congruent to a normal curve.

Conversely, if  $\alpha$  is plane normal curve with the equation  $\alpha - m = -\frac{r^2}{2}N - B$  where  $r \in \mathbb{R}^+$  and  $m = (m_1, m_2, m_3) \in \mathbb{E}_1^3$ , then we have  $k_2 = 0$ . Next, we get that  $m = \alpha + \frac{r^2}{2}N + B$  which differentiating in  $s$  gives  $m' = 0$ . Thus  $m = \text{constant} \in \mathbb{E}_1^3$ , (i.e.  $m$  is constant vector). Therefore,  $\alpha$  lies on  $\mathbb{S}_1^2(m, r)$ .  $\square$

**Theorem 3.5** *Let  $\alpha(s)$  be unit speed spacelike normal curve in  $\mathbb{E}_1^3$  with a null principal normal  $N$  and  $k_1 = 1$ . Then  $\alpha$  lies on pseudo-Riemannian hyperbolic space  $\mathbb{H}_0^2(m, r)$  if and only if  $\alpha$  is plane normal curve with the equation  $\alpha - m = \frac{r^2}{2}N - B$ , where  $r \in \mathbb{R}^+$*

**Proof.** The proof is similar with the proof of theorem 3.4.  $\square$

**Theorem 3.6** *Let  $\alpha(s)$  be unit speed spacelike normal curve in  $\mathbb{E}_1^3$  with a null principal normal  $N$  and  $k_1 = 1$ . Then  $\alpha$  lies on light cone  $C(m)$  with vertex at  $m$  if and only if  $\alpha$  is congruent to a normal curve with the equation  $\alpha(s) = -B(s)$ .*

**Proof.** Suppose that  $\alpha$  lies on light cone  $C(m)$  with vertex at point  $m \in \mathbb{E}_1^3$ . Then

$$g(\alpha - m, \alpha - m) = 0.$$

Differentiating the previous equation and using Frenet equations (2), we get  $g(\alpha - m, T) = 0$ ,  $g(\alpha - m, N) = -1$  and  $k_2 = 0$ . Next, decompose the vector  $\alpha - m$  by

$$\alpha - m = aT + bN + cB,$$

where  $a = a(s), b = b(s), c = c(s)$  are arbitrary functions of  $s$ .

Then  $g(\alpha - m, T) = 0 = a$ ,  $g(\alpha - m, N) = c = -1$ ,  $g(\alpha - m, B) = b$ . Differentiating  $g(\alpha - m, B) = b$ , we get  $b = b_0 = \text{constant}$ . It follows that

$$\alpha - m = b_0N - B.$$

Since  $g(\alpha - m, \alpha - m) = 0 = -2b_0$ , we get  $b_0 = 0$ . Thus  $\alpha - m = -B$ . Therefore, up to a translation for the vector  $m$ ,  $\alpha$  is congruent to a normal curve and  $\alpha = -B$ .

Conversely, assume that  $\alpha$  is congruent to a normal curve with the equation  $\alpha = -B$ . Differentiating this we get  $k_2 = 0$ . Let us consider the vector  $m = \alpha + B$ . Taking the derivative of the last equation, we find  $m = \text{constant}$  and finally  $g(\alpha - m, \alpha - m) = 0$ , which means that  $\alpha$  lies on the light cone  $C(m)$ .  $\square$

### Acknowledgements

I would like to express my hearty thanks to Dr. Emilija Nešović (from Kragujevac University - Serbia and Montenegro) for her encouragement during the preparation of this paper. Also I want to thank the referee for his valuable suggestions.

**References**

- [1] Bonnor, W. B.: Null Curves in a Minkowski space-time, Tensor, 20, 1969, 229-242.
- [2] Camcı, Ç., İlarıslan, K. and Šučurović, E.: On pseudohyperbolic curves in Minkowski space-time, Turkish Journal of Math., 27, 315-328, 2003.
- [3] Chen, B. Y.: When does the position vector of a space curve always lie in its rectifying plane?, Amer. Math. Monthly, 110, 2003, 147-152.
- [4] Ekmekci N. and İlarıslan, K.: Higher Curvatures of a Regular Curve in Lorentzian Space, Journal of Institute of Math. and Comp. Sci., Vol. 11, No. 2, 1998, 97-102.
- [5] İlarıslan, K., Nešović, E. and Petrović-Torgašev, M.: Some Characterizations of Rectifying Curves in the Minkowski 3-space, Novi Sad J. Math. Vol 33, No. 2, 2003, 23-32.
- [6] O'Neill, B.: Semi - Riemannian geometry, with applications to relativity. Academic Press. New York, 1983.
- [7] Petrović-Torgašev, M. and Šučurović, E.: Some characterizations of Lorentzian spherical spacelike curves with the timelike and null principal normal, Mathematica Moravica, 4, 2000, 83-92.
- [8] Petrović-Torgašev, M. and Šučurović, E.: Some characterizations of curves lying on the pseudohyperbolic space  $\mathbb{H}_0^2$  in the Minkowski space  $\mathbb{E}_1^3$ , Kragujevac J. Math., 22, 2000, 71-82.
- [9] Walrave, J.: Curves and Surfaces in Minkowski Space, Ph.D. thesis, K. U. Leuven, Fac. of Science, Leuven, 1995.
- [10] Wong, Y. C.: A global formulation of the condition for a curve to lie in a sphere, Monatshefte fur Mathematik, 67, 1963, 363-365.
- [10] Wong, Y. C.: On an explicit characterization of spherical curves, Proceedings of the American Math. Soc., 34, 1972, 239-242.

Kazım İLARSLAN  
 Department of Mathematics,  
 Faculty of Sciences and Arts,  
 Kocatepe University,  
 A.N. Sezer Campus, Afyon-TURKEY  
 e-mail: kilarslan@aku.edu.tr

Received 01.10.2003