

1-1-2004

A Generalization of a Result on Torsion-Free Groups With all Subgroups Subnormal

TAHİRE ÖZEN

Follow this and additional works at: <https://journals.tubitak.gov.tr/math>



Part of the [Mathematics Commons](#)

Recommended Citation

ÖZEN, TAHİRE (2004) "A Generalization of a Result on Torsion-Free Groups With all Subgroups Subnormal," *Turkish Journal of Mathematics*: Vol. 28: No. 4, Article 5. Available at: <https://journals.tubitak.gov.tr/math/vol28/iss4/5>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Mathematics by an authorized editor of TÜBİTAK Academic Journals. For more information, please contact academic.publications@tubitak.gov.tr.

A Generalization of a Result on Torsion-Free Groups With all Subgroups Subnormal

Tahire Özen

Abstract

The main result in this paper is the following: Let G be a torsion-free locally nilpotent group and let F be a finitely generated subgroup of G . If every subgroup of G containing F is subnormal in G , then G is nilpotent.

1. Introduction

In [1] and [5] it is proved that a torsion-free group with all subgroups subnormal is nilpotent. A generalization of this result is given by Smith in [8].

In this note we consider locally nilpotent torsion-free groups such that every subgroup containing a fixed finitely generated subgroup is subnormal and prove the following result :

Theorem. *Let G be a torsion-free locally nilpotent group and let F be a finitely generated subgroup of G . If every subgroup of G containing F is subnormal in G , then G is nilpotent.*

To prove the theorem we mainly exploit Möhres', Casolo's and Detomi's ideas in [4], [1] and [3], respectively.

2. Proof of the Theorem

Let G be a group and H be a subgroup of G . The isolator of H in G is the set defined by $I_G(H) = \{g \in G : g^n \in H, \text{ for some non-negative natural number } n\}$. See [4] for some interesting properties of isolators.

The following Lemma is analogous to Lemma 4 of [4].

Lemma 1. *Let G be a non-nilpotent, locally nilpotent, torsion-free group and let F be a finitely generated subgroup of G . If every subgroup of G containing F is subnormal in G , then there exist a non-negative natural number r , a non-nilpotent subgroup K of G and a finitely generated subgroup H of K containing F such that the subnormality index in K of every subgroup of K containing H is at most r . If G is countable, then K can be chosen such that $I_G(K) = G$.*

Proof. Since G is a non-nilpotent group, G has a non-nilpotent countable subgroup containing F . So we may assume that G is countable. Suppose that the assertion is false. Put $H_0 = F$ and let $x_0 \in G \setminus F$. By Lemma 2 of [4], there exists a subgroup K_1 of G containing F such that $x_0 \notin K_1$ and $I_G(K_1) = G$. Since G is a non-nilpotent group, by Lemma 1.3.5 of [3], K_1 is a non-nilpotent subgroup of G . Hence by Lemma 3 of [4] there exists a finitely generated subgroup H_1 of G such that $H_0 \leq H_1 \leq K_1$ and $s(K_1 : H_1) > 1$, where $s(K_1 : H_1)$ is the subnormality index of H_1 in K_1 . This implies that there exists an element $x_1 \in [K_1, H_1] \setminus H_1$ and clearly $x_0, x_1 \notin H_1$. Assume that there exist a non-nilpotent subgroup K_{i-1} of G and a finitely generated subgroup H_{i-1} of K_{i-1} such that $x_0, \dots, x_{i-1} \notin H_{i-1}$ for a natural number $i \geq 1$. Then we can obtain a non-nilpotent subgroup K_i of K_{i-1} containing a finitely generated subgroup H_i such that $s(K_i : H_i) > i$ and $H_{i-1} \leq H_i$ as above. Thus there exists $x_i \in [K_{i,i} H_i] \setminus H_i$ and clearly $x_0, \dots, x_i \notin H_i$. Now put $H = \cup_{i=1}^{\infty} H_i$. Since $F \leq H$, H is subnormal in G by hypothesis. If d is the subnormality index of H in G , then $x_d \in [K_{d,d} H_d] \leq [G, {}_d H] \leq H$. Hence $x_d \in H_i$ for $i > d$, a contradiction. That completes the proof. \square

Lemma 2. *Let G be a locally nilpotent, torsion-free, metabelian group and let F be a finitely generated subgroup of G . If every subgroup of G containing F is subnormal in G , then G is nilpotent.*

Proof. Assume that G is not nilpotent. Hence we may assume that G is countable. By Lemma 1 there exist a natural number r , a non-nilpotent subgroup K of G and a finitely generated subgroup H of K containing F such that every subgroup of K containing H has defect at most r in K .

Let $A = HK'$ and $L = I_K(A')$. Since H is subnormal in A and K' is nilpotent, by Lemma 1 of [7] A is nilpotent and so $I_K(A')$ is a nilpotent group by Lemma 1.3.5 of [3]. So K/L is non-trivial and torsion-free. Since AL/L is an abelian normal subgroup of K/L and every subgroup of K/L containing HL/L has defect in K/L at most r , K/L

is nilpotent by Lemma 3 of [1]. Hence we obtain that K is nilpotent by Lemma 4 of [1]. This contradiction completes the proof. \square

Lemma 3. *Let G be a locally nilpotent, torsion-free group such that G' is nilpotent and let F be a finitely generated subgroup of G . If every subgroup of G containing F is subnormal in G , then G is nilpotent.*

Proof. Since $G/I_G(G'')$ is nilpotent by Lemma 2 and G' is nilpotent, G is nilpotent by Lemma 4 of [1].

Proof of the Theorem. Assume that the assertion is false. Then we may assume that G is countable and by Lemma 1 there exist a non-negative natural number r , a non-nilpotent subgroup K of G and a finitely generated subgroup H of K containing F such that the subnormality index in K of every subgroup of K containing H is at most r . We proceed by induction on r . If $r = 1$, then every subgroup containing H is normal in K . By 5.3.7 (Dedekind-Baer) of [6] $K/I_K(H)$ is abelian. By Lemma 1.3.5 of [3] $I_K(H)$ is nilpotent and Lemma 3 gives the contradiction that K is nilpotent. If $r > 1$ then $N = H^K$ is nilpotent. Because if R is a subgroup of N such that $R \geq H$, then $R^K = N$ and so R has defect at most $r - 1$ in $H^K = N$. Thus by induction hypothesis, N is a nilpotent group. By Corollary to Theorem 1 (Roseblade Theorem) of [2], $K/I_K(N)$ is a nontrivial soluble group. Let the derived length of $K/I_K(N)$ be d . Now we prove that K is nilpotent by induction on d . If $d = 1$ then by Lemma 1 (v) of [4], $K/I_K(N')$ is a metabelian group which satisfies the conditions of Lemma 2. Thus $K/I_K(N')$ is nilpotent. By Lemma 4 of [1] K is nilpotent. If $d > 1$ then since the derived length of $K'I_K(N)/I_K(N)$ is $d - 1$, by induction hypothesis $K'I_K(N)$ is nilpotent. Since $K/I_K((K'I_K(N))')$ is a metabelian group which satisfies the conditions of Lemma 2, it is nilpotent. Thus by Lemma 4 of [1] K is nilpotent. But this is a contradiction. \square

The author is grateful to Professor Carlo Casolo for his helpful suggestions in proof of the Theorem.

References

- [1] C. Casolo. Torsion-free groups with all subgroups subnormal. *Rend. Circ. Mat. Palermo* **2** 50 (2001), 321–324.

ÖZEN

- [2] J.E.Roseblade. On groups in which every subgroup is subnormal. *J. Algebra* **2** (1965), 402-412.
- [3] E. Detomi. Generalizing subnormality and nilpotency in infinite groups. PhD Thesis, Firenze University.
- [4] W. Möhres. Torsionsfreie Gruppen, deren Untergruppen alle subnormal sind. *Math. Ann.* **284** (1989), 245-249.
- [5] H. Smith. Torsion-free groups with all subgroups subnormal. *Arch. Math.(Basel)* **76** (2001),1-6.
- [6] D.J.S. Robinson. A course in the theory of groups.Springer-Verlag New York. Heidelberg, Berlin, (1982).
- [7] W. Möhres. Torsionsgruppen, deren Untergruppen alle subnormal sind. *Geom.Dedicata* **31** (1989), 237-244.
- [8] H. Smith. Torsion-free groups with all non-nilpotent subgroups subnormal. *Topics in infinite groups, Quad. Mat.* **8**, Aracne, Rome (2001), 297-308.

Tahire ÖZEN
Faculty of Arts and Sciences,
Department of Mathematics,
06500, Ankara-TURKEY
e-mail: tahire@gazi.edu.tr

Received 27.06.2003