

1-1-2005

## Some Reproductive Parameters and Biochemical Properties in Akkaraman and Awassi Rams

MUSTAFA GÜNDOĞAN

MUSTAFA SERTESER

Follow this and additional works at: <https://journals.tubitak.gov.tr/veterinary>



Part of the [Animal Sciences Commons](#), and the [Veterinary Medicine Commons](#)

---

### Recommended Citation

GÜNDOĞAN, MUSTAFA and SERTESER, MUSTAFA (2005) "Some Reproductive Parameters and Biochemical Properties in Akkaraman and Awassi Rams," *Turkish Journal of Veterinary & Animal Sciences*: Vol. 29: No. 3, Article 4. Available at: <https://journals.tubitak.gov.tr/veterinary/vol29/iss3/4>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Veterinary & Animal Sciences by an authorized editor of TÜBİTAK Academic Journals. For more information, please contact [academic.publications@tubitak.gov.tr](mailto:academic.publications@tubitak.gov.tr).

## Some Reproductive Parameters and Biochemical Properties in Akkaraman and Awassi Rams

Mustafa GÜNDOĞAN

Department of Reproduction and Artificial Insemination, Faculty of Veterinary Medicine, Afyon Kocatepe University,  
ANS Campus, 03200, Afyonkarahisar - TURKEY  
E-mail: mgundogan@aku.edu.tr

Mustafa SERTESER

Department of Biochemistry, Faculty of Medicine, Afyon Kocatepe University, Afyonkarahisar - TURKEY

Received: 13.11.2002

**Abstract:** The objective of the present study was to describe some reproductive parameters, hormonal levels and some biochemical properties of blood serum in rams under the Afyonkarahisar province conditions. Semen (via an artificial vagina) and blood were collected five times during the breeding season from a total of 10 mature and healthy fat-tailed rams (Akkaraman and Awassi) in Afyonkarahisar. Breed effects were significant for relative testes volume, sperm concentration, percentage of abnormal spermatozoa, triiodothyronine ( $T_3$ ), albumin/globulin ratio, total lipid ( $P < 0.05$ ) and cholesterol ( $P < 0.01$ ). Some reproductive parameters and biochemical properties of Akkaraman and İvesi rams indicated that the values were recorded in the optimum range during the breeding season under Afyonkarahisar conditions. In addition, some reproductive parameters of rams show a correlation with some biochemical and enzymatic properties.

**Key Words:** Biochemical properties, hormone, ram, semen.

### İvesi ve Akkaraman Koçlarda Bazı Reprodüktif Parametreler ve Biyokimyasal Özellikler

**Özet:** Afyonkarahisar koşullarında yetiştirilen koçlarda bazı reprodüktif parametreler ve hormon miktarları ile kan serumu biyokimyasal özelliklerin araştırılması amacıyla yapılan bu çalışmada, aşıım sezonu esnasında, yağlı kuyruklu ve sağlıklı toplam 10 koçtan (Akkaraman ve İvesi) beşer numune kan ve sun'i vagen yardımıyla da sperma toplandı. Relatif testis hacmi, spermatozoa yoğunluğu, anormal spermatozoa oranı, triiodothyronine ( $T_3$ ) miktarı, albumin/globulin oranı, toplam lipid miktarı ( $P < 0.05$ ) ve kolesterol miktarı ( $P < 0,01$ ) yönünden koçlar arasındaki farklılıkların önemli olduğu gözlemlendi. Afyonkarahisar koşullarında aşıım sezonu esnasındaki İvesi ve Akkaraman koçların bazı reprodüktif değerlerinin ve bazı biyokimyasal özelliklerinin optimal sınırlar içerisinde olduğu ve ayrıca koçların bazı reprodüktif parametrelerinin bazı biyokimyasal ve enzimatik özellikler ile ilişkileri olduğu gözlemlendi.

**Anahtar Sözcükler:** Biyokimyasal özellik, hormon, koç, sperma.

### Introduction

Most sheep show a seasonal pattern in reproductive activity that is shaped by seasonal changes in their habitats. Biochemical estimates of blood serum are used for semen evaluation, since using semen characteristics alone is not completely satisfactory for semen appraisal in the current practice of commercial artificial insemination (1-4). However, the biochemical evaluation of ram blood and its relationship with semen characteristics are completely unknown. With better knowledge of ram reproductive physiology a more accurate andrological evaluation could be conducted, which would improve

reproductive efficiency and enhance breeding schemes and the rate of genetic gain. The present study is concerned with investigating some major biochemical constituents of blood serum and their relationship with hormonal levels and some reproductive parameters in Akkaraman and Awassi rams during the breeding season under Afyonkarahisar province conditions.

### Materials and Methods

This work was carried out in the breeding season (September-October 2001) at the Research and

Manipulation Livestock of the Faculty of Veterinary Medicine, University of Afyon Kocatepe located in Afyonkarahisar province (L: 1021 m, 38° 45' N, 30° 32' W), Turkey. Semen evaluation was carried out at the Faculty of Veterinary Medicine, Afyon Kocatepe University. Hormonal and biochemical analyses were carried out at the Medicine School of Afyon Kocatepe University.

### Animals and management

In a factorial randomized complete-block design experiment, 10 sexually mature and healthy rams were used. Each of five Akkaraman and Awassi were used to study the relationship between reproductive parameters and biochemical constituents of blood serum in rams groups during the breeding season. All animals were aged 3-4 years. Animals were offered rough and concentrate supplement according to their body weight requirements (5).

In terms of sexual performance, scrotal sac volume was measured by the water displacement method described by Demirci (6) and was then divided by body weight to calculate the relative testes volume. Sexual behavior of rams was recorded by using the reaction time criterion. The time elapsed between introducing the ram to a female ewe at estrous and semen collection was measured using a stopwatch.

### Semen evaluation

Total 5 ejaculates were collected weekly by using an artificial vagina. Ejaculates were placed in a water bath at 38 °C. Progressive motility (%) was subjectively estimated at 400 X magnification using a light microscope. Sperm concentration was measured by a haemocytometer slide. Percentages of abnormal spermatozoa were assessed by fluid fixation method according to Demirci (6).

### Blood Collection

Total 5 blood samples were collected weekly from each animal via the jugular vein. Serum was separated by centrifugation at 3000 r.p.m. for 15 min. and it was collected and stored at -20 °C until analysis.

### Hormonal assay and determination of blood serum constituents

The concentration of triiodothyronine ( $T_3$ ) and testosterone in the blood serum was measured by using commercial Chemiluminescent kits (ADVIA Centaur

Competitive Immunoassay System, Bayer Diagnostic Corporation, U.S.) according to Fernandez-Ullao and Maxon (7) and Spratt et al. (8).

Total blood serum protein was measured by the Biuret method and total albumin (A) concentration was determined by the method of Doumas et al. (9). Total globulin (G) concentration was calculated as the difference between blood serum total protein and blood serum albumin, then A/G ratio was calculated. Total lipids and cholesterol concentrations were measured by a colorimetric method (10). Aspartate aminotransferase (AST) and alanine aminotransferase (ALT) activities were measured in a Hitachi 917 Clinical Chemistry Analyzer by using commercially available diagnostic kits supplied from Roche Diagnostics GmbH (D-68298, Mannheim, Germany). Both enzyme activities were determined photometrically in which the decrease in NADH levels were directly proportional to enzyme activities and AST/ALT ratios were calculated (11).

### Statistical analysis

The groups were compared by Student's t-test and the partial correlation coefficients and probabilities among different parameters were worked out (12).

### Results

Overall mean values of some reproductive parameters and biochemical and enzymatic properties of blood serum in rams during the breeding season are presented in the Table. The partial correlation coefficients and probabilities among different parameters were worked out. Relative testes volume from reproductive parameters were negatively correlated with sperm concentration ( $P < 0.01$ ) and sperm motility ( $P < 0.05$ ) in all rams. Reaction times were positively correlated with testosterone levels in Akkaraman ( $P < 0.01$ ) and Awassi ( $P < 0.05$ ) rams. Percentages of abnormal spermatozoa were negatively ( $P < 0.01$ ) correlated with testosterone levels and sperm motility but positively correlated ( $P < 0.01$ ) with sperm concentration in all rams. Total protein from biochemical properties of blood serum were positively ( $P < 0.01$ ) correlated with albumin, globulin and ALT activity but negatively ( $P < 0.01$ ) correlated with total lipid, cholesterol and AST activity. Total lipids were positively ( $P < 0.01$ ) correlated with cholesterol and ALT activity but negatively ( $P < 0.01$ ) correlated with AST activity and AST/ALT ratio. AST activity was negatively ( $P$

< 0.01) correlated with cholesterol and ALT activity and positively ( $P < 0.01$ ) correlated with AST/ALT ratio in all rams.

Total protein and globulin were positively ( $P < 0.05$ ) correlated with sperm motility, sperm concentration and testosterone level in all rams. Total lipid levels were negatively ( $P < 0.01$ ) correlated with sperm motility and concentration and positively ( $P < 0.05$ ) correlated with percentage of abnormal spermatozoa in all rams. In addition, total lipids and cholesterol were positively ( $P < 0.01$ ) correlated with  $T_3$  levels and negatively ( $P < 0.05$ ) correlated with testosterone levels in all rams. Percentage of abnormal spermatozoa was negatively ( $P < 0.01$ ) correlated with alanine amino transaminase (ALT) activity but were positively ( $P < 0.05$ ) associated with aspartate amino transaminase (AST) activity and AST/ALT ratio in all rams.

Table. Overall mean values of some reproductive parameters, some biochemical and enzymatic properties of blood serum in rams during the breeding season in Afyon province conditions ( $X \pm$  s.e.m.) (n:25).

	Akkaraman	Awassi
Rel. testes volume (ml/kg)	10.6 $\pm$ 0.34*	10.9 $\pm$ 0.32*
Reaction time (s)	6.8 $\pm$ 0.35	6.6 $\pm$ 0.40
Motility (%)	77.8 $\pm$ 0.73	75.4 $\pm$ 1.00
Sperm concentration ( $\times 10^9$ /ml)	4.7 $\pm$ 0.08*	5.0 $\pm$ 0.07*
Abnormal sperm (%)	3.9 $\pm$ 0.12*	3.5 $\pm$ 0.10*
Testosterone (ng/ml)	4.7 $\pm$ 0.46	4.8 $\pm$ 0.64
$T_3$ ( $\mu$ g/100ml)	1.22 $\pm$ 0.10*	0.87 $\pm$ 0.07*
Total protein (g/100ml)	7.2 $\pm$ 0.23	7.5 $\pm$ 0.08
Albumin (A) (g/100ml)	3.3 $\pm$ 0.05	3.3 $\pm$ 0.08
Globulin (G) (g/100ml)	3.9 $\pm$ 0.20	4.3 $\pm$ 0.19
A/G	0.85 $\pm$ 0.04*	0.77 $\pm$ 0.02*
Total lipid (mg/100ml)	69.7 $\pm$ 2.24*	73.7 $\pm$ 2.07*
Cholesterol (mg/100ml)	28.9 $\pm$ 1.41**	33.6 $\pm$ 1.24**
AST	116.7 $\pm$ 4.03	115.8 $\pm$ 4.11
ALT	15.7 $\pm$ 0.65	14.6 $\pm$ 1.89
AST/ALT	7.4 $\pm$ 0.62	7.9 $\pm$ 0.32

\* $P < 0.05$

\*\* $P < 0.01$

## Discussion

Relative testes volumes were lower than the findings of Taha et al. (13) in Awassi rams and similar to those in Barki rams. The mean reaction times were lower than the

findings of Taha et al. (13). The differences might be due to breed, age, feeding and season. In addition, because of the farm conditions the rams and sheep were maintained closely therefore the effects of sheep could be concerned. The percentage of motile spermatozoa were higher than the findings of Taha et al. (13) (68.91% and 64.3%) in sexually mature Awassi and Barkies, lower than that of Gündoğan et al. (14) (80.9%) in 16 month old Akkaramans, of Kaya et al. (15) (85.7%) and Piriñçi et al. (16) (96%) and similar to the findings of Aral and Tekin (17), Ataman et al. (18) and Gündoğan (19). The spermatozoa concentrations were lower than the findings of Taha et al. (13) ( $5.2 \times 10^9$ /ml in Awassi), higher than those of Aral and Tekin (17) ( $3.1 \times 10^9$ /ml), Gündoğan (19) ( $3.8 \times 10^9$ /ml) and Kaya et al. (15) ( $3.6 \times 10^9$ /ml) and similar to those of Ataman et al. (18) in 15 month old Merino rams, Piriñçi et al. (16) ( $4.6 \times 10^9$ /ml for 2 year old Akkaramans) and Taha et al. (13) ( $4.7 \times 10^9$ /ml in Barkies). Percentages of abnormal spermatozoa rates were similar to the findings of Aral and Tekin (17) (3.3%), Gündoğan (19) (3.5%) and Kaya et al. (15) (3.6%), lower than those of Pérez et al. (4) (9.4%) and Taha et al. (13) (14.2%) and higher than those of Ataman et al. (18) (1.6%) and Piriñçi et al. (16) (2.7%). Breed effect was significant ( $P < 0.05$ ) for sperm concentration and percentage of abnormal spermatozoa. The differences in the spermatological parameters might be due to breed, age, feeding and management, evaluation technique, semen collection time and season.

Blood serum testosterone levels were similar to the results of Piriñçi et al. (20) and Taha et al. (13) in Awassi (Imported), higher than those of Gündoğan (19) (2.7 ng/ml) and lower than those of Pérez et al. (4) (19.7 nmol/l = 12.85 ng/ml) and Taha et al. (13) (13.41 ng/ml) in Barki rams. Triiodothyronine levels were lower than the findings of Taha et al. (13) in Awassi (Imported) and Barki rams and similar to those in Awassi (locally born) rams. The differences in hormonal levels might be due to breed, age, feeding and management, sample collection time and evaluation procedure.

The biochemical and enzymatic properties of blood serum in rams were in the optimum range and in accordance with the references (21-24). In addition, breed effect was significant for A/G ratio, total lipid ( $P < 0.05$ ) and cholesterol ( $P < 0.01$ ) and relationships between some biochemical and enzymatic properties of rams' blood serum were observed.

Many studies have shown that the low content of seminal plasma proteins is associated with poor seminal quality (1,2,25-28). In the present study, it was found that there were positive relationships between sperm motility, sperm concentration, testosterone level and level of total proteins and globulin ( $P < 0.05$ ) but there were negative relationships between percentage of abnormal spermatozoa and albumin and A/G ratio ( $P < 0.05$ ) in all of rams' blood serum.

Kelso et al. (29) reported that reductions in sperm concentration and motility were associated with a decrease in seminal plasma lipids content and also with sperm aging (poor semen quality). In the present study, total lipid levels were negatively ( $P < 0.01$ ) correlated with sperm motility and concentration and positively ( $P < 0.05$ ) correlated with percentage of abnormal spermatozoa in all rams.

The blood cholesterol level drops before the metabolic rate rises (30,31). In the present study total lipids and cholesterol were positively ( $P < 0.01$ ) correlated with triiodothyronine levels and negatively ( $P < 0.05$ ) correlated with testosterone levels in all rams. The differences between rams may be due to breed and genetics.

The assay of transaminase enzyme activity (AST and ALT) is a good indicator of semen quality because it measures sperm membrane stability (32,33). In the present study, positive relationships were found between percentage of abnormal spermatozoa and AST activity and AST/ALT ratio ( $P < 0.05$ ) and negative relationships between ALT activity ( $P < 0.01$ ) in ram blood serum and negative correlations were observed with sperm concentration, AST activity and AST/ALT ratio ( $P < 0.05$ ) in the two ram breeds.

In conclusion, some of the reproductive parameters and biochemical analyses of rams' blood serum indicated that the values were recorded in the normal range during the breeding season under Afyonkarahisar province conditions. Some reproductive parameters of rams show a correlation with some biochemical and enzymatic properties and could be used to evaluate the reproductive performance of a ram. In addition, total protein, total lipid, cholesterol, AST activity and AST / ALT ratio from biochemical and enzymatic properties of blood serum could be used to evaluate the reproductive performance of an Akkaraman and Awassi ram.

## References

1. Ashworth, P.J., Harrison, R.A., Miller, N.G., Plummer, J.M., Watson, P.F.: Survival of ram spermatozoa at high dilution: protective effect of simple constituents of culture media as compared with blood serum. *Reprod. Fert. Dev.*, 1994; 6: 173-180.
2. Barrios, B., Perez-Pe, R., Gallego, M., Tato, A., Osada, J., Muino-Blanco, T., Cebrian-Perez, J.A.: Blood serum proteins revert the cold-shock damage on ram sperm membrane. *Biol. Reprod.*, 2000; 63: 1531-1537.
3. Lincoln, G.A., Lincoln, C.E., McNeilly, A.S.: Seasonal cycle in the blood plasma concentration of FSH, inhibin and testosterone, and testicular size in rams of wild, and domesticated breeds of sheep. *J. Reprod. Fert.*, 1990; 88: 623-633.
4. Pérez, C.R., López, A., Castrillejo, A., Bielli, A., Laborde, D., Gastel, T., Tagle, R., Queirolo, D., Franco, J., Forsberg, M., Rodriguez-Martinez, H.: Reproductive seasonality of corriedale rams under extensive rearing conditions. *Acta Vet. Scand.*, 1997; 38: 109-117.
5. Church, D.C.: Digestive physiology and nutrition of ruminants. 3th ed., O. and B. Books Inc., Oregon. 1983.
6. Demirci, E.: Evcil hayvanlarda reproduksiyon, sun'i tohumlama ve androloji ders notları. F.Ü. Vet. Fak. Ders Teksiri, No: 53, Elazığ. 2002.
7. Fernandez-Ullao, M., Maxon, H.R. 1989. Thyroid. In: Kaplan, L.A. and Pesce, A.J. Eds. *Clinical chemistry: theory analysis, and correlation*. 2nd ed. St. Louis: CV, Mosby. 1989; 620-638.
8. Spratt, D.I., Bigos, S.T., Beitins, I.: Both hyper and hypogonadotropic hypogonadism occur transiently in acute illness: bio and immunoactive gonadotropins. *J. Clin. Endocrinol. Metab.*, 1992; 7: 1562-1570.
9. Doumas, B.T., Wastson, W.A., Biggs, H.G.: Albumin standards and the measurement of serum albumine with bromocresol green. *Clin. Chim. Acta.*, 1977; 31: 87-96.
10. Kaneko, J.J.: *Clinical biochemistry of domestic animals*. 3rd ed. Academic Press, New York. 1980.
11. Reitman, S., Frankel, S.: *Grand wohl's clinical laboratory methods and diagnosis*. 7th ed. C.V. Mosby Company, St. Louis. 1970.
12. Anonymous: SPSS, 5.0.1. for windows release, standard version. Sterling Technologies, Inc., 444 N., Michigan Avenue, Chicago, IL, 60611. 1992.
13. Taha, T.A., Abdel-Gawad, E.I., Ayoub, M.A.: Monthly variations in some reproductive parameters of barki and awassi rams throughout 1 year under subtropical conditions. 1- Semen characteristics and hormonal levels. *Anim. Sci.*, 2000; 71: 317-324.

14. Gündoğan, M., Demirci, E., Bozkurt, T., Sönmez, M.: The changes before, during and later the breeding season in the semen characteristics of the rams. *Yüzüncü Yıl Üniv. Vet. Fak. Derg.*, 1997; 8: 40-42.
15. Kaya, A., Yıldız, C., Lehimcioğlu, N.C., Ergin, A., Aksoy, M.: Konya merinosu koçlarında sperma kalitesi, testis ölçüleri ve kan testosteron düzeylerine ilişkin mevsimsel değişikliklerin araştırılması. *Hay. Araş. Derg.*, 1999; 9: 1-5.
16. Piriñçi, I., Karahan, I., Bozkurt, T., Güler, O.: Koçlarda prostaglandin F<sub>2</sub>α, furosemid ve indometasinin sperma özellikleri üzerine etkilerinin araştırılması. *Vet. Bil. Derg.*, 2001; 17: 83-88.
17. Aral, F., Tekin, N.: Koçlarda sperma kalitesi üzerine mevsimin etkisi. *Hay. Araş. Derg.*, 1996; 6: 15-20.
18. Ataman, M.B., Kaya, A., Karaca, F., Yıldız, C., Çoyan, K., Ergin, A., Aksoy, M.: Toklularda testisin sezon içi ve sezon dışı morфометrik ölçüleriyle spermatolojik özellikler arasındaki ilişkinin belirlenerek damızlık seçiminde kullanılabilirliğinin araştırılması. *Hay. Araş. Derg.*, 1996; 6: 1-7.
19. Gündoğan, M.: Koçların testis ölçülerinin spermatolojik özellikler ve kan serumu testosteron miktarları ile ilişkisi. *Hay. Araş. Derg.*, 1999; 9: 49-52.
20. Piriñçi, I., Karahan, I., Gürsu, F., Bozkurt, T., Güler, O.: Koçlarda prostaglandin F<sub>2</sub>α, furosemid ve indometasinin serum testosteron düzeyleri üzerine etkilerinin araştırılması. *Vet. Bil. Derg.*, 2001; 17: 5-12.
21. Alonso, A.J., Teresa, R., Garcia, M.J.R., Vallejo, M., De-Teresa, R.: The effects of age and reproductive status on serum and blood parameters in Merino breed sheep. *J. Vet. Med. A.*, 1997; 44: 223-231.
22. Jelinek, P., Illek, J., Helenova, I., Frajs, Z.: Biochemical and hematological values of the blood in rams during rearing. *Acta Vet. Brno.*, 1984; 53: 143-150.
23. Pavlovic, O., Vitic, J.: Proteins and lipoproteins of sheep blood serum. *Acta Vet. Yugoslavia*, 1979; 29: 147-154.
24. Prabhakaran, S.K., Rao, R.G.: Total proteins, albumin, globulin and A/G ratio in Madras red sheep. *Cheiron*, 1994; 23: 62-64.
25. White, I.G., Goh, P., Voglmayr, J.K.: Effect of male reproductive tract fluids and proteins on the metabolism and motility of ram spermatozoa. *Arch. Androl.*, 1987; 19: 115-125.
26. Zedda, M.T., Bini, P.P., Pav, S., Sbernardori, U.: Constituents of blood serum and blood serum of the ram. *Bull. Soc. Ital. Biol. Sper.*, 1996; 72: 227-230.
27. Paz, G., Lewin, L.M.: Chemical composition of seminal fluid as a diagnostic aid in clinical andrology. *Harefuah*, 1992; 123: 328-330.
28. Georgiev, G.S.: Biological and biochemical studies of the sperm and blood serum of rams with degraded sperm production. *Vet. Med. Nauki*, 1983; 20: 99-104.
29. Kelso, K.A., Redpath, A., Noble, R.C., Speake, B.K.: Lipid and antioxidant changes in spermatozoa and blood serum throughout the reproductive period of bulls. *J. Reprod. Fert.*, 1997; 109: 1-6.
30. Marai, İ.F.M., Daader, A.H., Mekki, M.Y., Gabr, H.A., İbrahim, H.: Effect of housing system, season of year and age on some physiological and hematological parameters and blood constituents of Ossimi sheep. *J. Arid. Environ.*, 1992; 22: 277-285.
31. Ramos, J.J., Verde, M.T., Marca, M.C., Fernandez, A.: Clinical chemical values and variations in Rasa Aragonesa /Aragonesa/ ewes and lambs. *Small Rum. Res.*, 1994; 13: 133-139.
32. Corteel, J.M.: Effects of blood serum on the survival and fertility of spermatozoa kept in vitro. *Reprod. Nutr. Dev.*, 1980; 20: 1111-1123.
33. Khokhar, B.S., Mehar, S., Chaudhary, K.C.: Transaminases in cattle and buffalo (*Bubalis bubalis*) semen in relation to fertility and seminal characteristics during moderate and colder seasons. *Anim. Reprod. Sci.*, 1987; 13: 177-182.