

1-1-2004

On θ -Euclidean L-Fuzzy Ideals of Rings

AYTEN KOÇ

EROL BALKANAY

Follow this and additional works at: <https://journals.tubitak.gov.tr/math>



Part of the [Mathematics Commons](#)

Recommended Citation

KOÇ, AYTEN and BALKANAY, EROL (2004) "On θ -Euclidean L-Fuzzy Ideals of Rings," *Turkish Journal of Mathematics*: Vol. 28: No. 2, Article 4. Available at: <https://journals.tubitak.gov.tr/math/vol28/iss2/4>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Mathematics by an authorized editor of TÜBİTAK Academic Journals. For more information, please contact academic.publications@tubitak.gov.tr.

On θ -Euclidean L -Fuzzy Ideals of Rings

Ayten Koç, Erol Balkanay

Abstract

In this paper we define a θ -Euclidean level subset and a θ -Euclidean level ideal. We also give some properties of a θ -Euclidean level subset.

Key Words: Ideal; Fuzzy ideal; Level subset; θ -Euclidean level ideal; θ -Euclidean L -fuzzy ideal.

1. Introduction

In this paper we define a new set denoted by $\mu_{\theta y}$. We call it a θ -Euclidean level subset. Then we will show that for $0 \neq y \in R$, the set $\mu_{\theta y}$ is an ideal of R if $\mu: R \rightarrow L$ is an L -fuzzy ideal of R . We also give a theorem on θ -Euclidean L -fuzzy ideal of R .

2. Preliminaries

Throughout this paper, R denotes a commutative ring with identity. L denotes a lattice with the least element 0 and the greatest element 1. Unless stated otherwise L is complete and completely distributive in the sense that it satisfies the following law:

$$\bigvee \{a_i \mid i \in I\} \wedge \bigvee \{b_j \mid j \in J\} = \bigvee \{a_i \wedge b_j \mid i \in I, j \in J\} \quad [3]$$

for all $a_i, b_j \in L$.

Definition 2.1 [3]. *An L -fuzzy ideal is a function $J: R \rightarrow L$ satisfying the following axioms:*

- (i) $J(x + y) \geq J(x) \wedge J(y)$,
- (ii) $J(-x) = J(x)$,
- (iii) $J(xy) \geq J(x) \vee J(y)$.

Since we are considering L -fuzzy ideals over a fixed lattice L , we shall call them fuzzy ideals only.

Definition 2.2 [5]. Let μ be any fuzzy subset of a set S and let $t \in [0, 1]$. The set $\mu_t = \{x \in S \mid \mu(x) \geq t\}$ is called a level subset of μ .

Proposition 2.3 [2, 3].

- (i) A function $J: R \rightarrow L$ is a fuzzy ideal iff $J(x - y) \geq J(x) \wedge J(y)$ and $J(xy) \geq J(x) \vee J(y)$.
- (ii) If $J: R \rightarrow L$ is a fuzzy ideal, then
 - (a) $J(0) \geq J(x) \geq J(1)$, for all $x \in R$;
 - (b) $J(x - y) = J(0)$ implies $J(x) = J(y)$ for all $x, y \in R$;
 - (c) the level cuts $J_\alpha = \{x \in R \mid J(x) \geq \alpha\}$ are ideals of R , for $\alpha \leq J(0)$. Conversely, if each J_α is an ideal, then J is a fuzzy ideal.

3. θ -Euclidean L -fuzzy ideal

Definition 3.1 [1]. Let $\theta: R \rightarrow L$ be a non-constant fuzzy subset of R . A function $\mu: R \rightarrow L$ is called a θ -Euclidean L -fuzzy ideal if μ satisfies the following axioms:

- (i) $\mu(x + y) \geq \mu(x) \wedge \mu(y)$ for all x, y in R ,
- (ii) $\mu(-x) = \mu(x)$;
- (iii) $\mu(xy) \geq \mu(x) \vee \mu(y)$;
- (iv) For any $x, y \in R$, with $y \neq 0$, there exist elements $q, r \in R$ such that $x = yq + r$ where either $r = 0$ or else $\mu(r) \vee \theta(r) \geq \mu(y) \vee \theta(y)$.

Now we will define a new set called a θ -Euclidean Level subset and examine this set.

Definition 3.2 Let $\mu: R \rightarrow L$ and $\theta: R \rightarrow L$ be fuzzy sets. For $0 \neq y \in R$, the set $\mu_{\theta_y} = \{x \in R \mid \text{there exist elements } q, r \in R \text{ such that } x = yq + r \text{ where either } r = 0 \text{ or else } \mu(r) \geq \mu(y) \vee \theta(y)\}$ is called a θ -Euclidean level subset of μ .

Theorem 3.3 *Let $\mu: R \rightarrow L$ be an L -fuzzy ideal of R . Then for $0 \neq y \in R$, μ_{θ_y} is an ideal of R . Also $\mu_{\theta_y} \neq \{0\}$.*

Proof. Let x_1 and x_2 be elements of μ_{θ_y} . We have to show that $x_1 + x_2 \in \mu_{\theta_y}$.

Then there exist elements $q_1, r_1 \in R$ such that $x_1 = yq_1 + r_1$ where either $r_1 = 0$ or else $\mu(r_1) \geq \mu(y) \vee \theta(y)$. And then there exist elements $q_2, r_2 \in R$ such that $x_2 = yq_2 + r_2$ where either $r_2 = 0$ or else $\mu(r_2) \geq \mu(y) \vee \theta(y)$. Hence $x_1 + x_2 = y(q_1 + q_2) + r_1 + r_2$ and $q_1 + q_2, r_1 + r_2 \in R$. Also we know that $\mu(r_1 + r_2) \geq \mu(r_1) \wedge \mu(r_2)$ since μ is an L -fuzzy ideal of R . If $r_i \neq 0$ for $i = 1$ or $i = 2$, then $\mu(r_i) \geq \mu(y) \vee \theta(y)$. Therefore we can say that there exist elements $q_1 + q_2, r_1 + r_2 \in R$ such that $x_1 + x_2 = y(q_1 + q_2) + r_1 + r_2$ where either $r_1 + r_2 = 0$ or else $\mu(r_1 + r_2) \geq \mu(y) \vee \theta(y)$. This means that $x_1 + x_2$ is an element of μ_{θ_y} .

Now, we need to show that $ax \in \mu_{\theta_y}$ for all $x \in \mu_{\theta_y}$ and $a \in R$. Let $a \in R$ and $x \in \mu_{\theta_y}$. Then there exist elements $q, r \in R$ such that $x = yq + r$ where either $r = 0$ or else $\mu(r) \geq \mu(y) \vee \theta(y)$. Suppose that $r \neq 0$. Then $\mu(r) \geq \mu(y) \vee \theta(y)$. Since μ is an L -fuzzy ideal of R , we can write

$$\mu(ar) \geq \mu(a) \vee \mu(r) \geq \mu(r) \geq \mu(y) \vee \theta(y).$$

Hence there exist elements $aq, ar \in R$ such that $ax = yaq + ar$ where either $ar = 0$ or else $\mu(ar) \geq \mu(y) \vee \theta(y)$. Therefore we get $ax \in \mu_{\theta_y}$. This means that μ_{θ_y} is an ideal of R . We can write $y = y.1 + 0$ where $q = 1, r = 0 \in R$. Then $0 \neq y \in \mu_{\theta_y}$. Thus $\mu_{\theta_y} \neq \{0\}$. \square

Definition 3.4 *Let μ be any L -fuzzy ideal of a ring R . The ideals μ_{θ_y} are called θ -Euclidean level ideals of μ .*

Corollary 3.5 *Let $\mu: R \rightarrow L$ be an L -fuzzy ideal of R and $\theta: R \rightarrow L$ be a fuzzy set. If $\mu(1) \geq \mu(y) \vee \theta(y)$ for $0 \neq y \in R$, then $\mu_{\theta_y} = R$.*

Proof. μ_{θ_y} is an ideal of R from Theorem 3.3. So $\mu_{\theta_y} \subseteq R$. To obtain $R \subseteq \mu_{\theta_y}$, we have to show that $1 \in \mu_{\theta_y}$. Since R is a ring with identity, we can write $1 = y.0 + 1$ where $r = 1$ and $q = 0$.

There exist elements $0 = q, 1 = r \in R$ such that $1 = y.0 + 1$ where $\mu(r) = \mu(1) \geq \mu(y) \vee \theta(y)$. So $1 \in \mu_{\theta_y}$. Since $1 \in \mu_{\theta_y}$ and μ_{θ_y} is an ideal of R , we obtain that $R \subseteq \mu_{\theta_y}$. Thus $R = \mu_{\theta_y}$. \square

Definition 3.6 Let $\mu: R \rightarrow L$ and $\theta: R \rightarrow L$ be fuzzy sets. For $0 \neq y \in R$ we define $\mu_{\theta y}^*$ as follows:

$$\mu_{\theta y}^* = \{x \in R \mid \text{there exist elements } q, r \in R \text{ such that } x = yq + r \text{ where}$$

$$\theta(r) \geq \mu(y) \vee \theta(y)\}.$$

Theorem 3.7 Suppose that L is a chain and that $\mu: R \rightarrow L$ and $\theta: R \rightarrow L$ are fuzzy sets. Then $\mu: R \rightarrow L$ is a θ -Euclidean L -fuzzy ideal of R if and only if μ_α is an ideal of R for $\alpha \leq \mu(0)$ and $R = \mu_{\theta y} \cup \mu_{\theta y}^*$ for all $0 \neq y \in R$.

Proof. Suppose first that $\mu: R \rightarrow L$ is a θ -Euclidean L -fuzzy ideal of R . Then $\mu: R \rightarrow L$ is an L -fuzzy ideal of R . Because of Proposition 2.3, μ_α is an ideal of R . Now it must be shown that $R = \mu_{\theta y} \cup \mu_{\theta y}^*$ for all $0 \neq y \in R$. Let $0 \neq y \in R$ and $x \in \mu_{\theta y} \cup \mu_{\theta y}^*$. Then $x \in R$. So $\mu_{\theta y} \cup \mu_{\theta y}^* \subseteq R$.

Let $a \in R$. Since $\mu: R \rightarrow L$ is a θ -Euclidean L -fuzzy ideal of R , for $0 \neq y, a \in R$ there exist elements $q, r \in R$ such that $a = yq + r$ where either $r = 0$ or else $\mu(r) \vee \theta(r) \geq \mu(y) \vee \theta(y)$. If $\mu(r) \vee \theta(r) = \mu(r)$, then we get $\mu(r) \geq \mu(y) \vee \theta(y)$. So $a \in \mu_{\theta y}$.

If $\mu(r) \vee \theta(r) = \theta(r)$, then $a \in \mu_{\theta y}^*$. So $a \in \mu_{\theta y} \cup \mu_{\theta y}^*$. Thus $R \subseteq \mu_{\theta y} \cup \mu_{\theta y}^*$. We obtain that $R = \mu_{\theta y} \cup \mu_{\theta y}^*$.

Conversely, suppose that μ_α is an ideal of R for $\alpha \leq \mu(0)$ and $R = \mu_{\theta y} \cup \mu_{\theta y}^*$ for all $0 \neq y \in R$. Because of Proposition 2.3, $\mu: R \rightarrow L$ is an L -fuzzy ideal.

Let $0 \neq y, x \in R$. Since $\mu_{\theta y} \cup \mu_{\theta y}^* = R$, $x \in \mu_{\theta y} \cup \mu_{\theta y}^*$. If $x \in \mu_{\theta y}$, then there exist elements $q, r \in R$ such that $x = yq + r$ where either $r = 0$ or else $\mu(r) \geq \mu(y) \vee \theta(y)$. We can write $\mu(r) \vee \theta(r) \geq \mu(r) \geq \mu(y) \vee \theta(y)$. If $x \in \mu_{\theta y}^*$, then there exist elements $q_1, r_1 \in R$ such that $x = yq_1 + r_1$ where $\theta(r_1) \geq \mu(y) \vee \theta(y)$. Also $\mu(r_1) \vee \theta(r_1) \geq \theta(r_1) \geq \mu(y) \vee \theta(y)$. Finally $\mu: R \rightarrow L$ is a θ -Euclidean L -fuzzy ideal of R . \square

Properties of $\mu_{\theta y}$

(i) Let $\mu: R \rightarrow L$ and $\theta: R \rightarrow L$ be fuzzy sets. For $0 \neq y \in R$,

(a) $\mu_\alpha \subseteq \mu_{\theta y}$ for all $\alpha \in L$ such that $\alpha \geq \mu(y) \vee \theta(y)$;

(b) $(y) \subseteq \mu_{\theta y}$;

(c) $\mu_{\theta_1} = R$ (for $y = 1$).

(ii) Let L be a chain, $\mu: R \rightarrow L$ be an L -fuzzy ideal, $\theta: R \rightarrow L$ be a fuzzy set and $0 \neq y \in R$. If $x, y \in \mu_{\theta_y} \cap \mu_{\theta_x}$ and $\theta(x) = \theta(y)$, then $\mu_{\theta_x} = \mu_{\theta_y}$.

Proof.

(a) Let $x \in \mu_\alpha$. Then $x \in R$ and $\mu(x) \geq \alpha$. We can write $x = y.0 + x$ where $q = 0, r = x \in R$. Also $\mu(r) = \mu(x) \geq \alpha \geq \mu(y) \vee \theta(y)$. Hence $x \in \mu_{\theta_y}$. So $\mu_\alpha \subseteq \mu_{\theta_y}$.

(b) We know that $y \in \mu_{\theta_y}$. So $(y) \subseteq \mu_{\theta_y}$.

(c) Let $a \in R$. We can write $a = 1.a + 0$ where $r = 0, q = a \in R$. Thus $a \in \mu_{\theta_1}$ and $R \subseteq \mu_{\theta_1}$. Also $\mu_{\theta_1} \subseteq R$. So $R = \mu_{\theta_1}$.

(ii) Since $x \in \mu_{\theta_y}$, there exist elements $q, r \in R$ such that $x = yq + r$ where either $r = 0$ or else $\mu(r) \geq \mu(y) \vee \theta(y)$.

Let $t \in \mu_{\theta_x}$. Then there exist elements $q_1, r_1 \in R$ such that $t = xq_1 + r_1$ where either $r_1 = 0$ or else $\mu(r_1) \geq \mu(x) \vee \theta(x)$. We can write

$$t = (yq + r)q_1 + r_1 = yqq_1 + rq_1 + r_1$$

and $qq_1, rq_1 + r_1 \in R$.

If $r = r_1 = 0$, then $rq_1 + r_1 = 0$.

If $r = 0$ and $r_1 \neq 0$, then $\mu(rq_1 + r_1) = \mu(r_1) \geq \mu(x) \vee \theta(x) = \mu(yq + r) \vee \theta(y)$
 $= \mu(yq) \vee \theta(y) \geq \mu(y) \vee \theta(y)$.

If $r \neq 0$ and $r_1 = 0$, then $\mu(rq_1 + r_1) = \mu(rq_1) \geq \mu(r) \geq \mu(y) \vee \theta(y)$.

Suppose that $r \neq 0$ and $r_1 \neq 0$. Then $\mu(rq_1 + r_1) \geq \mu(rq_1) \wedge \mu(r_1) \geq \mu(r) \wedge \mu(r_1)$.

If $r \neq 0, r_1 \neq 0$ and $\mu(r) \wedge \mu(r_1) = \mu(r)$, then $\mu(rq_1 + r_1) \geq \mu(r) \geq \mu(y) \vee \theta(y)$.

If $r \neq 0, r_1 \neq 0$ and $\mu(r) \wedge \mu(r_1) = \mu(r_1)$, then

$$\begin{aligned} \mu(rq_1 + r_1) &\geq \mu(r_1) \geq \mu(x) \vee \theta(x) = \mu(yq + r) \vee \theta(y) \\ &\geq \{\mu(y) \wedge \mu(r)\} \vee \theta(y) = \mu(y) \vee \theta(y). \end{aligned}$$

Finally we can say that there exist elements $qq_1, rq_1 + r_1 \in R$ such that $t = y(qq_1) + (rq_1 + r_1)$ where either $rq_1 + r_1 = 0$ or else

$$\mu(rq_1 + r_1) \geq \mu(y) \vee \theta(y).$$

This means that $\mu_{\theta_x} \subseteq \mu_{\theta_y}$.

We obtain $\mu_{\theta_y} \subseteq \mu_{\theta_x}$ in a similar way. So $\mu_{\theta_y} = \mu_{\theta_x}$. □

Acknowledgement

The authors would like to thanks the referees for their valuable suggestions.

References

- [1] A. Koç and E. Balkanay, θ -Euclidean L -fuzzy Ideals of Rings, Turkish Journal of Mathematics 26 (2002) 149-158.
- [2] H.V. Kumbhojkar and M.S. Bapat, Correspondence theorem for fuzzy ideals, Fuzzy Sets and Systems 41 (1991) 213-219.
- [3] H.V. Kumbhojkar and M.S. Bapat, On semiprime fuzzy ideals, Fuzzy Sets and Systems 60 (1993) 219-223.
- [4] R.Kumar, Fuzzy cosets and some fuzzy radicals, Fuzzy Sets and Systems 46 (1992) 261-265.
- [5] R.Kumar, Fuzzy prime spectrum of a ring, Fuzzy Sets and Systems 46 (1992) 147-154.

Ayten KOÇ, Erol BALKANAY
İstanbul Kültür University
Department of Mathematics Computer,
Şirinevler 34510 İstanbul-TURKEY
e-mail: akoc@iku.edu.tr
e-mail: e.balkanay@iku.edu.tr

Received 26.12.2002